

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
13 February 2003 (13.02.2003)

PCT

(10) International Publication Number
WO 03/011842 A1(51) International Patent Classification⁷: **C07D 263/32**,
A61K 31/421, C07C 69/734, C07D 413/04, 277/24,
417/04, 413/10, A61P 3/04Street, Stratford, CT 06615 (US). **LIVINGSTON, James**,
N. [US/US]; 170 Dromara Road, Guilford, CT 06437
(US).

(21) International Application Number: PCT/US02/23614

(74) Agents: **GREENMAN, Jeffrey, M.** et al.; BAYER COR-
PORATION, 400 Morgan Lane, West Haven, CT 06516
(US).

(22) International Filing Date: 25 July 2002 (25.07.2002)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
60/308,500 27 July 2001 (27.07.2001) US
60/373,048 16 April 2002 (16.04.2002) US(71) Applicant (*for all designated States except US*): **BAYER**
CORPORATION [US/US]; 100 Bayer Road, Pittsburgh,
PA 15205 (US).

(72) Inventors; and




(75) Inventors/Applicants (*for US only*): **LOWE, Derek**,
B. [US/US]; 31 Bittersweet Lane, Hamden, CT 06518
(US). **WICKENS, Philip, L.** [CA/US]; 9 Michaels Way,
Wallingford, CT 06492 (US). **MA, Xin** [CN/US]; 46 Hill-
top Road, Bethany, CT 06524 (US). **ZHANG, Mingbao**
[CN/US]; 75 Nyselius Place, Stamford, CT 06905 (US).
BULLOCK, William, H. [US/US]; 60 Hermann Lane,
Easton, CT 06612 (US). **COISH, Philip, D.G.** [CA/US];
548 Orange Street, Suite 401, New Haven, CT 06511
(US). **MUGGE, Ingo, A.** [DE/US]; 88 Vista Terrace, New
Haven, CT 06515 (US). **STOLLE, Andreas** [DE/DE]; Am
Kirchenfeld 13, 52327 Wuppertal (DE). **WANG, Ming**
[US/US]; 32 Milford Hunt Lane, Milford, CT 06460 (US).
WANG, Yamin [CN/US]; 10 Russet Road, Sandy Hook,
CT 06482 (US). **ZHANG, Chengzhi** [CA/US]; 193 Mul-
berry Lane, Orange, CT 06477 (US). **ZHANG, Hai-Jun**
[CN/US]; 260 Pine Street, Middletown, CT 06457 (US).
ZHU, Lei [US/US]; 101 Oak Ridge Lane, Milford, CT
06460 (US). **TSUTSUMI, Manami** [JP/US]; 70 Maureen(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU,
SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG,
UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS,
MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB,
GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF,
BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,
TD, TG)*

[Continued on next page]

WO 03/011842 A1

(54) Title: INDANE ACETIC ACID DERIVATIVES AND THEIR USE AS PHARMACEUTICAL AGENTS, INTERMEDIATES,
AND METHOD OF PREPARATION(57) Abstract: This invention relates to novel indane acetic acid derivatives of formula (I) which are useful in the treatment of
diseases such as diabetes, obesity, hyperlipidemia, and atherosclerotic diseases. The invention also relates to intermediates of formula
(II) and (III) of indane acetic derivatives and to methods of preparation.



- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- with international search report

**INDANE ACETIC ACID DERIVATIVES AND THEIR USE AS PHARMACEUTICAL
AGENTS, INTERMEDIATES, AND METHOD OF PREPARATION**

This application claims benefit of U.S. Provisional Application Serial No. 60/308,500, filed July 27, 2001, and U.S. Provisional Application Serial No. 60/373,048, filed on April 16, 2002, the contents of which are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

This invention is directed to indane acetic acid derivatives and their use in pharmaceutical compositions for the treatment of diseases such as diabetes, obesity, hyperlipidemia, and atherosclerotic disease. The invention is also directed to intermediates useful in preparation of indane acetic derivatives and to methods of preparation.

BACKGROUND OF THE INVENTION

Type II diabetes is the more common form of diabetes, with 90-95% of hyperglycemic patients experiencing this form of the disease. In Type II diabetes, there appears to be a reduction in the pancreatic β -cell mass, several distinct defects in insulin secretion, and a decrease in tissue sensitivity to insulin. The symptoms and consequences of this form of diabetes include fatigue, frequent urination, thirst, blurred vision, frequent infections and slow healing of sores, diabetic nerve damage, retinopathy, micro and macro blood vessel damage, and heart and renal disease.

Resistance to the metabolic actions of insulin is one of the key features of Type II diabetes. Insulin resistance is characterized by impaired uptake and utilization of glucose in insulin-sensitive target organs, for example, adipocytes and skeletal muscle, and by impaired inhibition of hepatic glucose output. Functional insulin deficiency, insulin resistance in the periphery, and the failure of insulin to suppress hepatic glucose output results in fasting hyperglycemia. Pancreatic β -cells compensate for the insulin resistance by secreting increased levels of insulin. However, the β -cells are unable to maintain this high output of insulin, and eventually, the glucose-induced insulin secretion falls, leading to the deterioration of glucose homeostasis and to the subsequent development of overt diabetes. Hyperinsulinemia is also linked to insulin resistance, hypertriglyceridemia, low high-density lipoprotein (HDL) cholesterol, and increased plasma concentration of low-density lipoproteins (LDL). The association of insulin resistance and hyperinsulinemia

with these metabolic disorders has been termed "Syndrome X," and has been strongly linked to an increased risk of hypertension and coronary artery disease.

Obesity is an excessive accumulation of adipose tissue. Excess adipose tissue is associated with the development of serious medical conditions, for example, Type II diabetes, hypertension, coronary artery disease, hyperlipidemia, obesity, and certain malignancies. The adipocyte may also influence glucose homeostasis through the production of tumor necrosis factor α (TNF α) and other molecules.


Atherosclerotic disease is known to be caused by a number of factors, for example, hypertension, diabetes, low levels of HDL, and high levels of LDL. Atherosclerotic-related diseases include cardiovascular disease, coronary heart disease (CHD), cerebrovascular disease, and peripheral vessel disease. Coronary heart disease includes CHD death, myocardial infarction, and coronary revascularization. Cerebrovascular disease includes ischemic or hemorrhagic stroke, and transient ischemic attacks.

Accordingly, despite the presence of some pharmaceuticals that are used to treat these diseases, there remains a need for new pharmaceuticals that are both safe and effective agents for the treatment of disease, and for useful methods to prepare them.

The present invention relates to compounds which are useful in the treatment of diabetes and related disorders such as Syndrome X, impaired glucose tolerance, impaired fasting glucose, and hyperinsulinemia; obesity; atherosclerotic disease, dyslipidemia, and related disorders such as hypertriglyceridemia, low HDL cholesterol, and hypercholesterolemia; cardiovascular disease; and cerebrovascular disease.

DESCRIPTION OF THE INVENTION

The present invention encompasses the compounds of Formula I,



wherein

R is H or C₁ - C₆ alkyl;

R¹ is H, COOR, C₃-C₈ cycloalkyl, or

C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;

R² is H, halo, or C₁-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro,

or

R² is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl, each of which may be unsubstituted or substituted with R⁶;

R³ is H, C₁-C₆ alkyl, or phenyl which may be unsubstituted or substituted with R⁶;

X is O or S;

R⁴ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl, either of which may be unsubstituted or substituted with fluoro, oxo, or C₁-C₆ alkoxy which may be unsubstituted or substituted with C₁-C₆ alkoxy, or phenyl optionally substituted with R⁶,

or either of which may be substituted with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶,

or

C₁-C₆ alkyl may also be substituted with C₃-C₈ cycloalkyl or with phenoxy which may be unsubstituted or substituted with R⁶ or with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶,

or

R⁴ is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, benzodioxolyl, dihydrobenzofuranyl, indolyl, pyrimidinyl or phenoxy,

each of which may be unsubstituted or substituted with R⁶;

R⁵ is H, halo or C₁-C₆ alkyl optionally substituted with oxo;

R⁶ is halo, CF₃, C₁-C₆ alkyl optionally substituted with oxo or hydroxy, or

C₁-C₆ alkoxy optionally substituted with fluoro;

and pharmaceutically acceptable salts and esters thereof.

The terms identified above have the following meaning throughout:

"C₁-C₆ alkyl" means straight or branched chain alkyl groups having from one to about six carbon atoms. Such groups include methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, *tert*-butyl, *n*-pentyl, *neo*-pentyl, 2-pentyl, *n*-hexyl, 2-hexyl, 3-hexyl, 2,3-dimethylbutyl, and the like.

"C₂-C₆ alkenyl" means straight or branched chain alkenyl groups having from two to about six carbon atoms and containing one or more double bonds. Such groups include ethenyl, propenyl, isopropenyl, 2-isobutenyl, 4-pentenyl, 5-hexenyl, and the like.

"C₃-C₈ cycloalkyl" means saturated monocyclic alkyl groups having from 3 to about 8 carbon atoms and includes such groups as cyclopropyl, cyclopentyl, cyclohexyl, and the like.

"C₁-C₆ alkoxy" means straight or branched chain alkoxy groups having from one to about six carbon atoms and includes such groups as methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, isobutoxy, *sec*-butoxy, *tert*-butoxy, and the like.

"Halo" means fluoro, chloro, bromo, or iodo.

When an alkyl, cycloalkyl, alkenyl, or alkoxy group is described as being substituted with fluoro, it may be substituted with one or more fluorine atoms at any available carbon atom up to the perfluoro level.

When an alkyl substituent is described as being substituted by oxo, it means substitution by a doubly bonded oxygen atom, which forms together with the carbon to which it is attached, a carbonyl group $-(C=O)-$.


When any moiety is described as being substituted, it can have one or more of the indicated substituents that may be located at any available position on the moiety. When there are two or more substituents on any moiety, each substituent may be defined independently of any other substituent and may, accordingly, be the same or different.

The term "optionally substituted" means that the moiety so modified may be unsubstituted or substituted with the identified substituent(s).




R^3 may be attached to the heterocyclic moiety of the compound of Formula I at either the 4 or 5 position (i.e., at either available carbon atom) and, accordingly, the remaining portion of the molecule will be attached at the remaining available carbon atom.



Examples of the compounds of Formula I, which are illustrative of the present invention but not limiting in any way, are listed in Table 1.


Table 1. Illustrative Examples of Compounds of Formula I


















Entry No.	R^1	R^2	R^3	R^4	R^5	X
1	H	H	H	CH ₃	H	O
2	H	H	H	n-butyl	H	O
3	H	H	H	cyclopropyl	H	O
4	H	H	H	cyclopentyl	H	O
5	H	H	H	cyclooctyl	H	O
6	H	H	H	Ph	H	O
7	H	H	H	Ph	H	S
8	H	H	H	2-Cl Ph	H	O






Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
9	H	H	H	2,3-d-F Ph	H	O
10	H	H	H	2,4-di-CH ₃ Ph	H	O
11	H	H	H	2-thienyl	H	O
12	H	H	H		H	O
13	H	H	H	2-furyl	H	O
14	H	H	H	2-furyl	H	S
15	H	H	H	2-(4-CH ₃)furyl	H	O
16	H	H	H		H	O
17	H	H	H	4-F Ph	H	O
18	H	H	H	4-F Ph	H	S
19	H	H	CH ₃	4-F Ph	H	O
20	H	H	Et	4-F Ph	H	O
21	H	H	Et	4-F Ph	H	S
22	H	H	Et	3-pyridyl	H	O
23	H	H	Et		H	O
24	H	H	isopropyl	4-F Ph	H	O
25	H	H	isopropyl	2,4-di-F Ph	H	O
26	H	H	n-butyl	2,4-di-F Ph	H	O
27	H	H	n-hexyl	2,4-di-F Ph	H	O
28	H	H	Ph	2,4-di-F Ph	H	O
29	H	H	4-F Ph	2,4-di-F Ph	H	O
30	H	CH ₃	Et	Ph	H	O
31	H	CH ₃	Et	Ph	H	S
32	H	CH ₃	Et	3-CH ₃ O Ph	H	O
33	H	H	Et	3-CH ₃ O Ph	H	O










Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
34	H	H	Et	3-CH ₃ O Ph	H	S
35	H	H	Et	4-CH ₃ O Ph	H	O
36	H	H	Et	4-CH ₃ O Ph	H	S
37	H	H	Et	4-EtO Ph	H	S
38	H	H	Et	4-EtO Ph	H	O
39	H	H	Me		H	O
40	H	H	Me	PhCH ₂	H	O
41	H	H	Me	3-Cl-4-F-Ph	H	O
42	H	H	Me	3-F-4-Me-Ph	H	O
43	H	H	Me	3-Me-4-F-Ph	H	O
44	H	H	Me	3-NH ₂ -4-Me-Ph	H	O
45	H	H	Et	4-Et-Ph	H	O
46	H	H	Me	4-Et-Ph	H	O
47	H	H	Et	4-CN-Ph	H	O
48	H	H	Et	4-(Et) ₂ N-Ph	H	O
49	H	H	Me	4- <i>i</i> -Pr-Ph	H	O
50	H	H	Me	4- <i>t</i> -Bu-Ph	H	O
51	H	H	Me	4-Et-Ph	H	O
52	H	H	Me	4- <i>n</i> -Bu-Ph	H	O
53	H	H	Et	4- <i>n</i> -Pr-Ph	H	O
54	H	CH ₃	Et	4-CH ₃ O Ph	H	O
55	H	CH ₃	Et	4-CH ₃ O Ph	H	S
56	H	CH ₃	Et	4-CH ₃ O Ph	CH ₃	O
57	H	CH ₃	Et	3,4-di-CH ₃ O Ph	CH ₃	O
58	H	CH ₃	Et	4-Ph Ph	CH ₃	O
59	H	CH ₃	Et	4-Ph Ph	CH ₃	S
60	H	CH ₃	Et		CH ₃	O
61	H	CH ₃	Ph	cyclopropyl	H	O
62	H	CH ₃	Ph	cyclohexyl	H	O
63	H	CH ₃	Ph	cyclohexyl	H	S








Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
64	H	CH ₃	p-F Ph	cyclohexyl	H	O
65	H	Cl	i-Pr	Ph	H	O
66	H	Cl	i-Pr	Ph	H	S
67	H	Cl	i-Pr	Ph	Cl	O
68	H	Cl	i-Pr	4-CH ₃ Ph	Cl	O
69	H	Br	CH ₃	Ph	Br	O
70	H	Br	CH ₃	3-F Ph	Br	O
71	H	Br	CH ₃	3-F Ph	Br	S
72	H	CH ₃ CO	CH ₃	n-propyl	CH ₃ CO	O
73	H	CH ₂ OCH ₃	Et	2-thienyl	H	O
74	H	Ph	H	2,4-di-Cl Ph	H	O
75	H	Ph	H	2,4-di-Cl Ph	H	S
76	H	Ph	CH ₃	2,4-di-Cl Ph	H	O
77	H	Ph	Et	2,4-di-Cl Ph	H	O
78	H	Ph	Ph	2,4-di-Cl Ph	H	O
79	H	Ph	Ph	2,4-di-Cl Ph	H	S
80	H	Ph	4-CH ₃ O-Ph	2,4-di-Cl Ph	H	O
81	H	4-F Ph	CH ₃	4-F Ph	H	O
82	H	4-F Ph	CH ₃	2,4-di-Cl Ph	H	O
83	H	3-pyridyl	CH ₃	2,4-di-Cl Ph	H	O
84	H	3-pyridyl	CH ₃	2,4-di-Cl Ph	H	S
85	H	2-thienyl	CH ₃	Ph	H	O
86	H	2-thienyl	CH ₃	2,4-di-Cl Ph	H	O
87	H	2-thienyl	CH ₃	2,4-di-Cl Ph	H	S
88	H	2-thienyl	CH ₃	3-pyridyl	H	O
89	H	2-thienyl	CH ₃	cyclopentyl	H	O
90	H	2-thienyl	CH ₃		H	O
91	H	2-thienyl	CH ₃	Ph	2-thienyl	O
92	CH ₃	H	H	Ph	H	O
93	CH ₃	H	H	Ph	H	S

Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
94	CH ₃	H	H	2-thienyl	H	O
95	CH ₃	H	H	2-thienyl	H	S
96	CH ₃	H	H		H	O
97	CH ₃	H	H		H	O
98	CH ₃	H	H		H	O
99	CH ₃	H	H	2-pyridyl	H	O
100	CH ₃	H	H		H	O
101	CH ₃	H	CH ₃	cyclobutyl	H	O
102	CH ₃	H	CH ₃	cyclohexyl	H	O
103	CH ₃	H	CH ₃	cyclohexyl	H	S
104	CH ₃	H	CH ₃	3,4-di-F Ph	H	O
105	CH ₃	H	CH ₃	3,4-di-F Ph	H	S
106	CH ₃	H	CH ₃	2-pyridyl	H	O
107	CH ₃	H	CH ₃		H	O
108	CH ₃	H	CH ₃		H	O
109	CH ₃	H	Et	Ph	H	O
110	CH ₃	H	Et	Ph	H	S
111	CH ₃	H	Et	4-CF ₃ Ph	H	O
112	CH ₃	H	Et		H	O
113	CH ₃	H	Et	2-naphthyl	H	O












Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
114	CH ₃	H	Et		H	O
115	CH ₃	H	Et		H	O
116	CH ₃	H	Et		H	S
117	CH ₃	H	Et		H	O
118	CH ₃	H	Et		H	O
119	CH ₃	H	i-Pr	Ph	H	O
120	CH ₃	H	i-Pr	Ph	H	S
121	CH ₃	H	i-Pr	3,4-di-F Ph	H	O
122	CH ₃	H	i-Pr	3,4-di-Cl Ph	H	O
123	CH ₃	H	i-Pr	4-Ph Ph	H	O
124	CH ₃	H	i-Pr	4-Ph Ph	H	S
125	CH ₃	H	i-Pr	4-(4-ClPh)Ph	H	O
126	CH ₃	H	i-Pr	4-(4-ClPh)Ph	H	S
127	CH ₃	H	i-Pr		H	O
128	CH ₃	H	i-Pr		H	O
129	CH ₃	H	i-Pr		H	O









Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
130	CH ₃	H	i-Pr		H	O
131	CH ₃	H	i-Pr	3-(5-CH ₃) pyridyl	H	O
132	CH ₃	H	i-Pr		H	O
133	CH ₃	H	i-Pr		H	S
134	CH ₃	H	i-Pr		H	O
135	CH ₃	CH ₃	i-Pr	3,4-di-Cl Ph	CH ₃	O
136	CH ₃	n-propyl	i-Pr	3,4-di-Cl Ph	n-propyl	O
137	CH ₃	Cl	i-Pr	4-Cl Ph	H	O
138	CH ₃	Cl	i-Pr	4-Cl Ph	H	S
139	CH ₃	Cl	i-Pr	3-CH ₃ O Ph	H	O
140	CH ₃	Cl	i-Pr	3-CH ₃ O Ph	Cl	O
141	CH ₃	Cl	i-Pr	3-CH ₃ O Ph	Cl	S
142	CH ₃	Cl	i-Pr		Cl	O
143	CH ₃	Br	i-Pr	Ph	H	O
144	CH ₃	Br	i-Pr	3-Cl Ph	H	O
145	CH ₃	Br	i-Pr	Ph	Br	O
146	CH ₃	Br	i-Pr	Ph	Br	S
147	CH ₃	CH ₃	i-Pr	Ph	H	O
148	CH ₃	CH ₃	i-Pr	Ph	H	S
149	CH ₃	CH ₃	i-Pr	2-Cl Ph	H	O



Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
150	CH ₃	CH ₃	i-Pr		H	O
151	CH ₃	CH ₃ CO	i-Pr	3-F Ph	H	O
152	CH ₃	CH ₃ CO	i-Pr	3-F Ph	H	S
153	CH ₃	n-PrCO	i-Pr	3-F Ph	H	O
154	CH ₃	n-BuCO	i-Pr	3-F Ph	H	O
155	CH ₃	H	n-Bu	Ph	H	O
156	CH ₃	H	n-Bu		H	O
157	CH ₃	H	n-Bu		H	S
158	CH ₃	H	n-Bu	2-Cl Ph	H	O
159	CH ₃	H	n-Bu	2,4 di-F Ph	H	O
160	CH ₃	H	n-Bu	3,4 di- CH ₃ O Ph	H	O
161	CH ₃	H	n-Bu		H	O
162	CH ₃	H	n-Bu	2-furyl	H	O
163	CH ₃	H	n-Bu		H	O
164	CH ₃	H	n-Bu		H	O
165	CH ₃	H	n-Bu		H	S
166	CH ₃	H	n-Bu		H	O
167	CH ₃	H	n-Bu		H	S




Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
168	CH ₃	H	n-Bu		H	O
169	CH ₃	H	n-Bu		H	O
170	CH ₃	Br	n-Bu	2,4 di-F Ph	Br	O
171	CH ₃	Cl	n-Bu	2,4 di-F Ph	H	O
172	CH ₃	H	n-pentyl	Ph	H	O
173	CH ₃	H	n-pentyl	2,4 di-F Ph	H	O
174	CH ₃	H	n-pentyl	2,4 di-F Ph	H	S
175	CH ₃	H	n-pentyl	4-pyridyl	H	O
176	CH ₃	H	n-pentyl		H	O
177	CH ₃	Cl	n-pentyl	Ph	H	O
178	CH ₃	Cl	n-pentyl	Ph	H	S
179	CH ₃	H	Ph		H	O
180	CH ₃	H	2-Cl Ph		H	O
181	CH ₃	H	2-Cl Ph		H	S
182	CH ₃	H	H	PhOCH ₂	H	O
183	CH ₃	H	H	(4-CH ₃ Ph)OCH ₂	H	O
184	CH ₃	H	H		H	O
185	CH ₃	H	CH ₃	Et	H	O
186	CH ₃	H	CH ₃	Et	H	S
187	CH ₃	H	CH ₃	CF ₃ CF ₂	H	O
188	CH ₃	H	CH ₃	t-butyl	H	O






Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
189	CH ₃	H	Et	3-(5-CH ₃) pyridyl	H	O
190	CH ₃	H	Et	4-pyridyl	H	O
191	CH ₃	H	Et	4-pyridyl	H	S
192	CH ₃	Et	CH ₃	PhOCH ₂	H	O
193	CH ₃	Et	CH ₃	PhOCH ₂	H	S
194	CH ₃	Et	CH ₃	PhCH ₂ OCH ₂	H	O
195	CH ₃	n-propyl	CH ₃	PhOCH ₂	H	O
196	CH ₃	n-propyl	CH ₃	PhOCH ₂	n-propyl	O
197	CH ₃	n-butyl	CH ₃	PhOCH ₂	H	O
198	CH ₃	n-hexyl	CH ₃	PhOCH ₂	H	O
199	CH ₃	n-hexyl	CH ₃	PhOCH ₂	H	S
200	CH ₃	n-hexyl	isopropyl	3-Cl Ph	H	O
201	CH ₃	n-hexyl	Ph	3-Cl Ph	H	O
202	CH ₃	CH ₃ OCH ₂	CH ₃	PhOCH ₂	H	O
203	CH ₃	Ph	n-butyl	3,4-di-F Ph	H	O
204	CH ₃	3-F Ph	CH ₃	1-naphthyl	H	O
205	CH ₃	4-pyridyl	H	4-CF ₃ Ph	H	O
206	CH ₃	4-pyridyl	H	4-CF ₃ Ph	H	S
207	CH ₃	Cl	CH ₃	3,5-di-F-Ph	H	O
208	CH ₃	Br	CH ₃	CF ₃ CF ₂	H	O
209	CH ₃	Br	n-butyl	CF ₃ CF ₂	H	O
210	CH ₃	Br	n-butyl	CF ₃ CF ₂	Br	O
211	CH ₃	Br	Ph	CF ₃ CF ₂	Br	O
212	CH ₃	2-furyl	CH ₃	isobutyl	H	O
213	CH ₃	2-furyl	CH ₃	isobutyl	H	S
214	CH ₃	2-furyl	CH ₃	2-F-4-CF ₃ Ph	H	O
215	CH ₃	2-furyl	CH ₃	2-naphthyl	H	O
216	CH ₃	2-furyl	i-Pr	isobutyl	H	O
217	CH ₃	EtCO	n-propyl	3-CH ₃ O Ph	EtCO	O
218	Et	H	H	cyclopropyl	H	O
219	Et	H	H	4-F Ph	H	O
220	Et	H	H	3,5-di-F-Ph	H	O
221	Et	H	H	4-Cl PhCH ₂	H	O







Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
222	Et	H	H	2-quinolinyl	H	O
223	Et	H	CH ₃	PhCH ₂	H	O
224	Et	H	CH ₃	4-F PhCH ₂	H	O
225	Et	H	CH ₃	3,4-di-F-PhOCH ₂	H	O
226	Et	H	CH ₃		H	O
227	Et	H	CH ₃		H	S
228	Et	H	CH ₃		H	O
229	Et	H	CH ₃		H	O
230	Et	H	CH ₃		H	S
231	Et	H	CH ₃		H	O
232	Et	H	CH ₃		H	O
233	Et	H	CH ₃		H	S
234	Et	H	CH ₃	2-quinolinyl	H	O
235	Et	H	CH ₃		H	O
236	Et	H	CH ₃		H	O
237	Et	H	CH ₃		H	O











Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
238	Et	H	CH ₃		H	O
239	Et	H	CH ₃		H	O
240	Et	H	CH ₃		H	O
241	Et	H	CH ₃		H	O
242	Et	H	CH ₃		H	O
243	Et	H	CH ₃	(4-CH ₃ O) PhCH ₂ CH ₂	H	O
244	Et	H	CH ₃		H	O
245	Et	Cl	CH ₃		H	O
246	Et	Br	CH ₃		H	O
247	Et	H	Et	4-Ph Ph	H	O
248	Et	H	Et	4-Ph Ph	H	S
249	Et	H	Et	4-(4-CH ₃ Ph)Ph	H	O
250	Et	CH ₃	CH ₃	2-F Ph	H	O
251	Et	CH ₃	CH ₃	2-F Ph	CH ₃	O
252	Et	CH ₃	CH ₃	2-F Ph	CH ₃	O
253	Et	CH ₃	CH ₃	2-F Ph	CH ₃	S
254	Et	3-Cl Ph	Et	4-Ph Ph	H	O
255	Et	3-Cl Ph	Et	4-Ph Ph	H	S








Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
256	Et	CH ₃ CO	H	4-F Ph	H	O
257	Et	CH ₃ CO	isopropyl	4-F Ph	H	O
258	Et	CH ₃ CO	Ph	4-F Ph	H	O
259	Et	CH ₃ CO	CH ₃	cyclohexyl	CH ₃ CO	O
260	Et	CH ₃ CO	CH ₃	4-F Ph	CH ₃ CO	O
261	Et	CH ₃ CO	Ph	4-F Ph	CH ₃ CO	O
262	Et	CH ₃ CO	Ph	4-F Ph	CH ₃ CO	S
263	Et	Cl	Et	4-(4-CH ₃ Ph)Ph	H	O
264	Et	Cl	Et	4-(4-CH ₃ Ph)Ph	Cl	O
265	Et	Cl	Et		Cl	O
266	Et	Br	Ph	2-OCH ₃ Ph	Br	O
267	CF ₃ CH ₂	H	H	n-butyl	H	O
268	CF ₃ CH ₂	H	H	Ph	H	O
269	CF ₃ CH ₂	H	H	3-pyridyl	H	O
270	CF ₃ CH ₂	H	CH ₃	cyclopentyl	H	O
271	CF ₃ CH ₂	H	CH ₃	4-(CF ₃ O)Ph	H	O
272	CF ₃ CH ₂	H	CH ₃	4-(CF ₃ O)Ph	H	S
273	CF ₃ CH ₂	H	CH ₃	4-(CHF ₂ O)Ph	H	O
274	CF ₃ CH ₂	H	CH ₃		H	O
275	CF ₃ CH ₂	H	n-butyl	(4-F Ph)OCH ₂	H	O
276	CF ₃ CH ₂	H	Ph	Ph	H	O
277	CF ₃ CH ₂	H	Ph	Ph	H	S
278	CF ₃ CH ₂	H	Ph	2-(5-CF ₃) furyl	H	O
279	CF ₃ CH ₂	H	Ph	2-thienyl	H	O
280	CF ₃ CH ₂	H	4-F Ph	Ph	H	O
281	CF ₃ CH ₂	CH ₃	H	2-F Ph	H	O
282	CF ₃ CH ₂	CH ₃	H	2-F Ph	H	S
283	CF ₃ CH ₂	CH ₃	H	2-F Ph	CH ₃	O
284	CF ₃ CH ₂	CH ₃	Et	3-CF ₃ Ph	H	O








Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
285	CF ₃ CH ₂	CH ₃	n-butyl	(4-F Ph)OCH ₂	H	O
286	CF ₃ CH ₂	CH ₃	n-butyl	(4-F Ph)OCH ₂	H	S
287	CF ₃ CH ₂	CH ₃	Ph	2-thienyl	H	O
288	n-propyl	H	H	CH ₃	H	O
289	n-propyl	H	H	CH ₃	H	S
290	n-propyl	H	H	n-propyl	H	O
291	n-propyl	H	H	cyclobutyl	H	O
292	n-propyl	H	H	cycloheptyl	H	O
293	n-propyl	H	H	3,4-di-CH ₃ Ph	H	O
294	n-propyl	H	H	2-thienyl	H	O
295	n-propyl	H	H	2-thienyl	H	S
296	n-propyl	H	H		H	O
297	n-propyl	H	CH ₃	CH ₃	H	O
298	n-propyl	H	CH ₃	CH ₃	H	S
299	n-propyl	H	CH ₃	3-CF ₃ Ph	H	O
300	n-propyl	H	CH ₃	2-thienyl	H	O
301	n-propyl	H	CH ₃	3-(4-(OCH ₃)thienyl)	H	O
302	n-propyl	H	CH ₃	2-(5-(CH ₃)thienyl)	H	O
303	n-propyl	H	CH ₃		H	O
304	n-propyl	H	CH ₃		H	O
305	n-propyl	CH ₃	CH ₃	3-Br Ph	H	O
306	n-propyl	CH ₃	CH ₃	3-Br Ph	H	S
307	n-propyl	CH ₃	CH ₃	3-Br Ph	CH ₃	O




Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
308	n-propyl	CH ₃	CH ₃		H	O
309	n-propyl	CH ₃	CH ₃		H	O
310	n-propyl	n-propyl	CH ₃	3-Cl Ph	H	O
311	n-propyl	n-propyl	CH ₃	3-Cl Ph	H	S
312	n-propyl	CH ₃ OCH ₂	CH ₃	3-Cl Ph	H	O
313	n-propyl	CH ₃ CO	CH ₃	3-Cl Ph	H	O
314	n-propyl	PrCO	CH ₃	3-Cl Ph	H	O
315	n-propyl	PrCO	CH ₃	3-Cl Ph	PrCO	O
316	n-propyl	Cl	CH ₃		H	O
317	n-propyl	Cl	CH ₃		H	O
318	n-propyl	Cl	CH ₃		H	O
319	n-propyl	Cl	H	Ph	Cl	O
320	n-propyl	Cl	CH ₃	Ph	Cl	O
321	n-propyl	Cl	CH ₃	Ph	Cl	S
322	n-propyl	Cl	n-propyl	3-CH ₃ O Ph	Cl	O
323	n-propyl	Cl	n-propyl	3-pyridyl	Cl	O




Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
324	isopropyl	H	H	Ph	H	O
325	isopropyl	H	H	2-quinolinyl	H	O
326	isopropyl	H	H		H	O
327	isopropyl	H	CH ₃	CH ₃	H	O
328	isopropyl	H	CH ₃	t-butyl	H	O
329	isopropyl	H	CH ₃	n-heptyl	H	O
330	isopropyl	H	CH ₃	n-heptyl	H	S
331	isopropyl	H	CH ₃	2,4-di-F Ph	H	O
332	isopropyl	H	CH ₃	2,4-di-F Ph	H	S
333	isopropyl	H	CH ₃	2-F-4-CF ₃ Ph	H	O
334	isopropyl	H	n-propyl	2-F-4-CF ₃ Ph	H	O
335	isopropyl	H	n-propyl	3, 5-di-Cl Ph	H	O
336	isopropyl	H	Ph	2,4-di-CF ₃ Ph	H	O
337	isopropyl	H	4-F Ph	2-F-4-CF ₃ Ph	H	O
338	isopropyl	CH ₃	Et		H	O
339	isopropyl	CH ₃	Et		H	O
340	isopropyl	CH ₃	Et		H	O
341	isopropyl	CH ₃	Et		H	S
342	isopropyl	CH ₃	Et		H	O








Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
343	isopropyl	CH ₃	Et		H	O
344	isopropyl	CH ₃	Et		H	O
345	isopropyl	Et	CH ₃	3-CF ₃ Ph	H	O
346	isopropyl	Et	CH ₃	3-Et Ph	H	O
347	isopropyl	n-propyl	H	PhOCH ₂	H	O
348	isopropyl	n-propyl	H	PhOCH ₂	n-propyl	O
349	isopropyl	n-propyl	H		H	O
350	isopropyl	n-propyl	H		n-propyl	O
351	isopropyl	n-propyl	H		H	S
352	isopropyl	n-propyl	H		H	O
353	isopropyl	n-propyl	n-butyl		H	O
354	isopropyl	n-propyl	Ph		H	O
355	isopropyl	n-butyl	H		H	O
356	isopropyl	n-hexyl	H		H	O
357	isopropyl	Ph	H	CH ₃	H	O
358	isopropyl	Ph	H	n-propyl	H	O
359	isopropyl	Ph	H	n-propyl	H	S





Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
360	isopropyl	Ph	H		H	O
361	isopropyl	Ph	H		H	S
362	isopropyl	Ph	CH ₃		H	O
363	isopropyl	Ph	CH ₃		H	O
364	isopropyl	Cl	Et	Ph	H	O
365	isopropyl	Cl	Et	Ph	H	S
366	isopropyl	Cl	Et	2-CH ₃ Ph	Cl	O
367	isopropyl	Cl	n-propyl	3-F Ph	H	O
368	isopropyl	Cl	isopropyl	3-F Ph	H	O
369	isopropyl	Cl	4-F Ph	3-F Ph	H	O
370	isopropyl	Br	Et	2-CH ₃ Ph	Br	O
371	isopropyl	Br	Et	2-CH ₃ Ph	Br	S
372	n-butyl	H	H	Cyclohexyl	H	O
373	n-butyl	H	H	Ph	H	O
374	n-butyl	H	H	4-F Ph	H	O
375	n-butyl	H	H	3, 5-di-Cl Ph	H	O
376	n-butyl	H	H	3, 5-di-Cl Ph	H	S
377	n-butyl	H	CH ₃	3,4-di-CH ₃ OPh	H	O
378	n-butyl	H	CH ₃	4-F PhOCH ₂	H	O
379	n-butyl	H	CH ₃		H	O
380	n-butyl	H	CH ₃	(4-CH ₃ O) PhCH ₂ CH ₂	H	O
381	n-butyl	H	CH ₃		H	O
382	n-butyl	H	CH ₃		H	S







Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
383	n-butyl	H	Et		H	O
384	n-butyl	H	n-propyl	cyclobutyl	H	O
385	n-butyl	H	n-propyl		H	O
386	n-butyl	H	isopropyl		H	O
387	n-butyl	H	Ph	n-propyl	H	O
388	n-butyl	H	Ph		H	O
389	n-butyl	H	Ph	Ph	H	O
390	n-butyl	H	Ph	Ph	H	S
391	n-butyl	CH ₃	CH ₃	4-CH ₃ Ph	H	O
392	n-butyl	CH ₃	CH ₃	4-CH ₃ Ph	CH ₃	O
393	n-butyl	CH ₃	Et	4-CH ₃ Ph	H	O
394	n-butyl	CH ₃	Ph	4-CH ₃ Ph	H	O
395	n-butyl	CH ₃ OCH ₂	CH ₃	2,4-di-CH ₃ Ph	H	O
396	n-butyl	Cl	CH ₃		H	O
397	n-butyl	Cl	CH ₃		H	O
398	n-butyl	Cl	Ph		H	O
399	n-pentyl	H	H	CH ₃	H	O
400	n-pentyl	H	H	CH ₃	H	S
401	n-pentyl	H	H	Et	H	O
402	n-pentyl	H	H	cyclopentyl	H	O
403	n-pentyl	H	H	cyclopentyl	H	S


Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
404	n-pentyl	H	H	cycloheptyl	H	O
405	n-pentyl	H	H	Ph	H	O
406	n-pentyl	H	H	Ph	H	S
407	n-pentyl	H	H	2-furyl	H	O
408	n-pentyl	H	H	2-(5-CF ₃) furyl	H	O
409	n-pentyl	H	H	2-thienyl	H	O
410	n-pentyl	H	H	3,4-di-Cl Ph	H	O
411	n-pentyl	H	CH ₃	n-butyl	H	O
412	n-pentyl	H	CH ₃	n-butyl	H	S
413	n-pentyl	H	CH ₃		H	O
414	n-pentyl	H	CH ₃	PhOCH ₂	H	O
415	n-pentyl	H	CH ₃	PhCH ₂ OCH ₂	H	O
416	n-pentyl	H	Et	2-F Ph	H	O
417	n-pentyl	H	Et	2-F Ph	H	S
418	n-pentyl	H	4-CH ₃ Ph	2-F Ph	H	O
419	n-pentyl	CH ₃	Et	4-CH ₃ Ph	H	O
420	n-pentyl	Cl	CH ₃	n-butyl	H	O
421	n-pentyl	Cl	CH ₃	Ph	H	O
422	n-pentyl	Cl	CH ₃	Ph	H	S
423	n-pentyl	Cl	CH ₃	4-Ph Ph	H	O
424	n-pentyl	Cl	CH ₃		H	O
425	n-pentyl	Cl	CH ₃		Cl	O
426	n-pentyl	PrCO	CH ₃	4-CH ₃ Ph	PrCO	O
427	n-pentyl	Ph	CH ₃	3-Br Ph	H	O
428	n-pentyl	2-thienyl	CH ₃	3-Br Ph	2-thienyl	O
429	n-hexyl	H	H	2-F Ph	H	O
430	n-hexyl	H	CH ₃	cyclopentyl	H	O
431	n-hexyl	H	CH ₃	cycloheptyl	H	O




Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
432	n-hexyl	H	CH ₃	2-F Ph	H	O
433	n-hexyl	H	CH ₃	2-F Ph	H	S
434	n-hexyl	H	Et	2-F Ph	H	O
435	n-hexyl	H	n-propyl	2-F Ph	H	O
436	n-hexyl	H	isopropyl	2-F Ph	H	O
437	n-hexyl	H	Ph	2-F Ph	H	O
438	n-hexyl	CH ₃ CO	CH ₃	2,4-di-CH ₃ Ph	H	O
439	n-hexyl	CH ₃ OCH ₂	CH ₃	2,4-di-CH ₃ Ph	H	O
440	n-hexyl	Ph	Et	Ph	H	O
441	n-hexyl	Ph	Et	Ph	H	S
442	n-hexyl	Ph	Et	4-pyridyl	H	O
443	n-hexyl	Br	Et	Ph	Br	O
444	n-hexyl	Br	Et	2-F Ph	Br	O
445	cyclopropyl	H	H	cyclopentyl	H	O
446	cyclopropyl	H	H	2, 4-di-Cl Ph	H	O
447	cyclopropyl	H	H		H	O
448	cyclopropyl	H	CH ₃	3-F Ph	H	O
449	cyclopropyl	H	CH ₃	3-F Ph	H	S
450	cyclopropyl	H	CH ₃		H	O
451	cyclopropyl	H	Et		H	O
452	cyclopropyl	H	n-propyl	4-CF ₃ Ph	H	O
453	cyclopropyl	H	isopropyl	Ph	H	O
454	cyclopropyl	H	isopropyl	3-pyridyl	H	O
455	cyclopropyl	H	n-butyl	4-CF ₃ Ph	H	O
456	cyclopropyl	H	n-hexyl	Ph	H	O
457	cyclopropyl	H	n-hexyl	4-CF ₃ Ph	H	O
458	cyclopropyl	H	Ph	Ph	H	O
459	cyclobutyl	H	CH ₃	4-CH ₃ Ph	H	O




Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
460	cyclobutyl	H	Et		H	O
461	cyclobutyl	H	Et		H	O
462	cyclobutyl	H	Et		H	O
463	cyclobutyl	H	Et		H	O
464	cyclobutyl	H	Et		H	O
465	cyclobutyl	H	4-F Ph		H	O
466	cyclobutyl	Cl	CH ₃	3-Cl Ph	Cl	O
467	cyclobutyl	Cl	CH ₃	3-Cl Ph	Cl	S
468	cyclopentyl	H	H	3-CF ₃ Ph	H	O
469	cyclopentyl	H	CH ₃	2,4-di-CF ₃ Ph	H	O
470	cyclopentyl	H	CH ₃	2,4-di-CF ₃ Ph	H	S
471	cyclopentyl	H	n-butyl		H	O
472	cyclopentyl	H	3-F Ph	4-CH ₃ Ph	H	O
473	cyclopentyl	CH ₃	CH ₃	Ph	H	O
474	cyclopentyl	CH ₃	CH ₃	3, 5-di-Cl Ph	H	O
475	cyclopentyl	CH ₃	CH ₃	Ph	H	S
476	cyclopentyl	Et	CH ₃	Ph	H	O
477	cyclopentyl	Cl	CH ₃	Ph	Cl	O
478	cyclopentyl	Cl	CH ₃	Ph	Cl	S
479	cyclohexyl	H	H	3-F Ph	H	O
480	cyclohexyl	H	H	2, 4-di-CH ₃ Ph	H	O





Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
481	cyclohexyl	H	H		H	O
482	cyclohexyl	H	CH ₃	n-propyl	H	O
483	cyclohexyl	H	CH ₃	n-propyl	H	S
484	cyclohexyl	H	CH ₃		H	O
485	cyclohexyl	H	CH ₃	3-Cl Ph	H	O
486	cyclohexyl	H	CH ₃	3-Cl Ph	H	S
487	cyclohexyl	H	CH ₃		H	O
488	cyclohexyl	H	Et		H	O
489	cyclohexyl	H	n-propyl	4-CF ₃ Ph	H	O
490	cyclohexyl	H	n-propyl	3-pyridyl	H	O
491	cyclohexyl	H	isopropyl	Ph	H	O
492	cyclohexyl	H	isopropyl	3-pyridyl	H	O
493	cyclohexyl	H	n-butyl	3-Cl Ph	H	O
494	cyclohexyl	H	n-pentyl	3-Cl Ph	H	O
495	cyclohexyl	H	n-hexyl	4-CF ₃ Ph	H	O
496	cyclohexyl	H	4-F Ph	Ph	H	O
497	cyclohexyl	CH ₃	CH ₃	3-CH ₃ Ph	H	O
498	cyclohexyl	CH ₃	CH ₃	3-CH ₃ Ph	H	S
499	cyclohexyl	CH ₃	Et	3-pyridyl	CH ₃	O
500	cyclohexyl	Et	CH ₃	2-F-4-CF ₃ Ph	Et	O
501	cyclohexyl	2-thienyl	i-Pr	3-pyridyl	H	O
502	cyclohexyl	Cl	CH ₃	2,3-di-CH ₃ Ph	H	O
503	cyclohexyl	Cl	CH ₃	2,3-di-CH ₃ Ph	H	S
504	2-propenyl	H	H	CH ₃	H	O
505	2-propenyl	H	H	isopentyl	H	O
506	2-propenyl	H	H	cyclopentyl	H	O
507	2-propenyl	H	H	Ph	H	O


Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
508	2-propenyl	H	H	Ph	H	S
509	2-propenyl	H	H	2-quinolinyl	H	O
510	2-propenyl	H	H		H	O
511	2-propenyl	H	CH ₃		H	O
512	2-propenyl	H	CH ₃	2,4-di-F Ph	H	O
513	2-propenyl	H	CH ₃	2,4-di-F Ph	H	S
514	2-propenyl	H	CH ₃	2-F-4-CF ₃ Ph	H	O
515	2-propenyl	H	Et	2-naphthyl	H	O
516	2-propenyl	H	Et	2-naphthyl	H	S
517	2-propenyl	H	Et		H	O
518	2-propenyl	H	Et		H	O
519	2-propenyl	H	n-propyl	2-F-4-CF ₃ Ph	H	O
520	2-propenyl	H	Ph	2,4-di-CF ₃ Ph	H	O
521	2-propenyl	H	4-F Ph	2-F-4-CF ₃ Ph	H	O
522	2-propenyl	CH ₃	Et		H	O
523	2-propenyl	Cl	CH ₃	3-CF ₃ Ph	Cl	O
524	2-propenyl	Cl	CH ₃	3-CF ₃ Ph	Cl	S
525	2-propenyl	Br	Et	3-CF ₃ Ph	Br	O
526	2-isobutenyl	H	H	3-pyridyl	H	O
527	2-isobutenyl	H	H		H	O

Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
528	2-isobutenyl	H	CH ₃	4-(CF ₃ O)Ph	H	O
529	2-isobutenyl	H	CH ₃	4-(CF ₃ O)Ph	H	O
530	2-isobutenyl	H	CH ₃	4-(CF ₃ O)Ph	H	S
531	2-isobutenyl	H	n-butyl	4-(CH ₃ O)Ph	H	O
532	2-isobutenyl	H	n-butyl	(4-F Ph)OCH ₂	H	O
533	2-isobutenyl	H	n-butyl	(4-CH ₃ O) PhCH ₂ CH ₂	H	O
534	2-isobutenyl	H	Ph	2-thienyl	H	O
535	2-isobutenyl	H	4-F Ph	Ph	H	O
536	2-isobutenyl	CH ₃ CO	Ph	cyclohexyl	H	O
537	2-isobutenyl	CH ₃ CO	Ph	3-F Ph	H	O
538	4-pentenyl	H	CH ₃	Ph	H	O
539	4-pentenyl	H	CH ₃	Ph	H	S
540	5-hexenyl	H	H	Ph	H	O
541	5-hexenyl	H	CH ₃	2-F Ph	H	O
542	5-hexenyl	H	CH ₃	2-F Ph	H	S
543	5-hexenyl	H	CH ₃		H	O
544	5-hexenyl	H	isopropyl	4-(CF ₃ O)Ph	H	O
545	5-hexenyl	H	Ph	4-(CF ₃ O)Ph	H	O
546	5-hexenyl	CH ₃ CO	CH ₃	2-CH ₃ Ph	CH ₃ CO	O
547	CH ₃ O	H	H	cyclobutyl	H	O

Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
548	CH ₃ O	H	H	2,4-di-F Ph	H	O
549	CH ₃ O	H	H	(4-CH ₃) PhCH ₂	H	O
550	CH ₃ O	H	H	2-quinolinyl	H	O
551	CH ₃ O	H	CH ₃	CH ₃	H	O
552	CH ₃ O	H	CH ₃	CH ₃	H	S
553	CH ₃ O	H	CH ₃	3-CF ₃ Ph	H	O
554	CH ₃ O	H	CH ₃	2-furyl	H	O
555	CH ₃ O	H	CH ₃	2-furyl	H	S
556	CH ₃ O	H	CH ₃	2-thienyl	H	O
557	CH ₃ O	H	CH ₃	3-(4-(OCH ₃)thienyl)	H	O
558	CH ₃ O	H	CH ₃		H	O
559	CH ₃ O	H	n-propyl	4-(CF ₃ O)Ph	H	O
560	CH ₃ O	H	4-F Ph	4-(CF ₃ O)Ph	H	O
561	CH ₃ O	Br	isobutyl	3-CF ₃ Ph	Br	O
562	CH ₃ O	H	CH ₃		H	O
563	EtO	3-F Ph	Et	cyclopentyl	H	O
564	EtO	H	H	CH ₃	H	O
565	EtO	H	H	CH ₃	H	S
566	EtO	H	H	3,4-di-CH ₃ Ph	H	O
567	EtO	H	CH ₃	n-propyl	H	O
568	EtO	H	CH ₃	cyclobutyl	H	O
569	EtO	H	CH ₃	cycloheptyl	H	O
570	EtO	H	CH ₃	cycloheptyl	H	S
571	EtO	H	CH ₃		H	O
572	EtO	H	CH ₃	3,4-di-F Ph	H	O

Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
573	EtO	H	CH ₃		H	O
574	EtO	H	n-butyl	2-thienyl	H	O
575	EtO	H	Ph	2-thienyl	H	O
576	EtO	CH ₃	CH ₃	4-Br Ph	H	O
577	EtO	Cl	CH ₃	n-hexyl	H	O
578	EtO	Cl	CH ₃	2-Cl Ph	H	O
579	EtO	Cl	CH ₃	2-Cl Ph	H	S
580	EtO	Cl	n-butyl	Ph	Cl	O
581	(i-Pr)O	H	H	CH ₃	H	O
582	(i-Pr)O	H	H	CH ₃	H	S
583	(i-Pr)O	H	H	3, 5-di-Cl Ph	H	O
584	(i-Pr)O	H	CH ₃		H	O
585	(i-Pr)O	H	CH ₃	3-Cl-5-F Ph	H	O
586	(i-Pr)O	H	CH ₃	3-Cl-5-F Ph	H	S
587	(i-Pr)O	H	CH ₃		H	O
588	(i-Pr)O	H	isopropyl	4-Br Ph	H	O
589	(i-Pr)O	H	4-F Ph	3,4-di-F Ph	H	O
590	(i-Pr)O	CH ₃	Et	2-thienyl	H	O
591	(i-Pr)O	CH ₃ CO	Et	2-thienyl	CH ₃ CO	O
592	(i-Pr)O	Cl	3-F Ph	2,4-di-F Ph	Cl	O
593	n-BuO	H	H	cyclopentyl	H	O
594	n-BuO	H	H	cyclooctyl	H	O
595	n-BuO	H	H	cyclooctyl	H	S
596	n-BuO	H	Et	cyclooctyl	H	O
597	n-BuO	H	Et	Ph	H	O
598	n-BuO	H	Et	2,4-di-F Ph	H	O
599	n-BuO	H	Et	PhOCH ₂	H	O
600	n-BuO	H	isopropyl	cyclooctyl	H	O

Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
601	n-BuO	H	n-hexyl	cyclooctyl	H	O
602	n-BuO	CH ₃	CH ₃	3,5-di-F Ph	H	O
603	n-BuO	PrCO	Et	3,5-di-CH ₃ Ph	H	O
604	n-BuO	Br	Ph	cyclooctyl	Br	O
605	(n-pentyl)O	H	CH ₃	3-Br Ph	H	O
606	(n-pentyl)O	H	CH ₃	3-Br Ph	H	S
607	(n-pentyl)O	H	CH ₃	2-naphthyl	H	O
608	(n-pentyl)O	H	CH ₃		H	O
609	(n-hexyl)O	H	CH ₃	cyclopropyl	H	O
610	(n-hexyl)O	H	CH ₃	n-pentyl	H	O
611	(n-hexyl)O	H	CH ₃	3-Br Ph	H	O
612	(n-hexyl)O	H	CH ₃	2-naphthyl	H	O
613	(i-hexyl)O	CH ₃ OCH ₂	Et	Ph	H	O
614	(i-hexyl)O	CH ₃ OCH ₂	Et	Ph	H	S
615	CO ₂ H	H	H	3, 5-di-Cl Ph	H	O
616	CO ₂ H	H	CH ₃	3, 5-di-Cl Ph	H	O
617	CO ₂ H	H	propyl	Ph	H	O
618	CO ₂ H	H	propyl		H	O
619	CO ₂ H	H	CH ₃	Ph	H	O
620	CO ₂ H	H	CH ₃		H	O
621	CO ₂ H	H	CH ₃		H	O
622	CO ₂ H	CH ₃	CH ₃	3, 5-di-Cl Ph	H	O
623	CO ₂ H	CH ₃	isopropyl	3-Br Ph	H	O
624	CO ₂ H	CH ₃	isopropyl	3-Br Ph	CH ₃	O
625	CO ₂ H	CH ₃	4-F Ph	propyl	H	O
626	CO ₂ H	Et	H	4-F Ph	H	O

Entry No.	R ¹	R ²	R ³	R ⁴	R ⁵	X
627	CO ₂ H	Et	H	4-F Ph	Et	O
628	CO ₂ H	Et	CH ₃	4-F Ph	Et	O
629	CO ₂ H	Et	propyl	Ph	H	O
630	CO ₂ H	Et	propyl	Ph	H	S
631	CO ₂ H	Ph	CH ₃	2-furyl	H	O
632	CO ₂ H	Ph	CH ₃	2-furyl	H	S
633	CO ₂ H	3-Br Ph	Ph	2-thienyl	H	O
634	CO ₂ H	n-PrCO	H	3-Cl Ph	H	O
635	CO ₂ H	n-PrCO	H	3-pyridyl	H	O
636	CO ₂ H	n-PrCO	H		H	O
637	CO ₂ H	n-PrCO	CH ₃	3-Cl Ph	H	O
638	CO ₂ H	n-PrCO	CH ₃	3-Cl Ph	n-PrCO	O
639	CO ₂ H	n-pentylCO	Ph	3-Cl Ph	H	O

A salt of a compound of Formula I may be prepared in situ during the final isolation and purification of a compound or by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed. Likewise, when the compound of Formula I contains a carboxylic acid moiety, (e.g., R = H), a salt of said compound of Formula I may be prepared by separately reacting it with a suitable inorganic or organic base and isolating the salt thus formed. The term "pharmaceutically acceptable salt" refers to a relatively non-toxic, inorganic or organic acid addition salt of a compound of the present invention (see, e.g., Berge et al., J. Pharm. Sci. 66:1-19, 1977).

Representative salts of the compounds of Formula I include the conventional non-toxic salts and the quaternary ammonium salts which are formed, for example, from inorganic or organic acids or bases by means well known in the art. For example, such acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cinnamate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride,

hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, itaconate, lactate, maleate, mandelate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, sulfonate, tartrate, thiocyanate, tosylate, undecanoate, and the like.

Base salts include, for example, alkali metal salts such as potassium and sodium salts, alkaline earth metal salts such as calcium and magnesium salts, and ammonium salts with organic bases such as dicyclohexylamine and N-methyl-D-glucamine. Additionally, basic nitrogen containing groups in the conjugate base may be quaternized with such agents as lower alkyl halides such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, and dibutyl sulfate; and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; aralkyl halides like benzyl and phenethyl bromides, and the like.

The esters of Formula I in the present invention are non-toxic, pharmaceutically acceptable esters, for example, alkyl esters such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or pentyl esters. Additional esters such as, for example, methyl ester or phenyl-C₁-C₅ alkyl may be used. The compound of Formula I may be esterified by a variety of conventional procedures including reacting the appropriate anhydride, carboxylic acid, or acid chloride with the alcohol group of the Formula I compound. The appropriate anhydride may be reacted with the alcohol in the presence of a base to facilitate acylation such as 1,8-bis[dimethylamino]naphthalene or N,N-dimethylaminopyridine. An appropriate carboxylic acid may be reacted with the alcohol in the presence of a dehydrating agent such as dicyclohexylcarbodiimide, 1-[3-dimethylaminopropyl]-3-ethylcarbodiimide, or other water soluble dehydrating agents which are used to drive the reaction by the removal of water, and optionally, an acylation catalyst. Esterification may also be effected using the appropriate carboxylic acid in the presence of trifluoroacetic anhydride and optionally, pyridine, or in the presence of N,N-carbonyldiimidazole with pyridine. Reaction of an acid chloride with the alcohol may be carried out with an acylation catalyst such as 4-DMAP or pyridine.

One skilled in the art would readily know how to successfully carry out these as well as other methods of esterification of alcohols.

Additionally, sensitive or reactive groups on the compound of Formula I may need to be protected and deprotected during any of the above methods for forming esters. Protecting groups in general may be added and removed by conventional methods well known in the art (see, e.g., T. W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*; Wiley: New York, (1999)).

The compounds of Formula I may contain one or more asymmetric centers, depending upon the location and nature of the various substituents desired. Asymmetric carbon atoms may be present in the (*R*) or (*S*) configuration. Preferred isomers are those with the absolute configuration which produces the compound of Formula I with the more desirable biological activity. In certain instances, asymmetry may also be present due to restricted rotation about a given bond, for example, the central bond adjoining two aromatic rings of the specified compounds.

Substituents on a ring may also be present in either *cis* or *trans* form, and a substituent on a double bond may be present in either *Z* or *E* form.

It is intended that all isomers (including enantiomers and diastereomers), either by nature of asymmetric centers or by restricted rotation as described above, as separated, pure or partially purified isomers or racemic mixtures thereof, be included within the scope of the instant invention. The purification of said isomers and the separation of said isomeric mixtures may be accomplished by standard techniques known in the art.

The particular process to be utilized in the preparation of the compounds of this invention depends upon the specific compound desired. Such factors as the selection of the specific X moiety, and the specific substituents possible at various locations on the molecule, all play a role in the path to be followed in the preparation of the specific compounds of this invention. Those factors are readily recognized by one of ordinary skill in the art.

In general, the compounds of this invention may be prepared by standard techniques known in the art and by known processes analogous thereto. The compounds of Formula I may generally be synthesized according to Reaction Schemes 1, 2, and 3. Reaction Schemes 1 and 2 demonstrate how to make intermediates that are coupled in Reaction Scheme 3 to provide the compounds of Formula I.

Route (A) of Reaction Scheme 1 provides a method to prepare compounds 4 and 5 where R¹ is C₁-C₆ lower alkyl or benzyl, R³ is not hydrogen, and X is O. The first step shows protection of the acid group of a commercially available aspartate derivative compound 1 by means well known in the art such as, for example, by forming a silyl ester, followed by N-acylation with the appropriate R⁴-acid derivative, R⁴COY, where Y is a leaving group such as halo. Finally, the compound is deprotected by means well known in the art such as, for example, in the case of a silyl ester, an aqueous work up, to give compound 2. Alternatively, condensation of the protected form of compound 1 with a free carboxylic acid such as R⁴COOH in the presence of a dehydrating reagent, such as DCC or EDCI, also provides compound 2. Compound 2 may then be converted to compound 3, where R³ is as defined for Formula I compounds by several methods. For

example, one such method, when $R^3 = \text{Me}$, is the well known Dakin-West reaction which is typically performed using acetic anhydride and pyridine. When R^3 is other than hydrogen, compound 2 may be converted to an acid chloride with a reagent such as thionyl chloride and reacted with a Grignard reagent such as $R^3\text{Mg-halo}$, to provide compound 3. Other methods for the formation of ketones of compound 3 from acids and acid derivatives may also be employed, for example, by using Weinreb amides, which are known to those skilled in the art. Compound 3 is then cyclized under acid dehydrative conditions using, for example, phosphorus oxychloride, or a mixture of sulfuric acid and acetic anhydride, generally with heating, to provide compound 4 where X is O and the R^3 group is attached at the 5 position.


It will be recognized by those skilled in the art that compound 4 and thus, compound 5, may exist in two regioisomeric forms with respect to the attachment point of the R^3 , $\text{CH}_2\text{CO}_2\text{R}''$, and $\text{CH}_2\text{CH}_2\text{OH}$ groups. Using Route (B), one can prepare compound 4 in which the R^3 is attached at the 4-position and carboxymethyl side chain is attached at the 5-position, that is, the groups are reversed from that of Route (A). In Route (B), a commercially available amino acid, compound 6, may be acylated under basic conditions, for example, with aqueous sodium hydroxide, with an appropriate R^4 -acid derivative, (e.g., $R^4\text{COY}$), where Y is a leaving group such as chloro, to provide the N-acylated product 7. Compound 7 may be then coupled with an acetic acid ester in the presence of a strong non-nucleophilic base to make the keto ester 8, where R'' is $\text{C}_1\text{-C}_6$ alkyl or benzyl. Cyclization of compound 8 using a dehydrating reagent such as POCl_3 provides compound 4 where $X = \text{O}$ and R^3 is attached at the 4 position. Reaction of compound 8 with a nucleophilic S reagent such as P_2S_5 in solvents such as pyridine or acetonitrile/triethylamine, with heating as necessary, gives compound 4 where $X = \text{S}$ and R^3 is attached at the 4 position.

Route (C) of Reaction Scheme 1 depicts the preparation of compound 4 from ketoesters 9 or 10, where Y is a leaving group such as halo and R'' is $\text{C}_1\text{-C}_6$ alkyl or benzyl. Either compound 9 or 10 may be chosen as the starting material depending on whether the R^3 group in the desired end product is hydrogen or is attached at the 4 or 5 position. Accordingly, compound 9 or 10 may be reacted with an amide or thioamide where X is either O or S to yield compound 4. Ketoesters 9 or 10 are commercially available, or may be prepared by methods well known in the art such as by bromination of commercially available ketoesters 9 and 10 where Y is hydrogen. Amides ($R^4\text{C}(=\text{X})\text{NH}_2$) where X is O may be commercially available carboxylic amides, or may be prepared from the corresponding available acids or acid chlorides by well known methods. Thioamides ($R^4\text{C}(=\text{X})\text{NH}_2$) where X is S may be commercially available thioamides, or may be

prepared from the corresponding available amides by known methods such as through the use of Lawesson's reagent. Reaction of ketoester 9 with an amide or thioamide in the presence of a base provides compound 4 as an oxazole or a thiazole, respectively, where R^3 is other than hydrogen and located at the 4-position. Reaction of ketoester 10 with an amide or thioamide in the presence of base provides compound 4 as an oxazole or thiazole, where R^3 is located at the 5-position.

Routes (A), (B), and (C) each provide compound 4 where R^3 and R^4 are each as described for a compound of Formula I and where R'' is a lower alkyl or benzyl. Compound 4 may then be reduced to compound 5 using reducing agents such as lithium aluminum hydride, lithium borohydride, or other suitable hydride donors under conditions well known in the art.

Reaction Scheme 1



Reaction Scheme 2 depicts the conversion of commercially available hydroxy ketone 11 to a protected derivative 12, by reaction with R^7-Y in the presence of a base,

where R^7 is C_1 - C_6 alkyl optionally substituted with phenyl or oxo, C_1 - C_6 trialkylsilyl, arylalkylsilyl, or COR^8 ; and R^8 is C_1 - C_6 alkyl or phenyl optionally substituted with C_1 - C_6 alkyl, halo, or nitro; and Y is a leaving group. "C₁-C₆ trialkylsilyl" means three independently selected straight or branched chain alkyl groups having from one to about six carbon atoms, each of which are bound to silicon and includes such groups as trimethylsilyl, *tert*-butyldimethyl silyl, and the like. "Arylalkylsilyl" means at least one phenyl or substituted phenyl group bound to silicon, with an appropriate number of independently selected straight or branched chain alkyl groups having from one to about six carbon atoms, each of which are also bound to silicon, and includes such groups as *t*-butyldiphenylsilyl, methyldiphenylsilyl, dimethylpentafluorophenylsilyl, and the like. "Leaving group" includes halides such as I, Br, and Cl; carboxylates such as acetates, and trifluoroacetates; and aryl and alkyl sulfonates such as methanesulfonates (mesylates) and *p*-toluene sulfonates (tosylates), and the like.

Compound 12 is substituted with R^2 (as described in Formula I) by means of, for example, reaction with a source of electrophilic halogen, or a Friedel-Crafts reaction in the presence of a Lewis acid and R^2 -Y where Y is as described above, to form a substituted ketone 13. Alternatively, a halogenated compound formed in this manner (for example, substituted with bromine or iodine) may be reacted with a range of coupling partners under metal catalysis, using complexes and compounds of elements such as palladium and nickel well known to those skilled in the art, to form further substituted ketone 13. Examples of such catalysts include tetrakis(triphenylphosphine)palladium(0) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), and similar nickel(0) and nickel(II) compounds; and examples of coupling partners include boronic acids and esters (the well known Suzuki coupling, carried out in solvents such as toluene in the presence of a base such as potassium carbonate), and organometallics such as Grignard reagents, organozincs (Negishi coupling), and organotin derivatives (Stille coupling), reaction conditions for which are widely known. Furthermore, such halogenated compounds may be coupled with secondary amines such as piperidine using similar palladium or nickel catalysts (Hartwig or Buchwald coupling) to provide further substituted ketones 13.

Further reaction of compound 13 with a halogen source or R^5 -Y, (where R^5 is as described in Formula I), under similar conditions gives disubstituted compound 14. A Wittig reaction, or the Horner-Emmons-Wadsworth variation, each well known in the art, may be used to convert 14 to compound 15. For example, reaction of compound 14 with a trialkylphosphonoacetate, where R'' is lower alkyl and R is as described in Formula I, in the presence of a strong base such as sodium hydride, provides compound 15. In like manner, compound 13 may be converted to compound 15 where R^5 is H.

Regardless of the isomeric mixture of isomers of 15 produced in the reaction, either isomer (*E* or *Z*) or a mixture of both, may be converted to the corresponding compound 17 by catalytic hydrogenation or reduction with a hydride reagent capable of 1,4 (conjugate) addition, which are known to those skilled in the art. This route is particularly advantageous for preparing compound 17 where R^1 is hydrogen.


Compound 17 where R^1 is COOR, may be prepared through standard condensation reactions, for example, the well known Knoevenagel reaction. In such cases, the ketone 13 or 14 may be reacted with a suitable active-hydrogen coupling partner, under the influence of acidic reagents such as titanium tetrachloride, or basic reagents such as piperidine, in appropriate solvents. The product 15b (compound 15 where R^1 is COOR), may be reduced to 17b (compound 17 where R^1 is COOR), which may be further alkylated with another R^1 group in the presence of base, hydrolyzed and decarboxylated to give 17d (compound 17 where R^1 is other than COOH and R is H). Reesterification of 17d and removal of the protecting group R^7 would afford 17c. Reesterification may be performed using standard conditions using the well known Fischer esterification by treatment with an acid and an alcohol or by reaction with diazoalkyl reagents or with an electrophilic species such as, for example, methyl iodide or dimethyl sulfate. Compound 17 where R^1 is alkoxy may be prepared by a similar condensation reaction of ketone 13 or 14 with a silylated enol ester of Formula $R^1CH=C(OR'')O\text{-alkylsilyl}$, where R^1 is alkoxy, under the influence of acidic reagents such as titanium tetrachloride, and reducing the intermediate compound 15, where R^1 is alkoxy, in the presence of hydrogen and a catalyst as described above.

A general coupling reaction of compound 13 or 14 via the Reformatsky reaction produces compound 16 (Formula II), when R^1 is alkyl, or compound 15a when R^1 is H. The ketone is condensed with an appropriate organozinc reagent prepared in situ from Zn and R^1CHYCO_2R , where Y is halo. The alpha-halo ester compounds of formula R^1CHYCO_2R , are either commercial reagents or are prepared by halogenation of commercially available $R^1CH_2CO_2R$ compounds by methods well known to those skilled in the art.

The conversion of 16 to 17 may be accomplished by standard hydrogenation conditions, for example, Pd/C and hydrogen; and deprotection of compound 17, where R^7 is a protecting group, to compound 17c, where R^7 is hydrogen, may be accomplished by standard means. For example, when the R^7 group is alkyl (e.g., methyl), the compound 17a may be generated by nucleophilic cleavage with a reagent such as an alkali metal thiolate. Alternatively, compound 17 when R^7 is methyl, may be converted to compound 17c by reaction with a Lewis acid such as a bromoborane. When R^7 is benzyl, the

compound 17 may be converted to 17c under hydrogenation conditions, typically carried out using a catalyst such as palladium. Other conditions for the removal of the protecting group R^7 from compound 17, where R^7 is other than hydrogen which produces the hydroxy compound 17c, are dependent on the specific protecting group chosen from among those which are well known by those skilled in the art.


Reaction Scheme 2



The final step in the preparation of Formula I compounds is shown in Reaction Scheme 3. The alcohol 5 (from Reaction Scheme 1) is coupled with the hydroxy indane 17c (from Reaction Scheme 2) via a Mitsunobu coupling, facilitated by an azodicarboxylate reagent such as DEAD, and a phosphine such as triphenylphosphine to make the compounds of Formula I. Alternatively, the hydroxy group of alcohol 5 is converted to a leaving group such as halo, tosylate (OTs), or mesylate (OMs), by reaction with a halogenating agent such as thionyl chloride or CCl₄/triphenylphosphine; or by reaction with a Y-halo compound, where Y is tosyl (Ts) or mesyl (Ms), in the presence of a base, providing compound 18. Compound 18 may be reacted with compound 17c in the presence of a base, providing the compounds of Formula I.


Compounds of Formula I in which R is alkyl, may be converted to compounds of Formula I in which R is H by treatment with a base (e.g., KOH) in a suitable solvent (e.g., methanol, THF, or water, or mixtures thereof) with heating. Alternatively, this conversion may be accomplished by reaction with a nucleophile such as iodide or cyanide, in a suitable solvent, such as pyridine. In addition, when R is benzyl, the cleavage to compounds of Formula I in which R is H may be effected through hydrogenolysis by means well known in the art.

Reaction Scheme 3



An alternative route to Formula I compounds, useful when X = S and the R⁴ group contains one or more R⁶ substituents labile to the reaction conditions of Scheme 1 or 2, is shown in Reaction Scheme 3a.

Reaction Scheme 3a



In Scheme 3a, a 2-aminothiazole **4** may be prepared using thiourea (similar to Route C, Reaction Scheme 1) and converted to a 2-halo thiazole **5a** as shown above (Erlenmeyer et al., *Helv. Chim. Acta* 28:362-363, 1945). Mitsunobu coupling of **5a** by a method analogous to Reaction Scheme 3 is then accomplished, and product **19** is further elaborated by a Palladium-catalyzed cross-coupling reaction to introduce the R⁴


substituent. Hydrolysis as described in Reaction Scheme 3 gives Formula I compounds where R = H.

The foregoing reaction schemes are further illustrated by the specific Examples described herein.


The salts and esters of this invention may be readily prepared by conventional chemical processes as described previously herein.

The invention is further directed to novel Formula II compounds (compound 16) and Formula III (compounds 17, including compounds 17a-d) compounds shown in Reaction Scheme 2. These compounds are useful in the preparation of the compounds of Formula I, and are further described as follows.

The present invention encompasses the compounds of Formula II and Formula III,



(II)




(III)

wherein

R, R¹, R², R³, R⁴, R⁵, R⁶, and X are as defined for Formula I above; and

R⁷ is H, C₁-C₆ alkyl optionally substituted with phenyl or oxo, C₁-C₆ trialkylsilyl, arylalkylsilyl, COR⁸, COOR⁸, or



R⁸ is C₁-C₆ alkyl, or phenyl optionally substituted with C₁-C₆ alkyl, halo, or nitro; and the salts thereof.

C₁-C₆ trialkylsilyl means three independently selected straight or branched chain alkyl groups having from one to about six carbon atoms, each of which are bound to silicon and includes such groups as trimethylsilyl, *tert*-butyldimethyl silyl, and the like.

Arylalkylsilyl means at least one phenyl or substituted phenyl group bound to silicon, with an appropriate number of independently selected straight or branched chain alkyl groups having from one to about six carbon atoms, each of which are also bound to silicon, and includes such groups as *t*-butyldiphenylsilyl, methyl diphenylsilyl, dimethylpentafluorophenylsilyl, and the like.

The salts of this invention may be readily prepared by conventional chemical processes as described previously herein.


The compounds of Formula II and Formula III may each contain one or more asymmetric centers, depending upon the location and nature of the various substituents desired. Asymmetric carbon atoms may be present in the (*R*) or (*S*) configuration. Preferred isomers are those with the absolute configuration which produces the compound of Formula II or Formula III that will be useful in producing the compounds of Formula I having a more desirable biological activity. In certain instances, asymmetry may also be present due to restricted rotation about a given bond, for example, the central bond adjoining two aromatic rings of the specified compounds.

Substituents on a ring may also be present in either *cis* or *trans* form, and a substituent on a double bond may be present in either *Z* or *E* form.

It is intended that all isomers (including enantiomers and diastereomers), either by nature of asymmetric centers or by restricted rotation as described above, as separated, pure or partially purified isomers or racemic mixtures thereof, be included within the scope of the present invention. The purification of said isomers and the separation of said isomeric mixtures may be accomplished by standard techniques known in the art, as well as by the novel means described herein.

For example, Formula II compounds may contain an asymmetric center (labeled **C-2**) and Formula III compounds may contain two asymmetric centers (labeled **C-2** and **C-1'**) which give rise to enantiomers and diastereomers. Examples of these and other compounds of Formula II and Formula III, which are illustrative of the present invention, are shown in Table 2.

Table 2
Illustrative Examples of Compounds II and III



Entry No.	Formula	absolute configuration		R ¹	R ²	R ⁵	R ⁷
		C-2	C-1'				
1	II	R	---	H	H	H	CH ₃

Entry No.	Formula	absolute configuration		R ¹	R ²	R ⁵	R ⁷
		C-2	C-1'				
2	III	R	R	H	H	H	CH ₃
3	II	R	---	Cl	H	H	t-Bu(CH ₃) ₂ Si
4	III	R	S	Cl	H	H	t-Bu(CH ₃) ₂ Si
5	II	S	---	H	H	H	CH ₃
6	III	S	S	H	H	H	CH ₃
7	II	R	---	CH ₃	H	H	CH ₃
8	III	R	R	CH ₃	H	H	CH ₃
9	II	S	---	CH ₃	H	H	CH ₃
10	III	S	R	CH ₃	H	H	CH ₃
11	II	R	---	CH ₃	H	H	PhCH ₂
12	III	R	S	CH ₃	H	H	PhCH ₂
13	II	S	---	CH ₃	H	H	PhCH ₂
14	III	S	S	CH ₃	H	H	PhCH ₂
15	II	R	---	CH ₃	H	H	t-Bu(CH ₃) ₂ Si
16	III	R	R	CH ₃	H	H	t-Bu(CH ₃) ₂ Si
17	II	S	---	CH ₃	H	H	t-Bu(CH ₃) ₂ Si
18	II	R	---	CH ₃	H	H	t-BuCO
19	III	R	S	CH ₃	H	H	t-BuCO
20	II	S	---	CH ₃	H	H	t-BuCO
21	III	S	S	CH ₃	H	H	t-BuCO
22	II	R	---	CH ₃	CH ₃	H	PhCH ₂
23	II	R	---	CH ₃	CH ₃ CO	H	PhCH ₂
24	II	S	---	CH ₃	2-thienyl	H	t-Bu(CH ₃) ₂ Si
25	III	S	R	CH ₃	2-thienyl	H	t-Bu(CH ₃) ₂ Si
26	II	S	---	CH ₃	Ph	H	CH ₃
27	II	R	---	CH ₃	Cl	H	CH ₃
28	II	S	---	CH ₃	Cl	H	CH ₃
29	III	S	S	CH ₃	Cl	H	CH ₃
30	II	R	---	CH ₃	Br	H	Ph(CH ₃) ₂ Si
31	III	R	R	CH ₃	Br	H	Ph(CH ₃) ₂ Si
32	II	S	---	CH ₃	Br	H	Ph(CH ₃) ₂ Si
33	III	S	R	CH ₃	Br	H	Ph(CH ₃) ₂ Si

Entry No.	Formula	absolute configuration		R ¹	R ²	R ⁵	R ⁷
		C-2	C-1'				
34	II	S	---	CH ₃	Cl	Cl	CH ₃
35	II	R	---	Et	H	H	CH ₃
36	III	R	R	Et	H	H	CH ₃
37	II	S	---	Et	H	H	PhCH ₂
38	III	S	S	Et	H	H	PhCH ₂
39	II	R	---	Et	H	H	t-Bu
40	II	S	---	Et	H	H	t-Bu
41	II	S	---	Et	CH ₃	H	Ph(CH ₃) ₂ Si
42	III	S	S	Et	CH ₃	H	Ph(CH ₃) ₂ Si
43	II	R	---	Et	n-propyl	H	CH ₃
44	II	S	---	Et	Ph	H	CH ₃
45	II	S	---	Et	3-Cl Ph	H	t-Bu(CH ₃) ₂ Si
46	III	S	R	Et	3-Cl Ph	H	t-Bu(CH ₃) ₂ Si
47	II	S	---	Et	4-pyridyl	H	t-Bu(CH ₃) ₂ Si
48	III	S	S	Et	4-pyridyl	H	t-Bu(CH ₃) ₂ Si
49	II	S	---	Et	CH ₃	H	Ph(CH ₃) ₂ Si
50	II	R	---	Et	n-propyl	Cl	CH ₃
51	II	R	---	Et	Br	Br	t-Bu(CH ₃) ₂ Si
52	III	R	R	Et	Br	Br	t-Bu(CH ₃) ₂ Si
53	II	S	---	CF ₃ CH ₂	H	H	CH ₃
54	II	S	---	CF ₃ CH ₂	CH ₃	CH ₃	(4-CH ₃ O)PhCH ₂
55	III	S	S	CF ₃ CH ₂	CH ₃	CH ₃	(4-CH ₃ O)PhCH ₂
56	II	S	---	n-propyl	H	H	(i-Pr) ₃ Si
57	II	R	---	n-propyl	PrCO	PrCO	t-Bu
58	II	R	---	n-propyl	Cl	Cl	(i-Pr) ₃ Si
59	III	R	R	n-propyl	Cl	Cl	(i-Pr) ₃ Si
60	II	S	---	isopropyl	CH ₃	H	CH ₃
61	III	S	R	isopropyl	CH ₃	H	CH ₃
62	II	R	---	isopropyl	n-hexyl	H	(4-CH ₃ O)PhCH ₂
63	III	R	S	isopropyl	n-hexyl	H	(4-CH ₃ O)PhCH ₂
64	II	S	---	n-butyl	H	H	PhCH ₂
65	II	S	---	n-butyl	CH ₃ OCH ₂	H	t-Bu(CH ₃) ₂ Si

Entry No.	Formula	absolute configuration		R ¹	R ²	R ⁵	R ⁷
		C-2	C-1'				
66	III	S	S	n-butyl	CH ₃ OCH ₂	H	t-Bu(CH ₃) ₂ Si
67	II	R	—	n-butyl	Cl	H	CH ₃
68	II	R	—	n-pentyl	Cl	Cl	(4-CH ₃ O)PhCH ₂
69	II	S	—	n-pentyl	2-thienyl	2-thienyl	CH ₃
70	III	S	S	n-pentyl	2-thienyl	2-thienyl	CH ₃
71	II	R	—	n-hexyl	CH ₃ CO	H	t-Bu(CH ₃) ₂ Si
72	III	R	S	n-hexyl	CH ₃ CO	H	t-Bu(CH ₃) ₂ Si
73	II	R	—	n-hexyl	Ph	H	Ph(CH ₃) ₂ Si
74	III	R	R	n-hexyl	Ph	H	Ph(CH ₃) ₂ Si
75	II	R	—	cyclopropyl	H	H	t-BuCO
76	II	S	—	cyclopropyl	CH ₃	H	(i-Pr) ₃ Si
77	II	S	—	cyclobutyl	H	H	CH ₃
78	III	S	S	cyclobutyl	H	H	CH ₃
79	II	S	—	cyclobutyl	Cl	Cl	(4-CH ₃ O)PhCH ₂
80	II	R	—	cyclopentyl	CH ₃	H	t-Bu(CH ₃) ₂ Si
81	III	R	S	cyclopentyl	CH ₃	H	t-Bu(CH ₃) ₂ Si
82	II	S	—	cyclohexyl	Et	Et	CH ₃
83	II	R	—	cyclohexyl	2-thienyl	H	CH ₃ CO
84	II	R	—	cyclohexyl	Cl	H	CH ₃
85	III	R	R	cyclohexyl	Cl	H	CH ₃
86	II	S	—	2-propenyl	H	H	t-Bu(CH ₃) ₂ Si
87	II	R	—	2-propenyl	CH ₃	H	CH ₃ CO
88	II	S	—	2-isobutenyl	CH ₃ CO	H	CH ₃
89	II	S	—	5-hexenyl	CH ₃ CO	CH ₃ CO	CH ₃
90	II	S	—	CH ₃ O	H	H	PhCH ₂
91	III	S	R	CH ₃ O	H	H	PhCH ₂
92	II	R	—	CH ₃ O	3-F Ph	H	(4-CH ₃ O)PhCH ₂
93	II	S	—	EtO	Cl	Cl	PhCH ₂
94	III	S	R	EtO	Cl	Cl	PhCH ₂
95	II	R	—	(i-Pr)O	H	H	PhCH ₂
96	III	R	R	(i-Pr)O	H	H	PhCH ₂



Entry No.	Formula	absolute configuration		R ¹	R ²	R ⁵	R ⁷
		C-2	C-1'				
97	II	S	---	(n-pentyl)O	CH ₃	H	t-Bu(CH ₃) ₂ Si
98	III	S	S	(n-pentyl)O	CH ₃	H	t-Bu(CH ₃) ₂ Si
99	II	S	---	CO ₂ H	H	H	(4-CH ₃ O)PhCH ₂

Another embodiment of the present invention is an improved process for the preparation of compounds having a specific isomeric configuration when that specific configuration is desired for the ultimate desired end product of Formula I. The improved process yields these intermediate compounds in significantly greater diastereomeric excess than was heretofore possible.

Previously, for example, in the absence of stereocontrol during the hydrogenation step of Reaction Scheme 2, hydrogenation of a Formula II compound, where R¹ is alkyl may produce an unequal mixture of diastereomeric products of Formula III, in which one pair of enantiomers is favored because of the asymmetric nature of the starting material. Separation of such compounds may be accomplished by stepwise separation of the enantiomeric pairs, then by resolution of each enantiomer by crystallization or by chiral HPLC. Prior resolution of the starting material into a single enantiomer produces mixtures with enrichment of a single enantiomer that may likewise be separated.

However, when a compound of a specific relative configuration, namely a syn form (defined below) is desired, the yield is low when R¹ is alkyl, because the conditions of the hydrogenation step described in the art may favor the other (i.e., anti) diastereomers.


The desired isomeric configurations realized from this improved process are in the syn form where, for example, in compounds of Formula Va and Vb (depicted in Reaction Schemes 4 and 5), the R⁹ group and the 2' methylene carbon of the cyclopentane ring are both below the plane or are both above the plane. Anti diastereomers are those compounds where, for example, R⁹ is above the plane and 2' methylene is below the plane. This is further exemplified in Figures 1 and 2 below, in which solid wedge bonds are used to indicate projection of the bond above the plane and dashed wedge bonds are used to indicate projection of the bond below the plane.

Figure 1. syn diastereomers of Formula V**Figure 2. anti diastereomers of Formula V**

The improved process of this invention yields compounds in the syn form (Formulas Va and Vb, as drawn in Figure 1 and Reaction Schemes 4 and 5) in significantly higher diastereomeric excess than was generally possible.

The intermediate compounds used as starting materials for this process (compound IV of Reaction Schemes 4 and 5) are related to the compounds of Formula II (compound 16) in Reaction Scheme 2, and may be prepared by the same or analogous methods. These intermediates may be reacted under certain conditions to yield Formula V compounds that are related to compounds of Formula III (compounds 17 and 17a of Reaction Scheme 2), or to directly yield compounds of Formula I. However, due to the constraints of the improved process, only certain substituents are appropriate for completing this process.

Accordingly, the present invention relates to an improved process for the preparation of a substantially enriched syn form of a compound of Formula V,



wherein

R⁹ is methoxy optionally substituted by fluoro,

C₂-C₆ alkoxy, C₁-C₆ alkyl, or C₄-C₈ cycloalkyl each optionally substituted by fluoro, methylenedioxyphenyl or phenyl optionally substituted with R¹³;

R¹⁰ is hydrogen, fluoro, methyl optionally substituted with fluoro, oxo, or

C₂-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl,

each of which may be unsubstituted or substituted with R¹³,

or

R¹⁰ is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl,


each of which may be unsubstituted or substituted with R¹³;

R¹¹ is halo or C₁-C₆ alkyl optionally substituted with oxo;

R¹² is hydrogen, methyl optionally substituted with fluoro or oxo,

C₂-C₆ alkyl optionally substituted with phenyl, fluoro, or oxo,

C₁-C₆ trialkylsilyl, arylalkylsilyl, COR¹⁴, COOR¹⁴, or



R¹³ is fluoro, CF₃, C₁-C₆ alkyl optionally substituted with oxo, or C₁-C₆ alkoxy optionally substituted with fluoro;

R¹⁴ is C₁-C₆ alkyl, or phenyl optionally substituted with C₁-C₆ alkyl or fluoro;

R¹⁵ is hydrogen, C₁-C₆ alkyl or phenyl substituted with R¹³;

R¹⁶ is methyl optionally substituted with fluoro, oxo or

with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R¹³, or

C₄-C₈ cycloalkyl or C₂-C₆ alkyl, either of which may be unsubstituted or substituted with fluoro, methoxy, C₂-C₆ alkoxy optionally substituted with phenyl or C₁-C₆ alkoxy, oxo or with, phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R¹³, or

C₂-C₆ alkyl which may also be substituted with C₄-C₈ cycloalkyl or with phenoxy which may be unsubstituted or substituted with R⁶ or with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R¹³,

or

R¹⁶ is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,


each of which may be unsubstituted or substituted with R¹³, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl,

pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, pyrimidinyl or phenoxy

each of which may be unsubstituted or substituted with R¹³, and

X is O or S;

comprising hydrogenation of a racemic mixture or isolated optical isomer of a compound of Formula IV,



(IV)

wherein the substituents are as defined above, in the presence of hydrogen source, a catalyst, optionally in the presence of a base.

Substantially enriched syn form means at least about seventy percent (70%) or greater of one or both of the compounds of the configuration of Va or Vb. This is equivalent to at least about 40% de (diastereomeric excess) of the syn diastereomer. Diastereomeric excess of the syn diastereomer is calculated from the following formula:

$$\% \text{ de (syn)} = \frac{[\text{syn}] - [\text{anti}]}{[\text{syn}] + [\text{anti}]} \times 100 = \% \text{ syn diastereomer} - \% \text{ anti diastereomer}$$

in which

% de (syn) represents the diastereomeric excess of the syn diastereomer

[syn] represents the concentration of the syn diastereomer

[anti] represents the concentration of the anti diastereomer,

and where

$$\% \text{ syn} + \% \text{ anti} = 100\%.$$

Thus, a 40% de of the syn diastereomer is calculated from a mixture of 70% syn diastereomer and 30% anti diastereomer:

$$40\% \text{ de (syn)} = 70\% \text{ syn diastereomer} - 30\% \text{ anti diastereomer}$$

Catalyst means any of the transition metal catalysts well known in the art to effect hydrogenation reactions (P.A. Chaloner, *Handbook of Co-ordination Catalysis in Organic Chemistry*, Butterworth, 1986), and includes homogeneous hydrogenation catalysts. A homogeneous catalyst is a catalyst which is at least partially soluble in the reaction medium and which effects the reduction of a double bond in the presence of hydrogen. Such catalysts include, for example, ClRh[P(Ph)₃]₃ (Wilkinson's catalyst), (1,5-

cyclooctadiene)tricyclohexylphosphinepyridinoiridium(I) hexafluorophosphate, (1,5-cyclooctadiene)bis(methyldiphenylphosphine)iridium(I) hexafluorophosphate (Crabtree's catalysts), and the like.

Base means a substance with a pK_b sufficient to form a salt in situ with a carboxylic acid (see, e.g., *Advanced Organic Chemistry*, 3rd Ed., Jerry March, pp 220-222). The base which is used in this reaction may be any inorganic or organic base, and may be soluble in the reaction medium. Such bases include, for example, mono, di, and tri(C_1 - C_6 alkyl)amines such as isopropyl amine, diisopropyl amine, triethylamine, and the like; additional primary amines such as, for example, cyclohexane methylamine and ethanolamine; additional secondary amines such as, for example, morpholine and piperidine; and additional tertiary amines such as, for example, 1,8-diazaobicyclo[5.4.0]undec-7-ene and 1,5-diazabicyclo[4.3.0]non-5-ene as well as inorganic bases such as alkali metal and alkaline earth hydroxides, carbonates, bicarbonates, and optically active bases such as quinine, cinchonine or (+)- or (-)-alpha-methylbenzylamine.

Such bases also include, for example, the chiral bases named below that are useful for resolution.

Hydrogen source refers to any means of delivering hydrogen to the reaction medium and includes the use of hydrogen gas. Hydrogenation may be performed under a broad range of hydrogen pressures, that is, from about atmospheric pressure to about 1000 psi, preferably from about 20 to about 100 psi. Suitable hydrogenation solvents include, but are not limited to, protic solvents such as ethanol, methanol, water, 2-propanol, *tert*-butanol, methyl cellosolve and the like, and mixtures thereof, or optionally mixtures thereof with a miscible aprotic solvent such as THF, such that the hydrogenation catalyst, the base, and the starting material are each at least partially soluble.

The resolution of the starting indene acetic acid derivatives of Formula IV or of the indane acetic acid derivatives of Formula V may be accomplished by means well known in the art, for example, by using optically active bases as resolving agents such as, for example, a readily available base such as quinine, cinchonine or (+)- or (-)-alpha-methylbenzylamine. Choice of the base will depend on the solubility properties of the salt formed, so that resolution by differential recrystallization may be readily accomplished. By selecting bases with opposite absolute configuration, separation of the salt of each enantiomer may be accomplished. For example, for the embodiment illustrated in Reaction Scheme 4, the desired enantiomer IVc or IVd may be separated, and the undesired isomer may be recycled by racemization under basic conditions to the starting material of Formula IV.

Suitable crystallization solvents refer to those solvents in which one diastereomeric salt of a mixture is more soluble than the other, enabling them to be separated by recrystallization. Such solvents include, for example, acetonitrile, acetone, *t*-butanol, 2-propanol, ethanol, methanol, and the like, and mixtures thereof.

Aqueous mineral acids include, for example, the commonly used inorganic acids such as hydrochloric or sulfuric acid, and the like.


The process may be carried out starting with a racemate of Formula IV (see Reaction Scheme 4), or with a Formula V compound with the configuration at one asymmetric carbon which corresponds to that desired for the end product (see Reaction Scheme 5). Starting with the generally pure configuration is preferred, although either process will yield the desired configuration of the end product (V) in substantially enriched syn form.

One embodiment of this process is shown in the example of Reaction Scheme 4 and comprises the steps of

- (1) formation of diastereomeric salts of IVc and IVd by treatment of IV with a suitable basic resolving agent,
- (2) separation of the diastereomeric salts IVc and IVd by crystallization in a suitable crystallization solvent,
- (3) optionally liberating the individual antipodes IVa and IVb from the separated salts by treatment with aqueous mineral acid, and
- (4) reduction of either the separated diastereomeric salts IVc and Vd or the individual antipodes IVa and IVb by hydrogenation in the presence of a homogeneous hydrogenation catalyst, a suitable solvent and a base, wherein M⁺ is a cation selected from an alkali metal, alkaline earth metal, ammonium, and mono-, di-, tri- or quaternary alkylammonium or aralkylammonium, and R⁹-R¹² are as defined above.

The enantiomeric purity of the product Va and Vb will correspond to the enantiomeric purity of the isomer IVa or IVb used, respectively, but will not include any substantial amount of the other (anti) diastereoisomer.


Reaction Scheme 4



A second embodiment of this process is shown in Reaction Scheme 5 and comprises the steps of

- (1) reduction of the indene carboxylic acid of Formula IV by hydrogenation in the presence of a homogeneous hydrogenation catalyst, a suitable solvent, and a base,
- (2) formation of diastereomeric salts of Vc and Vd by treatment of V with a suitable basic resolving agent,
- (3) separation of the diastereomeric salts Vc and Vd by crystallization in a suitable crystallization solvent, and
- (4) liberating the individual antipodes Va and Vb from the separated salts by treatment with aqueous mineral acid.

Reaction Scheme 5




The resolution of the racemate of either Formula IV or Formula V compounds may be accomplished by means well known in the art, such as by chiral HPLC, crystallization of chiral salt derivatives, chiral ester derivatives, and the like.


The determination of absolute chirality of IVa, IVb, IVc, IVd, Va, and Vb may be accomplished by several means known to those skilled in the art. X-ray crystallographic

methods may provide such information under certain well-established conditions. For example, the presence in the crystallographic unit cell of another component of known chirality, such as a chiral resolving agent or auxiliary in the form of a salt, complex, or covalently attached group, may allow such determination. Another method known in the art, heavy atom scattering technique may be utilized when the compound to be assayed contains an atom of sufficient mass (for example, bromine or iodine). Other methods involving optical properties and the use of plane-polarized light may also be employed. For example, one skilled in the art would recognize that such techniques as circular dichroism may be applicable to a given structure or structural class.

Specific examples of the intermediates that may be made with the process of the present invention are shown below by way of example, and not by way of limitation, and may be used for the preparation of compounds of Formula I of the same absolute configuration.



Compounds of Formula III in which R¹ = H may also be prepared in an optically active fashion by the methods summarized in Reaction Scheme 6. Resolution of racemic ester **17a** (Formula III, where R¹ is H) may be accomplished by selective enzymatic hydrolysis using Amano Lipase PS to yield **17f**. Alternatively, **17e**, which may be prepared by hydrolysis of **17a**, may be resolved by crystallization of the diastereomeric salts formed with an optically active amine, for example, (*S*)-(-)- α -methyl-benzylamine, followed by regeneration of the carboxylic acid by treating the salt with mineral acid. Further conversion of **17f** to the intermediates **17g** and **17h** may be accomplished by means analogous to that described for the preparation of **17c** in Reaction Scheme 2: reesterification and removal of the R⁷ protecting group.

Reaction Scheme 6

The compounds of Formula I are effective in the treatment of Type II diabetes mellitus (including associated diabetic dyslipidemia and other diabetic complications), as well as for a number of other pharmaceutical uses associated therewith, such as hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, Syndrome X, and insulin resistance. In addition, the compounds of the present invention are also effective in the regulation of appetite and food intake in such disorders as obesity, and in the treatment of atherosclerotic disease, hyperlipidemia, hypercholesteremia, low HDL levels, hypertension, cardiovascular disease (including atherosclerosis, coronary heart disease, coronary artery disease, and hypertension), cerebrovascular disease and peripheral vessel disease; and for the treatment of lupus, polycystic ovary syndrome, carcinogenesis, and hyperplasia. The compounds of Formula I are also useful for treating physiological disorders related to, for example, cell differentiation to produce lipid accumulating cells, regulation of insulin sensitivity and blood glucose levels, which are involved in, for example, abnormal pancreatic beta cell function, insulin secreting tumors and/or autoimmune hypoglycemia due to autoantibodies to insulin, autoantibodies to the insulin receptor, or autoantibodies that are stimulatory to pancreatic beta cells), macrophage differentiation which leads to the formation of atherosclerotic plaques, inflammatory response, carcinogenesis, hyperplasia, adipocyte gene expression, adipocyte differentiation, reduction in the pancreatic β -cell mass, insulin secretion, tissue sensitivity to insulin, liposarcoma cell growth, polycystic ovarian disease, chronic anovulation, hyperandrogenism, progesterone production, steroidogenesis, redox potential and oxidative stress in cells, nitric oxide synthase (NOS) production, increased gamma glutamyl transpeptidase, catalase, plasma triglycerides, HDL, and LDL cholesterol levels, and the like.

Particularly useful compounds of Formula I of the present invention are those with efficacy in lowering blood glucose concentration and serum triglyceride levels, and raising serum HDL cholesterol levels.

Therefore, the compounds of Formula I of this invention are expected to be valuable as therapeutic agents. Accordingly, an embodiment of this invention includes a method of treating the various conditions identified above in a patient (including mammals) which comprises administering to said patient a composition containing an amount of the compound of Formula I that is effective in treating the target condition.

As indicated above, a compound of Formula I may be administered alone or in combination with one or more additional hypoglycemic agents. Combination therapy includes administration of a single pharmaceutical dosage formulation which contains a compound of Formula I and one or more additional hypoglycemic agent, as well as

administration of the compound of Formula I and each additional hypoglycemic agents in its own separate pharmaceutical dosage formulation. For example, a compound of Formula I and hypoglycemic agent may be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent may be administered in separate oral dosage formulations.

Where separate dosage formulations are used, the compound of Formula I and one or more additional hypoglycemic agents may be administered at essentially the same time (e.g., concurrently) or at separately staggered times (e.g., sequentially).

For example, the compound of Formula I may be administered in combination with one or more of the following additional hypoglycemic agents: insulin; biguanidines such as metformin or buformin; sulfonylureas such as acetohexamide, chloropropamide, tolazamide, tolbutamide, glyburide, glipizide, glyclazide; or any other insulin secretagogue such as, for example, repaglinide and nateglinide; α -glycosidase inhibitors such as acarbose, voglibose, or miglitol; or β_3 -adrenoreceptor agonists such as CL-316,243.

The compounds of Formula I may also be utilized, in free base form or in compositions, as well as in research and diagnostics or as analytical reference standards, and the like, which are well known in the art. Therefore, the present invention includes compositions which are comprised of an inert carrier and an effective amount of a compound of Formula I, or a salt, or ester thereof. An inert carrier is any material which does not interact with the compound to be carried and which lends support, means of conveyance, bulk, traceable material, and the like to the compound to be carried. An effective amount of the compound is that amount which produces a result or exerts an influence on the particular procedure being performed.

In another aspect, the present invention provides a method for treating a disease state in a patient, wherein the disease is associated with a physiological detrimental level of insulin, glucose, free fatty acids (FFA), cholesterol, or triglycerides in the blood, comprising administering to the patient a therapeutically effective amount of a compound of Formula I. In a further embodiment, the present invention provides a method for treating a disease state in a patient, wherein the disease is associated with a physiological detrimental level of insulin, glucose, free fatty acids (FFA), or triglycerides in the blood, comprising administering to the patient a therapeutically effective amount of a compound of Formula I and also administering a therapeutically effective amount of an additional hypoglycemic agent such as, for example, insulin, a biguanidine compound, and the like.

Since sulfonylureas and other insulin secretagogues are known to be capable of stimulating insulin release, but are not capable of acting on insulin resistance, and

compounds of Formula I are able to act on insulin resistance, it is envisaged that a combination of these medicaments may be used as a remedy for conditions associated with both deficiency in insulin secretion and insulin resistance. Therefore, the invention also provides a method of treating Type II diabetes mellitus in a patient comprising administering a compound of Formula I and one or more additional hypoglycemic agents such as, for example, sulfonylureas, biguanidines, β -adrenoreceptor agonists, α -glycosidase inhibitors, and insulin. Also, compounds of Formula I may be used in combination with HMG Co-A reductase inhibitors (statins), bile acid binding resin, or fibric acid derivatives to improve the lipid profile of subjects with dyslipidemia and insulin resistance. Compounds of Formula I may also be used in combination with agents that regulate hypertension (e.g., inhibitors of angiotension converting enzyme (ACE), β -blockers, calcium channel blockers) and body weight of subjects with insulin resistance or type 2 diabetes.

The following specific examples are presented to illustrate the invention described herein, but should not be construed as limiting the scope of the invention in any way.

SPECIFIC EXAMPLES

HPLC-electrospray mass spectra (HPLC ES-MS) were obtained using a Hewlett-Packard 1100 HPLC equipped with a quaternary pump, a variable wavelength detector, a YMC Pro C18 2.0 mm x 23 mm column, and a Finnigan LCQ ion trap mass spectrometer with electrospray ionization. Gradient elution from 90% A to 95% B over 4 minutes was used on the HPLC. Buffer A was 98% water, 2% Acetonitrile, and 0.02% TFA, and Buffer B was 98% Acetonitrile, 2% water, and 0.018% TFA. Spectra were scanned from 140-1200 amu using a variable ion time according to the number of ions in the source.

Proton (^1H) nuclear magnetic resonance (NMR) spectra were measured with a General Electric GN-Omega 300 (300 MHz) spectrometer with either Me_4Si (δ 0.00) or residual protonated solvent (CHCl_3 δ 7.26; MeOH δ 3.30; DMSO δ 2.49) as standard. Carbon (^{13}C) NMR spectra were measured with a General Electric GN-Omega 300 (75 MHz) spectrometer with solvent (CDCl_3 δ 77.0; $\text{d}_3\text{-MeOD}$; δ 49.0; $\text{d}_6\text{-DMSO}$ δ 39.5) as standard.

Chiral separations were performed using a commercially available Chiracel® AD HPLC column, eluting with a gradient of isopropanol in hexane (from 1% to 15%) with addition of 0.1% trifluoroacetic acid.

ABBREVIATIONS AND ACRONYMS


When the following abbreviations are used herein, they have the following meaning:

Ac ₂ O	acetic anhydride
ADDP	1,1'-(azodicarbonyl)dipiperidine
anhy	anhydrous
BOC	<i>tert</i> -butoxycarbonyl
<i>n</i> -BuOH	<i>n</i> -butanol
<i>t</i> -BuOH	<i>tert</i> -butanol
<i>t</i> -BuOK	potassium <i>tert</i> -butoxide
CDI	carbonyl diimidazole
CD ₃ OD	methanol- <i>d</i> ₄
Celite [®]	diatomaceous earth filter agent, [®] Celite Corp.
CH ₂ Cl ₂	methylene chloride
CI-MS	chemical ionization mass spectroscopy
conc	concentrated
DCC	dicyclohexylcarbodiimide
DCM	dichloromethane
de	diastereomeric excess
DEAD	diethyl azodicarboxylate
dec	decomposition
DIA	diisopropyl amine
DIBAL-H	diisobutylaluminum hydroxide
DMAP	4-(<i>N,N</i> -dimethylamino)pyridine
DME	dimethoxyethane
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethylsulfoxide
EDCI	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
ee	enantiomeric excess
ELSD	evaporative light scattering detector
ES-MS	electrospray mass spectroscopy
EtOAc	ethyl acetate
EtOH	ethanol (100%)
EtSH	ethanethiol
Et ₂ O	diethyl ether
Et ₃ N	triethylamine
GC-MS	gas chromatography-mass spectroscopy

HPLC	high performance liquid chromatography
IPA	isopropylamine
LAH	lithium aluminum hydride
LC-MS	liquid chromatography-mass spectroscopy
LDA	lithium diisopropylamide
m/z	mass-to-charge ratio
MeCN	acetonitrile
NMM	4-methylmorpholine
Ph ₃ P	triphenylphosphine
Pd(dppf)Cl ₂	[1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II)
Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium(0)
Pd(OAc) ₂	palladium acetate
P(O)Cl ₃	phosphorous oxychloride
R _f	retention factor (TLC)
RT	retention time (HPLC)
rt	room temperature
TEA	triethyl amine
THF	tetrahydrofuran
TFA	trifluoroacetic acid
TLC	thin layer chromatography
TMAD	N,N,N',N'-tetramethylethylenediamine
TMSCI	trimethylsilyl chloride

Example 1

Preparation of methyl 2-(6-methoxy-1H-inden-3-yl) butanoate




An oven dried 5-L four-necked round-bottomed flask was fitted with a thermometer, a condenser, an addition funnel, and a mechanical stirrer. Under Ar protection, a suspension of 5-methoxy-1-indanone (80.0 g, 494 mmol), Zn powder (Lancaster, 56.2 g, 865 mmol) in 2 L anhydrous THF was stirred at 60°C (internal temperature), while a solution of methyl bromobutyrate (134.1 g, 741 mmol) in 400 mL anhydrous THF was added slowly through an addition funnel. After completion of the


addition, the reaction mixture was stirred at 60°C (internal temperature) for 1 hour. The reaction was followed by TLC analysis of aliquots following 1N aqueous HCl work-up. After the reaction was completed, it was cooled in an ice-water bath followed by slow addition of 3 L of 1N HCl solution. The pot temperature was kept below 20°C. The mixture was then extracted with 1 L EtOAc. The organic layer was washed with water until pH 6.0-7.0, then saturated NaCl solution, and dried over Na₂SO₄. The product (127 g, >99%), a yellow oil, was obtained after solvent removal and drying under vacuum. ¹H NMR (DMSO-*d*₆) δ 7.28(d, 1H), 7.05(d, 1H), 6.82(dd, 1H), 6.22(s, 1H), 3.72(s, 3H), 3.60(m, 1H), 3.58(s, 3H), 3.28(s, 2H), 1.95(m, 1H), 1.80(m, 1H), 0.88(t, 3H).

Example 2a


Preparation of 2-(6-methoxy-1*H*-inden-3-yl) butanoic acid



To a solution of the ester prepared in Example 1 (14.0 g, 58.9 mmol) in 140 mL MeOH, was added a solution of KOH (6.4 g, 113.8 mmol) in 5 mL water. The reaction mixture was stirred at 60°C (pot temperature) for 2 hours. TLC showed 70% conversion. A solution of KOH (3.0 g, 53.6 mmol) in 100 mL water was then slowly added to the pot. After 1 hour, the reaction was completed. After cooling to room temperature, the solvents were removed at a reduced pressure. The residue was dissolved in 500 mL water, and then washed with EtOAc. The aqueous layer was cooled in an ice-water bath, and then acidified with conc. HCl to pH<3.0. The product was extracted into 300 mL CH₂Cl₂, washed with water (2 x 100 mL), then dried over Na₂SO₄. After Na₂SO₄ was filtered off, the CH₂Cl₂ solution was stirred with 3.0 g of charcoal for 2 hours. The charcoal was removed by filtration through a pad of Celite®. The title product (12.5 g, 95%) was obtained as a light brown solid after solvent removal and vacuum drying. ¹H NMR (DMSO-*d*₆) δ 12.20(b, 1H), 7.30(d, 1H), 7.06(d, 1H), 6.82(dd, 1H), 6.22(s, 1H), 3.75(s, 3H), 3.45(t, 1H), 3.30(s, 2H), 1.90(m, 1H), 1.78(m, 1H), 0.90(t, 3H).

Example 2b**Preparation of 2-(6-methoxy-1H-inden-3-yl) propanoic acid**

This substrate was prepared using the same procedures as described for Examples 1 and 2a, starting with 5-methoxy-1-indanone and methyl 2-bromopropionate. Yield: 68%. ^1H NMR (CD_2Cl_2) δ 7.34 (d, $J = 9$, 1H), 7.07 (d, $J = 2$, 1H), 6.85 (dd, $J = 9$, $J = 2$, 1H), 6.32 (m, 1H), 3.82 (m, 4H), 3.36 (m, 2H), 1.56 (d, $J = 7$, 3H).

Example 3**Preparation of (2S)-2-(6-methoxy-1H-inden-3-yl)butanoic acid**

To a solution of the racemic indene acid prepared in Example 2a (300 g, 1.29 mol) in 4.5 L CH_3CN , was added quinine (324 g, 1.0 mol) at rt. The mixture was stirred for 1 hour, and became a solution. A small amount of insoluble particles was removed by filtration through a microfiber filter under vacuum. The filtrate was then mechanically stirred under Ar over night. After 24 hours, a small sample of solid was taken and analyzed, showing 76.2% ee. The agitation was stopped after two more days. The suspension was filtered. The filter cake was washed with CH_3CN (3 x 200 mL), and then dried under vacuum at 40°C for 3 hours. This solid was stirred with 4.5 L CH_3CN at 70°C until all solids went into solution. The solution was allowed to cool down to rt slowly. The resulting suspension was stirred at rt for 24 hours. The suspension was filtered. The filter cake was washed with CH_3CN (3 x 250 mL), and then dried under vacuum at 40°C for 24 hours. This quinine salt was collected as a white solid (254.6 g, 35.4% yield, 96.8% ee).


The quinine salt (544.3 g, 0.98 mol) was dissolved in 4.0 L CH_2Cl_2 to obtain a clear solution. It was stirred vigorously with 4.0 L of 2N HCl solution in a 22-L round-bottomed flask with a bottom valve. After 30 minutes, the mixture was allowed to settle. The bottom layer was separated and top aqueous layer was extracted with 1 L CH_2Cl_2 . The combined CH_2Cl_2 layers were washed with water (3 x 2.0 L) until pH 5.0-6.0, and then

dried over Na_2SO_4 . The product (230.8 g, 99%, 96.8% ee) was obtained as an off white solid after solvent removal and vacuum drying. ^1H NMR was identical to that of the racemic material described in Example 2a.

Treatment of the mother liquor in similar fashion gave the (*R*) isomer. Alternatively, the mother liquor may be subjected to aqueous basic conditions in order to effect racemization and recovery of racemic starting material.

Example 4


Preparation of (2*S*)-2-[(1*S*)-5-methoxy-2,3-dihydro-1*H*-inden-1-yl]butanoic acid



A solution of the product obtained in Example 3 (105 g, 453 mmol), $\text{ClRh}(\text{PPh}_3)_3$ (21.0 g, 5% eq.) and triethylamine (68.8 g, 679.5 mmol) in EtOH (945 mL) and THF (105 mL) was shaken in a 2-L pressure bottle under 60 psi H_2 for 16 hours. The solvents were removed at a reduced pressure. The resulting mixture was stirred in 1.5 L of 1N HCl solution and 1.5 L CH_2Cl_2 . The aqueous layer was extracted with CH_2Cl_2 (2 x 250 mL). The combined CH_2Cl_2 layers were washed with 1 L of 1N HCl solution and stirred with 1 L of 1N NaOH solution. The organic layer was extracted with 1N NaOH solution (2 x 0.5 L). The combined aqueous layer was washed with CH_2Cl_2 (2 x 250 mL), and acidified (pH 2.0-3.0) by a slow addition of conc. HCl solution at below 15°C . The acidic mixture was extracted with CH_2Cl_2 (2 x 1.5 L), and washed with water (2 x 0.5 L) until pH 5.0-6.0. After washing with brine and drying over anhydrous Na_2SO_4 , solvent was evaporated under a reduced pressure. The product (101.0 g, 95% yield, 96.8% ee) was obtained as a light yellow oil. ^1H NMR ($\text{DMSO}-d_6$) δ 12.20(s, 1H), 7.04(d, 1H), 6.78(d, 1H), 6.66(dd, 1H), 3.70(s, 3H), 3.28(m, 1H), 2.72(m, 2H), 2.32(m, 1H), 2.06(m, 1H), 1.80(m, 1H), 1.50(m, 1H), 1.36(m, 1H), 0.82(t, 3H).

Example 5a

Preparation of *syn*-2-[5-methoxy-2,3-dihydro-1*H*-inden-1-yl]butanoic acid



A suspension of racemic indene acid (Example 2, 980 mg, 4.2 mmol), $\text{ClRh}(\text{PPh}_3)_3$ (139 mg, 0.15 mmol), NaHCO_3 (378 mg, 4.5 mmol) in EtOH (20 mL), and H_2O (10 mL) was shaken in a 500 mL pressure bottle under 60 psi H_2 for 30 hours. Additional $\text{ClRh}(\text{PPh}_3)_3$ (300 mg, 0.33 mmol) was added to the reaction mixture and hydrogenation was continued for 3 more days. After this time, EtOH was removed at a reduced pressure and the residue was diluted with 200 mL water. The black solid was removed by filtration and the filtrate was washed with EtOAc (2 x 200 mL). The aqueous solution was then acidified with conc. HCl, and extracted with CH_2Cl_2 (2 x 100 mL). The combined CH_2Cl_2 layer was washed with brine and dried over Na_2SO_4 . Removal of the solvent in vacuum afforded the indane acid as light yellow oil (600 mg, 60%). The product mixture resulted a diastereomeric mixture (87:13) in favor of the *syn* isomers as determined by NMR analysis, using the ratio of integration of NMR peaks δ 7.11(d, 1H) for the *anti*, and δ 7.03(d, 1H) for the *syn* isomers.


Resolution of the product into optical isomers may be accomplished as follows: to a mechanically stirred solution of the *syn* indane acetic acid [(2*R*, 1*R*) and (2*S*, 1*S*), 14.69 g, 62.7 mmol] in acetonitrile (290 mL) at rt, was added (*R*)-(+)- α -methylbenzylamine (8.49 mL, 65.9 mmol) in one portion. The resulting mixture was stirred overnight. Little solid formation was observed. The reaction mixture was concentrated to dryness and the residue was redissolved in acetonitrile (200 mL) with heating. Magnetic stirring was begun to initiate precipitation. The mixture was stirred overnight. The solids were collected by filtration, and washed three times with a small amount of cold acetonitrile. The solids were then dried under vacuum for 1.5 hours (8.1 g, 86% ee). The slightly wet solids were recrystallized in acetonitrile (120 mL) to give 6.03 g of the (2*S*)-2-[(1*S*)-5-methoxy-2,3-dihydro-1*H*-inden-1-yl]butanoic acid, (*R*)- α -methylbenzylamine salt (94.4% ee). A second crop was collected from various filtrates (0.89 g, 97.6% ee). The overall yield of resolution was 31% (62% based on the maximum content of (2*S*, 1*S*) acid in the racemate). The material was identical to that obtained in Example 4.

Optical purity for this Example and that of Example 4 may also be analyzed by chiral HPLC; Column: Chiracel AD, 4.6 (I.D.) x 250 mm; Mobile Phase, A: 0.1% TFA (trifluoroacetic acid) in hexanes, B: 0.1% TFA in IPA (isopropyl alcohol); Method, Isocratic 95%A (5%B), 20 min.; Flow Rate, 1.5 mL/min.; Detector (UV), 284 nm. Retention times for the four possible diastereomers are 5.163 min. (2*S*, 1*R*), 6.255 min. (2*R*, 1*S*), 10.262 min. (2*R*, 1*R*) and 14.399 min. (2*S*, 1*S*). The first locator (2*S* or 2*R*) denotes the absolute configuration of the carbon adjacent to the carboxyl group (the 2-position); the second locator (1*S* or 1*R*) denotes the absolute configuration of the indane ring carbon (its 1-position).

The stereochemical assignment for each peak was determined by chiral HPLC analysis of a non-equal (syn/anti) racemic diastereomeric mixture of compound 5, which provided four baseline-resolved peaks. Peaks 3 and 4, and peaks 1 and 2 represented enantiomer pairs, based on UV integration. The absolute configuration of the compound of peak 4 was determined to be 2*S*, 1*S* by X-ray structural analysis. Peak 3, the corresponding enantiomer, was then assigned a 2*R*, 1*R* configuration with certainty. Peak 1 was assigned to the (2*S*, 1*R*)-diastereomer (retention time: 5.363 min., ca. 0.97% area) by comparison to the minor product obtained from the reduction of the chiral acid (Example 3) as described in Example 4. The remaining peak 2, could then be assigned with certainty to the compound with 2*R*, 1*S* configuration.

Example 5b


Preparation of *syn*-2-[5-methoxy-2,3-dihydro-1*H*-inden-1-yl]propanoic acid



The compound was prepared in 71% yield and >99% de using the same procedure as described for Example 4 starting with (racemic) Example 2b: ¹H NMR (DMSO-*d*₆) δ 12.18 (s, 1H), 7.03 (d, *J* = 8, 1H), 6.75 (d, *J* = 2, 1H), 6.67 (dd, *J*₁ = 8, *J*₂ = 2, 1H), 3.68 (s, 3H), 3.37 (m, 1H), 2.72 (m, 3H), 2.03 (m, 1H), 1.75 (m, 1H), 0.89 (d, *J* = 7, 3H); ¹³C NMR (CD₂Cl₂) δ 12.626, 28.228, 31.950, 43.300, 46.445, 55.607, 110.054, 112.510, 124.552, 136.702, 146.411, 159.464, 182.330.

Example 6


Preparation of methyl (2S)-2-[(1S)-5-methoxy-2,3-dihydro-1H-inden-1-yl]butanoate



A suspension of acid prepared in Example 4 (220.0 g, 0.94 mol), NaHCO_3 (237.0 g, 2.82 mol), CH_3I (200 g, 1.41 mol) in 2.0 L DMF was stirred under Ar at rt for 18 hours. NMR analysis showed 95% reaction. Adding CH_3I (100 g), and stirring for additional 24 hours at rt caused completion of the reaction. The reaction mixture was poured into 4.0 L water, and extracted with EtOAc (2 x 2 L). The organic layer was sequentially washed with water (2 x 1 L), 1 L of 1N NaOH solution, water (2 x 1 L), and 500 mL brine, and dried over Na_2SO_4 . The product (233 g, 99%) was obtained as a light yellow oil after solvent removal and vacuum drying. ^1H NMR ($\text{DMSO}-d_6$) δ 6.90(d, 1H), 6.78(d, 1H), 6.66(dd, 1H), 3.70(s, 3H), 3.60(s, 3H), 3.20(m, 1H), 2.80(m, 2H), 2.40(m, 1H), 2.08(m, 1H), 1.80(m, 1H), 1.58(m, 1H), 1.40(m, 1H), 0.80(t, 3H).

Example 7

Preparation of methyl (2S)-2-[(1S)-5-hydroxy-2,3-dihydro-1H-inden-1-yl]butanoate




To a cold solution (ice water bath) of the compound prepared in Example 6 (233 g, 0.94 mol) in 2.5 L CH_2Cl_2 , was added AlCl_3 (630 g, 4.7 mol) slowly under Ar. The pot temperature was kept below 20°C , and the color of the reaction turned purple. EtSH (345 mL, 4.7 mol) was added slowly via an addition funnel to the reaction mixture, and the inside temperature was kept below 15°C . After 2 hours of stirring at below 20°C , the reaction went to completion by NMR analysis. The pot mixture was slowly poured into 2.5 L ice water with a strong agitation. The organic layer was separated, and the aqueous layer was extracted with 1L CH_2Cl_2 . The combined CH_2Cl_2 layers were washed with water (4 x 1 L) until pH 6.0-7.0, and then dried over Na_2SO_4 . The product (216 g, 98%) was obtained as a white solid after solvent removal and vacuum drying. ^1H NMR

(DMSO- d_6) δ 9.10(s, 1H), 6.78(d, 1H), 6.58(d, 1H), 6.50(dd, 1H), 3.60(s, 3H), 3.20(q, 1H), 2.70(m, 2H), 2.40(m, 1H), 2.08(m, 1H), 1.80(m, 1H), 1.50(m, 2H), 0.80(t, 3H).

Example 8


Preparation of methyl 3-[(4-methylbenzoyl)amino]-4-oxopentanoate




To a suspension of L-aspartic acid β -methyl ester hydrochloride (250 g, 1.36 mol) in chilled ($<5^{\circ}\text{C}$) CH_2Cl_2 (4 L) was added Et_3N (440 g, 4.35 mol) in a steady flow followed by a slow addition of Me_3SiCl (324 g, 2.99 mol). The mixture was warmed to 25°C and stirred for one hour, cooled again ($<10^{\circ}\text{C}$), and *p*-toluoyl chloride (205 g, 1.36 mol) was added dropwise. The mixture was allowed to warm to ambient slowly with stirring for 16 hours. The reaction mixture was then diluted with CH_2Cl_2 (500 mL) and washed with 1N HCl (500 mL), brine (500 mL), and dried over Na_2SO_4 . The resultant amide product (310 g, 91%), a white solid, was obtained after solvent removal and drying under vacuum. It was then dissolved in pyridine (1.25 L) and DMAP (5 g) was added. Acetic anhydride (840 mL) was added slowly and then the reaction was heated at 90°C for 2 hours. The cooled solution was poured into 7 L ice water and extracted with 6 L EtOAc. The organic layer was washed with 2N HCl (3 x 1 L) and 1N NaOH (1 L), dried over MgSO_4 and concentrated to afford the title compound as a white solid (301 g, 93%).

Example 9


Preparation of methyl [5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]acetate



The intermediate prepared in Example 8 (280 g, 1.06 mol) was dissolved in acetic anhydride (650 mL) followed by slow addition of conc. H_2SO_4 (60 mL). The pot temperature reached 80°C . The reaction was then held at 85°C for 1 hour, cooled, and the acetic anhydride removed *in vacuo*. The residue was poured into ice water (2 L) and extracted with EtOAc (4 L total). The organic layer was then stirred with 1 N NaOH (500 mL) for 1 hour, separated, then dried with MgSO_4 and concentrated to afford the title ester as a clear oil (223 g, 87%), which slowly solidified to a white solid.

Example 10**Preparation of 2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethanol**

The oxazole ester prepared in Example 9 (300 g, 1.22 mol) was dissolved in THF (2.7 L) and solid LiBH_4 (26.6 g, 1.22 mol) was added in 5-g portions while maintaining temperature below 45°C . Reaction was complete within an hour after addition. Solvent was reduced to half volume and then poured into ice water (3 L). The mixture was then acidified by slowly adding 1 N HCl (1 L). A white precipitate formed and was collected by filtration and oven dried over P_2O_5 to give the desired oxazole ester (214 g, 83%).


Example 11**Preparation of methyl (2S)-2-((1S)-5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)butanoate**

A suspension of the hydroxyindane carboxylate prepared in Example 7 (208 g, 889 mmol), oxazole alcohol prepared in Example 10 (212 g, 977 mmol), ADDP (335 g, 1.33 mol), Ph_3P (348 g, 1.33 mol) in 6.0 L anhydrous THF was stirred at rt under Ar. The reaction was followed by ^1H NMR. No further progress was observed after 2 days. After solids were removed by filtration, THF was removed under reduced pressure. The remaining mixture was stirred in 3 L of 50/50 mixture EtOAc/hexane for 10 minutes, and more solids were formed and removed by filtration. The filtrate was concentrated and subjected to the same procedure with 25/75 mixture of EtOAc/hexane. After solvents were removed, the resulting oily mixture was purified on a silica gel (3.0 kg) column using CH_2Cl_2 (10.0 L) and 20% $\text{CH}_3\text{CN}/\text{CH}_2\text{Cl}_2$ (10.0 L) as solvent. Fractions containing product were collected, and then concentrated. The crude mixture was dissolved in 4.0 L CH_2Cl_2 , and the unreacted hydroxy compound was removed by washing with 1N NaOH (3 x 1 L). The CH_2Cl_2 layer was dried over Na_2SO_4 . The product (358 g, 93%) was obtained as a light yellow oil after solvent removal and vacuum drying. ^1H NMR (DMSO-


d_6) δ 7.78(d, 2H), 7.30(d, 2H), 6.90(d, 1H), 6.75(d, 1H), 6.65(dd, 1H), 4.15(t, 2H), 3.60(s, 3H), 3.25(q, 1H), 2.90(t, 2H), 2.75(m, 2H), 2.40(m, 1H), 2.35(s, 3H), 2.32(s, 3H), 2.05(m, 1H), 1.80(m, 1H), 1.50(m, 2H), 0.80(t, 3H).

Example 12

Preparation of (2S)-2-((1S)-5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)butanoic acid



To a solution of LiOH (90.4 g, 3.76 mol) in 1.3 L water and 1.3 L MeOH, was added a solution of the ester prepared in Example 11 (325 g, 0.75 mol) in 3.9 L THF at rt. The solution turned cloudy. This mixture was heated at 60°C (pot temperature) for 4 hours, and TLC (50% EtOAc/hexane) analysis showed ca. 50% conversion. A solution of LiOH (30.1 g, 1.25 mol) in water (200 mL) was added to the reaction mixture. After 2 hours, TLC analysis showed ca. 85% reaction. Again, a solution of LiOH (30.1 g, 1.25 mol) in water (200 mL) was added to the reaction mixture. After 2 hours, TLC analysis showed very little starting ester left. After the reaction mixture was cooled to rt, THF and MeOH were removed at a reduced pressure. The residue was diluted with water until the solids dissolved (a total of 60 L of water used). Conc. HCl solution was slowly added to this aqueous solution until pH 2.0-3.0. The solid was collected by filtration, and dried under house vacuum overnight. This solid was stirred with 15 L EtOAc and 2 L of 1N HCl solution for 30 minutes. The EtOAc layer was separated and washed with 1N HCl solution (2 x 1 L). The organic phase was then washed with water (4 x 2 L) until pH = 5.0-6.0. Under Ar protection, the EtOAc solution was reduced to 2.5 L by normal pressure distillation, then cooled to rt without disturbance. White solid precipitated out. After further cooling in an ice water bath for 2 hours, the solid was filtrated and washed with 500 mL cold EtOAc. After drying under high vacuum at 35°C to a constant weight, the final product (266 g, 81%, 98% ee,) was collected as a white crystal. ^1H NMR (CDCl_3) δ 7.82(d, 2H), 7.20(d, 2H), 7.05(d, 1H), 6.75(d, 1H), 6.70(dd, 1H), 4.20(t, 2H), 3.42(q, 1H), 2.95(t, 2H), 2.80(m, 2H), 2.50(m, 1H), 2.35(s, 3H), 2.32(s, 3H), 2.20(m, 1H), 1.90(m, 1H), 1.65(m, 1H), 1.45(m, 1H), 0.90(t, 3H). Chiral purity, 99% ee, $[\alpha]_D^{25} = +16.11(\text{CHCl}_3)$, mp 149.5-150.5°C.

Example 13**Preparation of 2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid**

Step 1. To a solution of 5-methoxy-indanone (10 g) dissolved in toluene (150 mL) was added AlCl_3 (15 g). The mixture was refluxed for 4 hours until a precipitate appeared. The resulting mixture was cooled and poured into ice water (150 mL). The precipitate was filtered and washed with water, then air-dried to give the desired product (8.5 g, 90%).

Step 2. Benzyl bromide (17 g), 5-hydroxyl-indanone (15 g), K_2CO_3 (20 g), and 200 mL acetone were mixed in a round-bottom flask (500 mL). The mixture was refluxed

for 1 hour. The K_2CO_3 was filtered off, and the filtrate was evaporated. The resulting residue was crystallized from EtOAc to give 18 g product (75%).

Step 3. A solution of 5-benzyloxyl-indanone (1.14 g, 4.79 mmol) and diethyl malonate (0.844 g, 5.29 mmol) in THF (20 mL) was cooled to 0°C under argon, and $TiCl_4$ (10 mL, 1M in CH_2Cl_2) was added dropwise. Pyridine (2 mL) was added finally. The resulting mixture was stirred overnight at rt. After filtration, EtOAc (30 mL) was added into the filtrate. The organic layer was washed with brine (20 mL x 3), dried with Na_2SO_4 , and evaporated. The residue was separated by silica gel chromatography to give 800 mg product (50%).

Step 4. The product of step 3 (1.7 g) was dissolved in MeOH (25 mL), and Pd-C (300 mg) was added as a slurry in MeOH, and placed under 60 psi H_2 in a Parr shaker for 6 hours. After filtration and concentration, 1.2 g product was obtained (92%).


Step 5. $P(Ph)_3$ (420 mg) and ADDP (420 mg) were dissolved in THF (5 mL) at 0°C, and stirred for 10 minutes. A THF solution of oxazole (300 mg) and phenol (430 mg) was added to the flask. The resulting mixture was stirred for 24 hours, and filtered. The filtrate was evaporated and the resulting residue was separated by silica gel chromatography to give product (320 mg, 45%).

Step 6. The intermediate prepared in step 5 (160 mg) was dissolved in THF (5 mL), and iodoethane (0.5 mL) and *t*-BuOK (50 mg) were added to the solution and stirred overnight. After filtration, the product was separated by using TLC, providing 100 mg (65%).


Step 7. The intermediate prepared in step 6 (30 mg) was dissolved in DMSO (1 mL). LiCl (160 mg) was added into the flask. The mixture was refluxed for 5 hours. From the resulting mixture, the product was separated by TLC, giving 13 mg (52%).

Step 8. The intermediate prepared in step 7 was subjected to hydrolysis in aqueous KOH as described for Example 2 to obtain the product: LC-MS, RT 3.57 min., $M+1$ 406; 1H NMR (CD_2Cl_2): δ 0.93 (t, 3H), 1.40-1.70 (m, 2H), 1.80-2.20 (m, 2H), 2.30 (s, 3H), 2.40 (m, 1H), 2.60-2.80 (m, 2H), 2.90 (t, 2H), 3.20-3.40 (m, 1H), 4.10 (t, 2H), 6.60 (dd, 1H), 6.70 (d, 1H), 7.00 (d, 1H), 7.30 (m, 3H), 7.90 (m, 2H).


By using the procedures from Examples 1-13 together in some cases with the chiral HPLC separation method described in the general section, and by using the appropriate starting materials, the following were prepared and characterized in a similar manner:

Example 14**2-(5-[2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl]butanoic acid**


LC-MS, RT 3.70 min., M+1 420; ^1H NMR (CD_2Cl_2): δ 0.93 (t, 3H), 1.40-1.70 (m, 2H), 1.80-2.20 (m, 2H), 2.30 (s, 3H), 2.35 (s, 3H), 2.40 (m, 1H), 2.60-2.80 (m, 2H), 2.90 (t, 2H), 3.20-3.40 (m, 1H), 4.10 (t, 2H), 6.60 (dd, 1H), 6.70 (d, 1H), 7.00 (d, 1H), 7.20 (m, 3H), 7.80 (m, 2H).

Example 15**(2S)-2-[(1S)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl]butanoic acid**


The enantiomer was isolated by chiral HPLC. LC-MS, RT 3.57 min., M+1 406; ^1H NMR (CD_2Cl_2): δ 0.93 (t, 3H), 1.48 (ddq, 1H), 1.68 (ddq, 1H), 1.93 (dddd, 1H), 2.18 (dddd, 1H), 2.34 (s, 3H), 2.50 (ddd, 1H), 2.77 (ddd, 1H), 2.87 (ddd, 1H), 2.96 (t, 2H), 3.42 (ddd, 1H), 4.19 (t, 2H), 6.68 (dd, 1H), 6.77 (d, 1H), 7.08 (d, 1H), 7.42 (m, 2H), 7.44 (m, 1H), 7.97 (dd, 2H). ^{13}C NMR: δ 10.4, 12.4, 22.4, 26.6, 29.5, 31.8, 46.5, 51.8, 67.2, 110.9, 113.0, 124.7, 126.2, 128.1, 129.1, 130.2, 133.2, 137.1, 145.6, 146.3, 158.7, 159.7, 180.4.

Example 16**(2S)-2-((1R)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl)butanoic acid**

The enantiomer was isolated by chiral HPLC. LC-MS, RT 3.57 min., M+1 406; ¹H NMR (CD₂Cl₂): δ 0.93 (t, 3H), 1.61 (ddq, 1H), 1.69 (ddq, 1H), 1.99 (dddd, 1H), 2.19 (dddd, 1H), 2.47 (s, 3H), 2.52 (ddd, 1H), 2.73 (ddd, 1H), 2.89 (ddd, 1H), 3.11 (t, 2H), 3.31 (ddd, 1H), 4.21 (t, 2H), 6.66 (dd, 1H), 6.74 (d, 1H), 7.13 (d, 1H), 7.55 (m, 2H), 7.61 (m, 1H), 8.05 (dd, 2H). ¹³C NMR: δ 10.5, 12.2, 23.8, 24.8, 30.3, 31.5, 46.4, 50.9, 66.1, 110.8, 112.6, 125.9, 127.4, 123.6, 129.8, 133.3, 129.7, 137.0, 148.4, 146.5, 158.2, 160.5, 181.0.

Example 17**(2R)-2-((1R)-5-[2-(5-methyl-2-[4-methylphenyl]-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl)butanoic acid**


The enantiomer was isolated by chiral HPLC. LC-MS, RT 3.70 min., M+1 420; ¹H NMR (CD₂Cl₂): δ 0.95 (t, 3H), 1.40 (m, 1H), 1.70 (m, 1H), 1.90 (m, 1H), 2.20 (m, 1H), 2.30 (s, 3H), 2.35 (s, 3H), 2.50 (m, 1H), 2.60-2.80 (m, 2H), 2.90 (t, 2H), 3.40 (dd, 1H), 4.20 (t, 2H), 6.60 (dd, 1H), 6.70 (d, 1H), 7.10 (d, 1H), 7.20 (m, 3H), 7.80 (m, 2H).

Example 18**2-(5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl)propanoic acid**

LC-MS, RT 3.41 min., M+1 392; ^1H NMR (CD_2Cl_2): δ 1.10 (d, 3H), 1.90 (m, 2H), 2.20 (m, 1H), 2.40 (s, 3H), 2.70-3.00 (m, 2H), 2.95 (t, 2H), 3.45 (m, 1H), 4.20 (t, 2H), 6.70 (dd, 1H), 6.80 (d, 1H), 7.10 (d, 1H), 7.45 (m, 3H), 8.00 (m, 2H).

Example 19


2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}malonic acid



LC-MS, RT 3.00 min., M+1 422; ^1H NMR (CD_2Cl_2): δ 1.90 (m, 2H), 2.40 (t, 3H), 2.60-3.00 (m, 3H), 3.40 (t, 2H), 3.70 (m, 1H), 4.20 (t, 2H), 6.60 (dd, 1H), 6.80 (d, 1H), 7.10 (d, 1H), 7.50 (m, 3H), 7.95 (m, 2H).

Example 20


3-ethoxy-2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}-3-oxopropanoic acid



LC-MS, RT 3.39 min., M+1 450; ^1H NMR (CD_2Cl_2): δ 1.20 (t, 3H), 2.00 (m, 1H), 2.30 (m, 1H), 2.40 (s, 3H), 2.90 (m, 2H), 3.10 (t, 2H), 3.80 (m, 1H), 4.20 (t & q, 4H), 6.70 (dd, 1H), 6.80 (d, 1H), 7.10 (d, 1H), 7.50 (m, 3H), 8.00 (m, 2H).

Example 21


2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}-5-phenylpentanoic acid



LC-MS, RT 3.98 min., M+1 396; ^1H NMR (CD_2Cl_2): δ 1.40-1.80 (m, 4H), 1.90-2.20 (m, 2H), 2.35 (s, 3H), 2.40-3.00 (m, 5H), 2.90 (t, 2H), 3.35 (m, 1H), 4.10 (t, 2H), 6.60 (dd, 1H), 6.70 (d, 1H), 6.90-7.20 (m, 6H), 7.30 (m, 3H), 7.95 (m, 2H).

Example 22


2-(5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)propanoic acid




LC-MS, RT 3.52 min., M+1 406; ^1H NMR (CD_2Cl_2): δ 1.10 (d, 3H), 1.90 (m, 2H), 2.20 (m, 1H), 2.30 (s, 3H), 2.31 (s, 3H), 2.70-3.00 (m, 2H), 2.95 (t, 2H), 3.40 (m, 1H), 4.10 (t, 2H), 6.60 (dd, 1H), 6.70 (d, 1H), 7.00 (d, 1H), 7.20 (d, 2H), 7.80 (d, 2H).

Example 23


2-(5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)hexanoic acid




LC-MS, RT 3.92 min., M+1 448; ^1H NMR (CD_2Cl_2): δ 0.93 (t, 3H), 1.10-1.30 (m, 4H), 1.40-1.70 (m, 2H), 1.80-2.20 (m, 2H), 2.30 (s, 3H), 2.31 (s, 3H), 2.40 (m, 1H), 2.60-2.80 (m, 2H), 2.90 (t, 2H), 3.20-3.40 (m, 1H), 4.10 (t, 2H), 6.60 (dd, 1H), 6.70 (d, 1H), 7.00 (d, 1H), 7.20 (d, 2H), 7.80 (d, 2H).

Example 24**4-methyl-2-(5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)pentanoic acid**


LC-MS, RT 4.00 min., M+1 448; ^1H NMR (CD_2Cl_2): δ 0.93 (m, 6H), 1.20 (m, 1H), 1.40-1.70 (m, 2H), 1.80-2.20 (m, 2H), 2.30 (s, 3H), 2.31 (s, 3H), 2.40 (m, 1H), 2.60-2.80 (m, 2H), 2.90 (t, 2H), 3.20-3.40 (m, 1H), 4.10 (t, 2H), 6.60 (dd, 1H), 6.70 (d, 1H), 7.00 (d, 1H), 7.40 (d, 2H), 8.40 (d, 2H).

Example 25**4-methyl-2-(5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)-4-pentenoic acid**


LC-MS, RT 3.74 min., M+1 446; ^1H NMR (CD_2Cl_2): δ 1.60 (s, 3H), 1.70 (m, 2H), 1.80-2.20 (m, 2H), 2.30 (s, 3H), 2.31 (s, 3H), 2.40 (m, 1H), 2.60-2.80 (m, 2H), 2.90 (t, 2H), 3.20-3.40 (m, 1H), 4.10 (t, 2H), 5.60 (m, 2H), 6.60 (dd, 1H), 6.70 (d, 1H), 7.00 (d, 1H), 7.20 (d, 2H), 7.80 (d, 2H).

Example 26**Preparation of 2-{6-chloro-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid****via****2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethyl methanesulfonate and methyl 2-(6-chloro-5-hydroxy-2,3-dihydro-1H-inden-1-yl)butanoate**


Step 1. To a solution of 2-phenyl-4-methyl-5-hydroxyethyloxazole (500 mg, 2.5 mmol) in 12.5 mL THF, was added methanesulfonyl chloride (0.21 mL, 2.75 mmol) and triethylamine (0.42 mL, 3 mmol). The reaction solution was stirred at rt under argon for two hours then concentrated *in vacuo*. The resulting residue was taken up in ethyl acetate, washed with 1% aqueous hydrochloric acid (three times) and brine. It was then dried over sodium sulfate, filtered, and concentrated *in vacuo* to provide (617 mg, 88%): ES-MS m/z 282 ((M+H)⁺); HPLC RT 2.67; ¹H NMR (d₆-DMSO) δ 2.33 (s, 3H), 2.89 (t, 2H), 3.13 (s, 3H), 4.41 (t, 2H), 7.47-7.51 (m, 3H), 7.88-7.91 (m, 2H).



Step 2. Sulfuryl chloride (0.035 mL, 0.43 mmol) was added to a solution of methyl-5-hydroxy-2,3-dihydro-1-(2-butanoate) (100 mg, 0.43 mmol) in 2.15 mL acetic acid. The reaction solution was stirred at rt for 30 minutes, then concentrated *in vacuo*. The resulting residue was taken up in ethyl acetate and washed with water, saturated aqueous sodium bicarbonate, and brine. It was then dried over sodium sulfate, filtered, and concentrated *in vacuo* to provide 63 mg of the desired intermediate as a crude yellow oil which was carried on without further purification: GC-MS 269, ((M+H)⁺); GC RT (min.) 8.71; ¹H NMR (d₆-DMSO) δ 0.81 (t, 3H), 1.40-1.63 (m, 2H), 1.77-1.88 (m, 1H), 2.00-2.15 (m, 1H), 2.40-2.80 (m, 3H), 3.15-3.22 (m, 1H), 3.50 (s, 3H), 6.76 (s, 1H), 7.13 (s, 1H), 9.84 (s, 1H).




Step 3. A solution of the product obtained in step 2 (30.5 mg, 0.12 mmol) in 0.6 mL DMF was cooled to 0°C in an ice bath. A 60% dispersion of sodium hydride in oil (5.2 mg, 0.13 mmol) was then added and the ice bath was removed. After stirring the reaction mixture for 1 hour at rt, the mesylate from step 1 (34 mg, 0.12 mmol) was added. The reaction mixture was heated at 50°C for 24 hours, then cooled to 0°C. An additional 9.6 mg NaH (60% dispersion in oil) was added and heating was resumed for two hours, after which the reaction mixture was cooled to rt and stirred for 48 hours. At this time, ethyl acetate was added and the organic solution was washed with water and brine (three times), dried over sodium sulfate, filtered, and concentrated *in vacuo*. The resulting residue was purified through silica gel flash chromatography by using 5:1 hexane:ethyl acetate as the eluant to provide product (19 mg, 35%) as a mixture of diastereomers (3:1): ES-MS m/z 454 ((M+H)⁺); HPLC RT (min.) 4.21; ¹H NMR (d₆-DMSO) δ 0.80 (t, 3H), 1.38-1.63 (m, 2H), 1.79-1.90 (m, 1H), 2.02-2.14 (m, 1H), 2.34 (s, 3H), 2.51-2.57 (m, 1H), 2.63-2.84 (m, 2H), 2.91 (t, 2H), 3.19-3.25 (m, 1H), 3.49 (s, 2.3H), 3.58 (s, 0.7H), 4.22 (t, 2H), 7.00 (s, 1H), 7.21 (s, 1H), 7.43-7.51 (m, 3H), 7.85-7.90 (m, 2H).



Step 4. Under the standard hydrolysis conditions, the ester from step 3 was converted to the acid (a mixture of diastereomers 3:2): ES-MS m/z 440 ((M+H)⁺); HPLC RT (min.) 3.69; ¹H NMR (d₆-DMSO) δ 0.83 (t, 3H), 2.34 (s, 3H), 2.92 (t, 2H), 4.21 (t, 2H), 7.00-7.02 (d, 1H), 7.12 (s, 0.24H), 7.21 (s, 0.37H), 7.47-7.48 (m, 3H), 7.87-7.90 (m, 2H).

Example 27


Preparation of ethyl 2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}pentanoate




An oven dried 15 mL round-bottom flask and stir bar, cooled under a stream of Ar(g), was charged with ethyl 2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl} acetate (0.070 g, 0.17 mmol) followed by addition of 0.2 mL THF. The stirred solution was then cooled to -78°C followed by dropwise addition of lithium bis(trimethylsilyl)amide (1.0 M hexane solution, 0.86 mL, 0.86 mmol). Upon complete addition of base, the solution was allowed to stir at -78°C for 1 hour, then iodopropane (0.142 g, 0.86 mmol) was added via syringe. The contents were then slowly warmed to rt and maintained for 1 hour. The contents of the flask were poured into 5 mL $\text{NH}_4\text{Cl}(\text{aq})$, then extracted with ethyl acetate (3 x 10 mL). The organic layers were combined and dried over Na_2SO_4 and concentrated *in vacuo* yielding 3.0 mg (4.0% yield) of a colorless film. The product had: ^1H NMR (300 MHz, d_6 -acetone) δ 7.96 (dd, 8.1, 1.5 Hz, 2H), 7.48 (m, 3H), 6.99 (d, 8.4 Hz, 1H), 6.79 (d, 2.7 Hz, 1H), 6.70 (dd, 8.1, 2.7 Hz, 1H), 4.22 (t, 6.9 Hz, 2H), 4.11 (q, 7.2 Hz, 2H), 3.33 (q, 6.6 Hz, 1H), 2.94 (t, 6.9 Hz, 2H), 2.78 (m, 3H), 2.54 (m, 1H), 2.39 (s, 3H), 2.14 (m, 2 H), 1.91 (m, 1H), 1.63 (qt, 10.2, 3.9Hz, 2H), 1.21 (t, 7.2 Hz, 3 H), 0.852 (t, 7.5 Hz, 3 H); mass spectroscopy gave MH^+ of 448.2 (calc'd molecular weight for $\text{C}_{28}\text{H}_{33}\text{NO}_4 = 447.57$).

Example 28

Preparation of 2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}pentanoic acid




Hydrolysis of the product of Example 27 by the method described above for Example 2 gave a product with the following ^1H NMR (300 MHz, d_6 -acetone); δ 7.96 (dd, 8.1, 1.5 Hz, 2H), 7.48 (m, 3H), 7.10 (d, 8.4 Hz, 1H), 6.79 (d, 2.7 Hz, 1H), 6.71 (dd, 8.1, 2.7 Hz, 1H), 4.22 (t, 6.9 Hz, 2H), 3.40 (m, 1H), 2.91 (t, 6.9 Hz, 2H), 2.74 (m, 1H), 2.58 (m, 1H), 2.39 (s, 3H), 2.26 (m, 1H), 2.11 (m, 1H), 1.95 (m, 2H), 1.84 (m, 1H), 1.62 (m, 2H), 0.859 (td, 6.9, 1.5 Hz, 3H); mass spectroscopy gave MH^+ of 420.1 (calc'd molecular weight for $\text{C}_{26}\text{H}_{29}\text{NO}_4 = 419.51$).

Example 29**Preparation of 2-{6-bromo-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid****via****methyl 2-(6-bromo-5-hydroxy-2,3-dihydro-1H-inden-1-yl)butanoate**


Step 1. A solution of bromine (0.032 mL, 0.60 mmol) in dioxane (3 mL) was cooled to 0°C for 15 minutes after which a solution of 2-(5-hydroxy-indan-1-yl)-butyric acid methyl ester (141 mg, 0.60 mmol) in dioxane (3 mL) was added. After 5 minutes, the ice bath was removed and the reaction was stirred at rt for 4 hours. Solvent was removed by rotary evaporation. The residue was purified by column chromatography (8% EtOAc in hexane) to obtain a colorless oil of mono-bromo intermediate (**A**) (145 mg, 77%) and dibromo intermediate (**B**) (20 mg).

A: R_f = 0.46 (4 : 1 hexane : EtOAc); GC-MS (+Cl): m/z = 313 (M^+); ^1H NMR (DMSO - d_6): δ 0.840 (m, 3H), 1.511 (m, 2H), 1.905 (m, 1H), 2.091 (m, 1H), 2.410 – 2.793 (m, 3H), 3.212 (m, 1H), 3.505 and 3.512 (s, 3H), 6.713 and 6.753 (s, 1H), 7.034 and 7.274 (s, 1H), 9.932 and 9.934 (s, OH).

B: R_f = 0.30 (4 : 1 hexane : EtOAc); GC-MS(+Cl): m/z = 393 (M^+); ^1H NMR (DMSO - d_6): δ 0.817 (m, 3H), 1.459 – 1.596 (m, 2H), 1.910 (m, 1H), 2.101 (m, 1H), 2.433 – 2.768 (m, 3H), 3.371 (m, 1H), 3.400 and 3.596 (s, 3H), 7.168 and 7.357 (s, 1H), 9.535 and 9.542 (s, OH).




Step 2. To a solution of (A) from step 1 above (118 mg, 0.38 mmol) in DMF (3.8 mL) at 0°C, was added NaH (60% in mineral oil, 30 mg). After 1 hour, the mesylate as prepared in step 1, Example 26 was added. The mixture was heated to 50°C for 30 hours. The solution was diluted with water, and then extracted with ethyl acetate three times. The combined organic layer was washed with water and brine, then dried (Na_2SO_4) and concentrated. The residue was purified by column chromatography (10% ethyl acetate in hexane) to give product (63 mg, 34%); $R_f = 0.46$ (2 : 1 hexane : EtOAc); ES/MS: $m/z = 498$ (MH^+); ^1H NMR ($\text{DMSO} - d_6$): δ 0.847 (m, 3H), 1.468 (m, 2H), 1.812 (m, 1H), 2.146 (m, 1H), 2.340 (s, 3H), 2.525 – 2.788 (m, 3H), 2.902 (m, 2H), 3.236 (m, 1H), 3.481 and 3.586 (s, 3H), 4.211 (m, 2H), 6.969 (s, 1H), 7.347 and 7.386 (s, 1H), 7.452 (m, 3H), 7.833 (m, 2H).




Step 3. To a solution of product from step 2 (5.6 mg) in methanol, was added 3 N KOH (1 mL) followed by addition of THF until the cloudy solution became clear. The mixture was refluxed overnight. Conc. HCl was added to adjust the pH to 2, then extracted three times with ethyl acetate. The organic layers were combined, dried, and concentrated to give white solid (4 mg). $R_f = 0.18$ (2:1 hexane:EtOAc); ES/MS: $m/z = 484$ (MH^+); ^1H NMR ($\text{DMSO} - d_6$): δ 0.832 (m, 3H), 1.468 (m, 2H), 1.812 (m, 1H), 2.146 (m, 1H), 2.405 (m, 1H), 2.788 (m, 2H), 2.904 (m, 2H), 3.015 (m, 1H), 3.136 and 3.138 (s, 3H), 4.209 (m, 2H), 6.987 and 7.344 (s, 1H), 6.972 and 7.251 (s, 1H), 7.487 (m, 3H), 7.882 (m, 2H).


Example 30

Preparation of 2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-6-phenyl-2,3-dihydro-1H-inden-1-yl}butanoic acid
via
methyl 2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-6-phenyl-2,3-dihydro-1H-inden-1-yl}butanoate






Step 1. A mixture of the product of step 2, Example 29 and $\text{Pd(PPh}_3)_4$ in THF (1.5 mL) was stirred at rt for 30 minutes. Phenylboronic acid (13.2 mg, 0.108 mmol) and 2 N NaOH were then added into the solution. The mixture was heated to reflux for 14 hours. The solution was allowed to cool down, diluted with water, and extracted with ethyl acetate three times. The combined organic layers were washed with brine and dried over sodium sulfate. The crude product was purified by column chromatography eluting with 5% ethyl acetate in hexane to obtain the desired product (8.6 mg). $R_f = 0.48$ (2:1 hexane:EtOAc); ESI-MS: $m/z = 496$ (MH^+); $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 0.804 (m, 3H), 1.541 (m, 2H), 1.880 (m, 1H), 1.987 (m, 1H), 2.090 (s, 3H), 2.247 – 2.698 (m, 3H), 2.791 (m, 2H), 3.199 (m, 1H), 3.524 and 3.537 (s, 3H), 4.190 (m, 2H), 6.970 (s, 1H), 7.062 (s, 1H), 7.275 (m, 5H), 7.472 (m, 3H), 7.868 (m, 2H).



Step 2. The ester was hydrolyzed by methods described above to give product: $R_f = 0.16$ (2 : 1 hexane : EtOAc); ESI-MS: $m/z = 482$ (MH^+); $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 0.923 (m, 3H), 1.504 (m, 2H), 1.812 (m, 1H), 2.146 (m, 1H), 2.188 (s, 3H), 2.334 (m, 2H), 2.432 (m, 2H), 2.539 (m, 1H), 2.625 (m, 1H), 4.287 (m, 2H), 7.059 (s, 1H), 7.160 (s, 1H), 7.351 (m, 5H), 7.544 (m, 3H), 7.971 (m, 2H).

Example 31

Preparation of methyl 2-{6-[4-(2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoate




A mixture of the product prepared in step 2, Example 29 (71.4 mg, 0.14 mmol), NaHCO_3 (14.3 mg, 0.17 mmol), 4-chlorophenylboronic acid (26.8 mg, 0.17 mmol) in ethylene glycol dimethyl ether (1.5 mL) and water (0.4 mL) was degassed for 20 minutes.

Pd(dppf)Cl₂ was then added to the solution. The mixture was heated to reflux for 2 days. The mixture was then concentrated and purified with column chromatography (10% EtOAc in hexane) to obtain desired product (25 mg). $R_f = 0.51$ (2:1 hexane:EtOAc); ES/MS: $m/z = 530$ (MH⁺); ¹H NMR (DMSO - *d*₆): δ 0.841 (m, 3H), 1.557 (m, 2H), 1.888 (m, 1H), 1.987 (m, 1H), 2.146 (s, 3H), 2.247 – 2.698 (m, 3H), 2.791 (m, 2H), 3.214 (m, 1H), 3.487 and 3.5538 (s, 3H), 4.189 (m, 2H), 6.993 (s, 1H), 7.080 (s, 1H), 7.308 (s, 4H), 7.493 (m, 3H), 7.868 (m, 2H).

By using the above described methods for Examples 26-31 and substituting the appropriate starting materials, the following were made and characterized:

Example 32


2-{6-chloro-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid




ES/MS: $m/z = 516$ (MH⁺); ¹H NMR (DMSO - *d*₆): δ 0.847 (m, 3H), 1.557 (m, 2H), 1.888 (m, 1H), 1.987 (m, 1H), 2.137 (s, 3H), 2.247 – 2.687 (m, 3H), 2.819 (m, 2H), 3.234 (m, 1H), 4.187 (m, 2H), 6.994 (s, 1H), 7.089 (s, 1H), 7.298 and 7.308 (m, 4H), 7.484 (m, 3H), 7.869 (m, 2H).

Example 33


Methyl 2-{6-methyl-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoate




$R_f = 0.23$ (2:1 hexane:EtOAc); ES/MS: $m/z = 434$ (MH⁺); ¹H NMR (DMSO - *d*₆): δ 0.804 (m, 3H), 1.522 (m, 2H), 1.830 (m, 1H), 1.987 (m, 1H), 2.037 (s, 3H), 2.335 (s, 3H), 2.410 – 2.550 (m, 3H), 2.901 (m, 2H), 3.146 (m, 1H), 3.507 (s, 3H), 4.163 (m, 2H), 6.777 (s, 1H), 6.939 (s, 1H), 7.483 (m, 3H), 7.875 (m, 2H).

Example 34**2-[6-methyl-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl]butanoic acid**


$R_f = 0.31$ (2:1 hexane:EtOAc); ESLC-MS: $m/z = 420$ (MH^+); 1H NMR (DMSO - d_6): δ 0.827 (m, 3H), 1.508 (m, 2H), 1.828 (m, 1H), 1.987 (m, 1H), 2.017 (s, 3H), 2.333 (s, 3H), 2.410 – 2.550 (m, 3H), 2.894 (m, 2H), 3.146 (m, 1H), 4.116 (m, 2H), 6.773 (s, 1H), 6.942 (s, 1H), 7.467 (m, 3H), 7.880 (m, 2H).

Example 35**Methyl 2-[5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-6-(2-thienyl)-2,3-dihydro-1H-inden-1-yl]butanoate**


$R_f = 0.60$ (2:1 hexane:EtOAc); ESLC-MS: $m/z = 502$ (MH^+); 1H NMR (DMSO - d_6): δ 0.801 (m, 3H), 1.535 (m, 2H), 1.891 (m, 1H), 1.987 (m, 1H), 2.299 (s, 3H), 2.410 – 2.550 (m, 3H), 2.988 (m, 2H), 3.146 (m, 1H), 3.506 (s, 3H), 4.337 (m, 2H), 7.011 – 7.041 (m, 2H), 7.405 – 7.493 (m, 5H), 7.884 (m, 2H).

Example 36**2-[5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-6-(2-thienyl)-2,3-dihydro-1H-inden-1-yl]butanoic acid**


$R_f = 0.18$ (2:1 hexane:EtOAc); ESLC-MS: $m/z = 488$ (MH^+); 1H NMR (DMSO - d_6): δ 0.801 (m, 3H), 1.535 (m, 2H), 1.891 (m, 1H), 1.987 (m, 1H), 2.299 (s, 3H), 2.410 – 2.550 (m, 3H), 2.988 (m, 2H), 3.146 (m, 1H), 4.337 (m, 2H), 7.078 (m, 2H), 7.472 (m, 5H), 7.896 (m, 2H).


Example 37**Methyl 2-{4,6-dibromo-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoate**

$R_f = 0.35$ (4:1 hexane:EtOAc); ES/MS: $m/z = 578$ (MH^+); 1H NMR (DMSO - d_6): δ 0.847 (m, 3H), 1.468 (m, 2H), 1.812 (m, 1H), 2.146 (m, 1H), 2.350 (s, 3H), 2.407 – 2.788 (m, 3H), 2.982 (m, 2H), 3.225 (m, 1H), 3.480 and 3.588 (s, 3H), 4.145 (m, 2H), 7.276 (s, 1H), 7.458 (m, 3H), 7.866 (m, 2H).


Example 38**2-{4,6-dibromo-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid**

$R_f = 0.17$ (2:1 hexane:EtOAc); ES/MS: $m/z = 564$ (MH^+); 1H NMR (DMSO - d_6): δ 0.847 (m, 3H), 1.468 (m, 2H), 1.812 (m, 1H), 2.146 (m, 1H), 2.361 (s, 3H), 2.414 – 2.781 (m, 3H), 2.995 (m, 2H), 3.123 (m, 1H), 4.125 (m, 2H), 7.345 (s, 1H), 7.437 (m, 3H), 7.886 (m, 2H).


Example 39**Preparation of 2-{6-acetyl-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid****via****methyl 2-(6-acetyl-5-methoxy-2,3-dihydro-1H-inden-1-yl)butanoate**




Step 1. To a solution of AlCl_3 (103 mg, 0.78 mmol) in methylene chloride (2.5 mL) at 0°C , was added acetyl chloride (0.044 mL, 0.63 mmol), followed by the addition of a solution of methyl 5-methoxy-2,3-dihydro-1H-indene-1-yl-butanoate (130 mg, 0.52 mmol) in methylene chloride (2.7 mL) dropwise. The mixture was stirred at 0°C for 15 minutes. The ice bath was removed and the mixture stirred at rt for 16 hours. The mixture was poured over ice and 4 drops of conc. HCl were added. This mixture was extracted with methylene chloride twice. The combined organic layers were washed with water, 0.05 N NaOH and water. The organic layer was dried, concentrated, and purified by chromatography with 10% EtOAc:hexane to give desired product (103 mg, 68%). $R_f = 0.28$ (4:1 hexane:EtOAc); GC-MS (+Cl): $m/z = 291$ (M^+); ^1H NMR ($\text{DMSO} - d_6$): δ 0.840 (m, 3H), 1.536 (m, 2H), 1.876 (m, 1H), 2.108 (m, 1H), 2.505 (s, 3H), 2.521 (m, 1H), 2.760 – 2.889 (m, 2H), 3.236 (m, 1H), 3.511 and 3.589 (s, 3H), 3.836 (s, 3H), 7.012 and 7.253 (s, 1H), 7.440 (s, 1H).



Step 2. To a solution of AlCl_3 (238 mg, 1.77 mmol) in CH_2Cl_2 (1 mL), was added the product of step 1 (103 mg, 0.35 mmol) in CH_2Cl_2 (2 mL). The mixture was cooled to 0°C for 5 minutes, then EtSH (0.13 mL, 1.77 mmol) was added slowly. The mixture was stirred at this temperature for 4.5 hours. The mixture was then poured over ice water, stirred for 10 minutes, and extracted with CH_2Cl_2 twice. The combined organic layers were washed with water, dried over sodium sulfate, and concentrated to give product (86 mg, 89%). $R_f = 0.51$ (4:1 hexane:EtOAc); GC-MS (+Cl): $m/z = 276$ (M^+); ^1H NMR ($\text{DMSO} - d_6$): δ 0.841 (m, 3H), 1.574 (m, 2H), 1.888 (m, 1H), 2.094 (m, 1H), 2.585 (s, 3H), 2.639 (m, 1H), 2.729 – 2.847 (m, 2H), 3.244 (m, 1H), 3.513 and 3.628 (s, 3H), 6.774 and 7.503 (s, 1H), 6.792 and 7.715 (s, 1H), 12.117 and 12.143 (s, 1H).



Step 3. The coupling of the hydroxy indene acetic acid ester of step 2 with the mesylate of step 2, Example 26. ESLC-MS: $m/z = 462$ (MH^+);




Step 4. The hydrolysis of the product from step 3 was carried out in similar fashion as described above to give product: $R_f = 0.08$ (2:1 hexane:EtOAc); ESLC-MS: $m/z = 448$ (MH^+); 1H NMR ($DMSO - d_6$): δ 0.848 (m, 3H), 1.468 (m, 2H), 1.812 (m, 1H), 2.146 (m, 1H), 2.305 (s, 3H), 2.368 (s, 3H), 2.405 (m, 1H), 2.788 (m, 2H), 2.971 (m, 2H), 3.015 (m, 1H), 4.332 (m, 2H), 7.039 and 7.441 (s, 1H), 7.446 (s, 1H), 7.465 (m, 3H), 7.875 (m, 2H).


Using a combination of the above described procedures and substituting the appropriate starting materials, a variety of compounds were prepared and are described below.

Example 40


Methyl 2-{5-[2-(2,5-diphenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoate




Yield: 0.09 g, 46%; 1H NMR ($CDCl_3$, 400 MHz) δ 0.83-0.93 (t, 3 H), 1.55-1.78 (m, 2 H), 1.87-1.97 (m, 1 H), 2.10-2.22 (m, 1 H), 2.44-2.52 (m, 1 H), 2.67-2.80 (m, 1 H), 2.81-2.93 (m, 1 H), 3.21-3.29 (m, 1 H), 3.23-3.33 (t, 2 H), 3.62 (s, 3 H), 4.34-4.43 (t, 2 H), 6.66-6.72 (m, 1 H), 6.76 (s, 1 H), 7.05-7.14 (d, 1 H), 7.33-7.39 (t, 1 H), 7.43-7.51 (m, 5 H), 7.78-7.84 (d, 2 H), 8.06-8.12 (m, 2 H).

Example 41**2-{5-[2-(2,5-diphenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid**

Yield: 0.07 g, 70%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.85-0.98 (m, 3 H), 1.23-1.47 (m, 1 H), 1.57-1.78 (m, 1 H), 1.88-2.07 (m, 1 H), 2.12-2.27 (m, 1 H), 2.43-2.56 (m, 1 H), 2.68-2.97 (m, 2 H), 3.27-3.35 (t, 2 H), 3.42-3.50 (m, 1 H), 4.34-4.41 (t, 2 H), 6.66-6.73 (d, 1 H), 6.77 (s, 1 H), 7.02-7.16 (d, 1 H), 7.34-7.40 (t, 1 H), 7.43-7.52 (m, 5 H), 7.78-7.83 (d, 2 H), 8.05-8.12 (m, 2 H).

Example 42**Methyl 2-{5-[2-(5-isopropyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoate**

Yield: 0.09 g, 45%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.78-0.96 (t, 3 H), 1.26-1.32 (d, 6 H), 1.51-1.62 (m, 1 H), 1.64-1.75 (m, 1 H), 1.81-1.93 (m, 1 H), 2.07-2.21 (m, 1 H), 2.40-2.51 (m, 1 H), 2.65-2.75 (m, 1 H), 2.77-2.98 (m, 1 H), 2.91-2.98 (t, 2 H), 3.09-3.16 (m, 1 H), 3.21-3.28 (m, 1 H), 3.62 (s, 3 H), 4.10-4.17 (t, 2 H), 6.60-6.68 (d, 1 H), 6.72 (s, 1 H), 7.01-7.13 (d, 1 H), 7.33-7.45 (m, 3 H), 7.94-8.00 (d, 2 H).


Example 43**2-{5-[2-(5-isopropyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid**

Yield: 0.08 g, 97%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.91-0.98 (t, 3 H), 1.30-1.36 (d, 6 H), 1.58-1.79 (m, 2 H), 1.89-2.05 (m, 1 H), 2.12-2.27 (m, 1 H), 2.44-2.57 (m, 1 H), 2.69-2.80 (m, 1 H), 2.83-2.96 (m, 1 H), 2.97-3.02 (t, 2 H), 3.10-3.21 (m, 1 H), 3.24-3.32 (m, 1

H), 4.14-4.21 (t, 2 H), 6.63-6.71 (d, 1 H), 6.75 (s, 1 H), 7.04-7.16 (d, 1 H), 7.36-7.45 (m, 3 H), 7.94-8.00 (d, 2 H).

Example 44


Methyl 2-{5-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-indenyl}butanoate



Yield: 0.14 g, 60%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.85-0.91 (t, 3 H), 1.25-1.35 (t, 3 H), 1.58-1.77 (m, 2 H), 1.85-1.97 (m, 1 H), 2.10-2.22 (m, 1 H), 2.44-2.64 (m, 2 H), 2.68-2.80 (q, 2 H), 2.82-2.93 (m, 1 H), 2.95-3.01 (t, 2 H), 3.25-3.34 (m, 1 H), 3.62 (s, 3 H), 4.16-4.25 (t, 2 H), 6.66-6.71 (d, 1 H), 6.75 (s, 1 H), 7.08-7.14 (d, 1 H), 7.38-7.46 (m, 3 H), 7.95-8.01 (m, 2 H).

Example 45


2-{5-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid



Yield: 0.05 g, 60%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.85-0.98 (m, 3 H), 1.21-1.33 (m, 3 H), 1.37-1.54 (m, 1 H), 1.56-1.78 (m, 2 H), 1.87-2.29 (m, 2 H), 2.45-2.60 (m, 1 H), 2.69-2.79 (q, 2 H), 2.85-2.95 (m, 1 H), 2.96-3.01 (t, 2 H), 3.27-3.49 (m, 1 H), 4.14-4.23 (t, 2 H), 6.65-6.71 (d, 1 H), 6.75 (s, 1 H), 7.03-7.17 (d, 1 H), 7.38-7.46 (m, 3 H), 7.95-8.01 (d, 2 H).

Example 46

Methyl 2-{5-[2-(2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-indenyl}butanoate




Yield: 0.18 g, 80%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.82-0.92 (t, 3 H), 1.56-1.66 (m, 1 H), 1.67-1.77 (m, 1 H), 1.88-1.99 (m, 1 H), 2.12-2.23 (m, 1 H), 2.43-2.52 (m, 1 H), 2.68-

2.81 (m, 1 H), 2.84-2.97 (m, 1 H), 3.02-3.11 (t, 2 H), 3.25-3.33 (m, 1 H), 3.63 (s, 3 H), 4.21-4.30 (t, 2 H), 6.69-6.74 (d, 1 H), 6.79 (s, 1 H), 7.11-7.16 (d, 1 H), 7.41-7.47 (m, 3 H), 7.55-7.58 (m, 1 H), 7.99-8.05 (m, 2 H).

Example 47


2-{5-[2-(2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid



Yield: 0.07 g, 46%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.84-1.01 (m, 3 H), 1.36-1.51 (m, 1 H), 1.59-1.81 (m, 1 H), 1.88-2.00 (m, 1 H), 2.11-2.29 (m, 1 H), 2.43-2.64 (m, 1 H), 2.68-2.81 (m, 1 H), 2.82-3.00 (m, 2 H), 3.02-3.11 (t, 2 H), 3.23-3.37 (m, 1 H), 4.17-4.28 (t, 2 H), 6.66-6.74 (d, 1 H), 6.78 (s, 1 H), 7.04-7.19 (m, 1 H), 7.39-7.47 (m, 2 H), 7.55 (s, 1 H), 7.98-8.05 (m, 2 H).

Example 48


Methyl 2-(5-{2-[2-(2,3-dihydro-1-benzofuran-6-yl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)butanoate



Yield: 0.17 g, 58%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.86-0.97 (t, 3 H), 1.41-1.53 (m, 1 H), 1.61-1.77 (m, 1 H), 1.92-2.01 (m, 1 H), 2.04-2.20 (m, 1 H), 2.40 (s, 3 H), 2.49-2.56 (m, 1 H), 2.71-2.92 (m, 2 H), 3.93-3.00 (t, 2 H), 3.21-3.32 (t, 2 H), 3.34-3.49 (m, 1 H), 3.75 (s, 3 H), 4.18-4.24 (t, 2 H), 4.54-4.70 (t, 2 H), 6.70-6.76 (d, 1 H), 6.79 (s, 1 H), 6.82-6.89 (d, 1 H), 6.92-7.01 (d, 1 H), 7.75-7.80 (d, 1 H), 7.87 (s, 1 H).

Example 49

2-(5-{2-[2-(2,3-dihydro-1-benzofuran-6-yl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)butanoic acid




Yield: 0.10 g, 99%; ^1H NMR (CDCl_3 , 400 MHz) δ 0.90-1.04 (t, 3 H), 1.41-1.54 (m, 1 H), 1.60-1.76 (m, 1 H), 1.83-1.97 (m, 1 H), 2.12-2.23 (m, 1 H), 2.35 (s, 3 H), 2.48-2.60 (m, 1 H), 2.69-2.90 (m, 2 H), 2.92-3.01 (t, 2 H), 3.18-3.28 (t, 2 H), 3.39-3.50 (m, 1 H), 4.08-4.12 (t, 2 H), 4.46-4.64 (t, 2 H), 6.76-6.71 (d, 1 H), 6.73 (s, 1 H), 6.77-6.84 (d, 1 H), 7.01-7.09 (d, 1 H), 7.71-7.78 (d, 1 H), 7.83 (s, 1 H).

Example 50


Preparation of ethoxy{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}acetic acid

via

ethyl [5-(benzyloxy)-2,3-dihydro-1H-inden-1-ylidene](ethoxy)ethanoate




Step 1. LDA (prepared from 11 mmol DIA and 11 mmole BuLi) was added to methyl 2-ethoxyacetate (10 mmol) in 50 mL THF at -78°C , stirred for 1 hour, then TMSCl (30 mmol) was added. The mixture was concentrated *in vacuo*, and was carried to the next step directly without purification.



Step 2. 5-Benzyloxy-1-indanone in CH_2Cl_2 (5 mL) was slowly added to TiCl_4 in CH_2Cl_2 (10 mL) at -78°C , stirred at -60°C for 10 minutes, and cooled to -78°C . The product of step 1 in CH_2Cl_2 (5 mL) was slowly added and stirred for 10 minutes. The reaction was quenched with saturated K_2CO_3 , filtered, extracted with ethyl acetate, and dried over sodium sulfate. Column chromatography yielded a colorless oil as product.

LC-MSMH⁺=353.1, RT = 4.00 min.; NMR (CDCl₃, 400 MHz) δ 7.9 (1H, d), 7.25 (5H, m), 6.78 (2H, m), 4.93 (2H, s), 4.15 (2H, q), 3.75 (2H, q), 3.05 (2H, m), 2.85 (2H, m), 1.22 (6H, m)





Step 3. Using the product of step 2 as starting material and procedures similar to that described for Example 13, steps 4-8, the desired final product was prepared and characterized: LC-MS [MH⁺] = 422.2, RT = 3.25 min.; NMR (CDCl₃, 400 MHz) δ 8.26 (1H, d), 7.55 (2H, m), 7.16 (2H, d), 6.70 (3H, m), 4.16 (2H, q), 3.63 (2H, t), 3.5 (2H, m), 3.30 (1H, m), 3.20 (1H, m), 2.50 (3H, s), 1.10 (3H, m).

Example 51


Preparation of 2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid

via


2-(4-methyl-2-phenyl-1,3-oxazol-5-yl)ethanol




Step 1. To a solution of sodium hydroxide (8.98 g, 224.49 mmol) in water (112.25 mL), was added at rt DL-Alanine (10 g, 112.25 mmol). The resulting solution was heated at 75°C and the benzoyl chloride (15.77 g, 112.25 mmol) was slowly added. The reaction was heated for 30 minutes, and cooled down to 0°C with an ice bath. Conc. HCl was added to adjust the pH to 1, then the white solid was filtrated through a fritted glass funnel and vacuum dried with P₂O₅ overnight. No purification was needed. This gave N-benzoylalanine (19.6 g, 90.4% yield) as white solid. ¹H NMR (DMSO-d₆) δ 12.61 (s br, 1H), 8.64 (d, 1H), 7.87-7.85 (m, 2H), 7.52-7.43 (m, 3H), 4.40 (q, 1H), 1.39 (d, 3H).



Step 2. In the first flask, N-benzoylalanine (2 g, 10.35 mmol) was dissolved in THF (20 mL), and carbonyl diimidazole (CDI) (1.84 g, 11.39 mmol) was added. The resulting mixture was stirred 1 hour at rt and cooled down to -78°C . Into a second flask, ethyl acetate (3.83 g, 43.48 mmol) in THF (40 mL) was cooled down to -78°C and LDA (24.3 mL, 48.51 mmol, 2 M in THF) pre-cooled to -78°C was added. The resulting solution was stirred 30 minutes at -78°C , and the lithium enolate generated was cannulated into the first flask. The resulting white slurry was stirred 30 minutes at -78°C and warmed up to -10°C . The reaction was quenched with a saturated aqueous solution of NH_4Cl . Phases were separated and the organics were dried over MgSO_4 and solvents removed under reduced pressure. The crude product was carried to the next step without purification. This gave ethyl 4-(benzoylamino)-3-oxopentanoate (2.6 g, 95.5% yield) as a white solid. ES-MS m/z 263.4 ($(\text{MH})^+$); HPLC RT (min.) 1.53; ^1H NMR ($\text{Acetone-}d_6$) δ 8.13 (s br, 1H), 7.93-7.91 (m, 2H), 7.58-7.43 (m, 3H), 4.72 (m, 1H), 4.19-4.01 (q, 2H), 3.67 (s, 2H), 1.47 (d, 3H), 1.15 (t, 3H).




Step 3. To a crude mixture of ethyl 4-(benzoylamino)-3-oxopentanoate (0.6 g, 2.28 mmol) in DMF (4 mL) at rt, was added POCl_3 (1.04 g, 6.84 mmol). The resulting solution was heated at 90°C for 1 hour, then cooled down to rt, and poured into ice for 30 minutes. The aqueous solution was carefully added to a saturated aqueous solution of NaHCO_3 . Phases were separated with EtOAc and the combined organic extracts were dried over MgSO_4 and solvent removed under reduced pressure. The crude material was purified on Biotage small column using a solvent gradient of 0 to 50% EtOAc/Hexane. This gave ethyl (4-methyl-2-phenyl-1,3-oxazol-5-yl)acetate (0.269 g 48% yield) as yellowish oil. ES-MS m/z 246.2 ($(\text{MH})^+$); HPLC RT (min.) 2.77; ^1H NMR (CDCl_3) δ 8.01-7.98 (m, 2H), 7.45-7.41 (m, 3H), 4.20 (q, 2H), 3.71 (s, 2H), 2.21 (s, 3H), 1.28 (t, 3H).




Step 4. Ethyl (4-methyl-2-phenyl-1,3-oxazol-5-yl)acetate (0.922 g, 3.76 mmol) in THF (6 mL) at rt, was added LiBH_4 2M/THF (9.41 mL, 4.70 mmol). The reaction was

stirred overnight at rt, then treated with 2 N HCl until pH 7. The solvent THF was removed under reduced pressure, EtOAc was added, and phases separated. The combined organic extracts were dried over MgSO_4 and solvent concentrated *in vacuo*. The crude material was purified by Biotage using a gradient of 10 to 100% EtOAc/Hexane as solvent mixture. This gave 2-(4-methyl-2-phenyl-1,3-oxazol-5-yl)ethanol (0.193 g, 25% yield) as colorless oil. ES-MS m/z 204.2 (MH^+); HPLC RT (min.) 2.02; ^1H NMR (Acetone- d_6) δ 7.98-7.95 (m, 2H), 7.52-7.42 (m, 3H), 3.95 (s br, 1H), 3.82 (t, 2H)m, 2.90 (t, 2H), 2.13 (s, 3H).




Step 5. DEAD (0.84 mL, 5.28 mmol) in THF (1.5 mL) was slowly added to a solution of the product of step 3 (4.95 mmol), methyl 5-hydroxy-2,3-dihydro-inden-1-yl-2-butanoate (0.78 g, 3.3 mmol), PPh_3 (1.4 g, 5.28 mmol) in THF (13 mL). The mixture was stirred at rt overnight. The mixture was filtered, washed with water, brine, dried over sodium sulfate, and concentrated. Column chromatography yielded a colorless oil as product. LC-MS $[\text{C}_{26}\text{H}_{29}\text{NO}_4\text{H}]^+ = 420.4$, RT = 4.00 min.; ^1H NMR (CDCl_3): δ 7.9 (2H,d), 7.45 (2H,dd), 7.1(d), 6.6-6.8 (3H, m), 4.2 (2H, t), 3.62 (3H,s), 3.3 (1H, m), 3.15 (2H,t), 2.6-3.0 (2H, m,br), 2.5 (1H, m), 2.21 (3H,s), 1.95 (1H, m), 1.56-1.6 (3H,br,m), 0.88 (3H,t).



Step 6. KOH (0.5 mL, 3 N) was added to a solution of the product of step 4 (42 mg, 0.1 mmol) in THF/MeOH (1 mL, THF:MeOH 8:2). The mixture was stirred at 70°C for 6 hours, then cooled down. The pH was adjusted to 4 with 1 N HCl. The mixture was extracted with ethyl acetate (3 x 2 mL). The combined organic layers were dried over sodium sulfate and concentrated *in vacuo*. Column chromatography (2:8/hexane:ethyl acetate) gave a white solid as the product (33 mg, 81%). LC-MS $[\text{C}_{25}\text{H}_{27}\text{NO}_4\text{H}]^+ = 406.3$, RT = 3.37 min.; ^1H NMR (CDCl_3): δ 8.0 (2H,d), 7.45 (2H,dd), 7.15 (1H,d), 6.7-6.8 (3H, m), 4.2 (2H, t), 3.3 (1H, m), 3.15 (2H,t), 2.6-3.0 (2H, m,br), 2.5 (1H, m), 2.21 (3H,s), 1.95 (1H, m), 1.56-1.6 (3H,br,m), 0.88 (3H,t).

By using the procedure described above for Example 51 and substituting the appropriate starting materials, the following were similarly prepared and characterized.


Example 52



LC-MS [$C_{26}H_{29}NO_4H$] $^+ = 420.3$, RT = 3.52 min.; 1H NMR ($CDCl_3$): δ 7.87 (2H,d), 7.25 (2H,dd), 7.1(1H,d), 6.6-6.8 (3H, m), 4.2 (2H, t), 3.45 (1H,m), 3.30 (1H, m), 3.15 (2H,t), 2.7-3.0 (2H, m, br), 2.5 (1H, m), 2.4 (3H, s) 1.95 (1H, m), 1.56-1.60 (3H, br ,m), 0.88 (3H,t)

Example 53

2-{5-[2-(4-methyl-2-propyl-1,3-oxazol-5-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}butanoic acid




LC-MS [$C_{22}H_{29}NO_4H$] $^+ = 372.3$, RT = 3.16 min.; 1H NMR ($CDCl_3$): δ 7.1 (1H,d), 6.6 (2H,d), 4.2 (2H, t), 3.3 (1H,m), 3.3 (1H, m), 2.8 (2H,t), 2.7 (1H, m), 2.6 (2H, t), 2.4 (2H,m), 2.2 (3H, s), 2.0-1.8 (2H,br,m), 0.88 (3H,t)




By using the methods described above for Examples 1-53 and by substituting the appropriate starting materials, compounds of Formula Ia, listed in Table 3 below, were similarly prepared.

Table 3


Preparative Examples of Compounds of Formula (Ia)




(Ia)


Ex. No.	R	R ¹	R ²	R ³	R ⁴	R ⁵	X	LC-MS [M+H] ⁺ or NMR
54	H	Et	H	Me	PhOCH ₂ -	H	O	436.2
55	H	Et	H	Me	PhCH ₂ -	H	O	420
56	H	H	H	Me	Ph	H	O	378.2
57	Me	Ph(CH ₂) ₃ -	H	Me	Ph	H	O	3.45/3.52 (t, 3H), 4.10 (t, 2H), 7.3 (m, 3H), 7.83 (d, 2H)
58	Et	EtO ₂ C-	H	Me	Ph	H	O	478.2
59	Et	Et	H	Me	Ph	H	O	434.3
60	H	MeO	H	Me	Ph	H	O	3.30 (s, 3H), 4.04 (d, 1H), 7.98 (m, 2H)
61	Et	EtO	H	Me	Ph	H	O	450.3
62	H	CF ₃ CH ₂ -	H	Me	Ph	H	O	2.51 (s, 3H), 4.36 (m, 2H), 8.32 (m, 2H)
63	Et	CF ₃ CH ₂ -	H	Me	Ph	H	O	1.18 (t, 3H), 4.21 (t, 2H), 7.98 (d, 2H)
64	Me	cyc-Pr	H	Me	Ph	H	O	432.3
65	H	cyc-Pr	H	Me	Ph	H	O	0.02 (m, 1H), 0.12 (m, 1H), 4.18 (m, 2H), 7.94 (m, 2H)
66	H		H	Me	Ph	H	O	512.3
67	H	Et	H	Me	Ph	H	S	422.3
68	H		H	Me	Ph	H	O	526.4
69	H	Et	H	Me	Ph	H	S	422.3
70		Et	H	Me	Ph	H	S	
71	Me	Et	H	Me	Ph	H	O	0.82 (t, 3H), 3.54 (s, 3H), 4.16 (t, 2H), 7.90 (m, 2H)
72	H	Et	H	<i>i</i> -Pr	Ph	H	O	434.3
73	H	Et	H	Ph	Ph	H	O	468.3
74	H	Me	H	Me	Ph	H	S	422.3
75	Me	Me	H	Me	Ph	H	S	
76	Me	Et	MeC(O)-	Me	Ph	H	O	462.4

Ex. No.	R	R ¹	R ²	R ³	R ⁴	R ⁵	X	LC-MS [M+H] ⁺ or NMR
77	Me	Et	4-MeO-Ph	Me	Ph	H	O	526.4
78	H	Et	4-MeO-Ph	Me	Ph	H	O	512.3
79	Me	Et	4-pyridyl	Me	Ph	H	O	497.3
80	H	Et	H	Me	cyc-Pentyl	H	O	398
81	H	Et	H	Me	cyc-Hexyl	H	O	412
82	H	Et	H	Me	4-Ph-Ph-	H	O	482
83	Et	EtO ₂ C-	H	Me	4-Me-Ph-	H	O	492.3
84	H	PhCH ₂ -	H	Me	4-Me-Ph-	H	O	482.4
85	Et	<i>n</i> -Bu	H	Me	4-Me-Ph-	H	O	476.3
86	Et	Me	H	Me	4-Me-Ph-	H	O	434.3
87	Et	PhCH ₂ -	H	Me	4-Me-Ph-	H	O	510.4
88	H	Et	H	Me	4-MeO-Ph	H	O	436.1
89	H	Et	H	Me	4- <i>i</i> -Pr-Ph	H	O	448.2
90	H	Et	H	Me	4-F-PhCH ₂ -	H	O	438.3
91	H	Et	H	Me	4-F-Ph	H	O	424.3
92	H	Et	H	Me	4-Et-Ph	H	O	434.3
93	H	Et	H	Me	4-Cl-PhOCH ₂ -	H	O	470.2
94	H	Et	H	Me	4-Cl-Ph	H	O	440
95	Me	Et	H	Me	4-Cl-Ph	H	S	470.3
96	Me	Et	H	Me	4-Cl-Ph	H	S	470.3
97	H	Et	H	Me	4-CF ₃ -Ph	H	S	490.3
98	Me	Et	H	Me	4-CF ₃ -Ph	H	S	504.3
99	H	Et	H	Me	4-CF ₃ -Ph	H	O	474.3
100	H	Et	H	Me	4-(<i>n</i> -Bu)-Ph	H	O	462.3
101	H	Et	H	Me	4-(<i>t</i> -Bu)-Ph	H	O	462.3
102	H	Et	H	Me	3-Me-Ph	H	O	420.4
103	H	Et	H	Me	3-MeO-Ph	H	O	436.3
104	H	Et	H	Me	3-Me-5-isoxazoly	H	O	411.3


Ex. No.	R	R ¹	R ²	R ³	R ⁴	R ⁵	X	LC-MS [M+H] ⁺ or NMR
105	H	Et	H	Me	3-F-Ph	H	O	424.2
106	H	Et	H	Me	3-F-4-Me-Ph	H	O	438.2
107	H	Et	H	Me	4-F-3-Me-Ph	H	O	438.3
108	Me	Et	H	Me	3-Cl-Ph	H	S	470.3
109	H	Et	H	Me	3-Cl-Ph	H	O	440.3
110	H	Et	H	Me	3-Cl-Ph	H	S	456.3
111	H	Et	H	Me	3-CF ₃ -Ph	H	O	474.2
112	H	Et	H	Me	3,5-(CF ₃) ₂ -Ph	H	O	542.1
113	H	Et	H	Me	3,4-Me ₂ -Ph	H	O	434.3
114	H	Et	H	Me	3,4-Cl ₂ -Ph	H	O	474.2
115	H	Et	H	Me	2,3-Cl ₂ -Ph	H	O	474.1
116	H	Et	H	Me	3,4-(MeO) ₂ -Ph	H	O	466.3
117	H	Et	H	Me	3,4-methylenedioxy-Ph	H	O	466.3
118	H	Et	H	Me	2-thienyl	H	O	412
119	H	Et	H	Me	2-naphthyl	H	O	456.3
120	H	Et	H	Me	2-Me-Ph	H	O	420.3
121	H	Et	H	Me	2-furyl	H	O	396
122	H	Et	H	Me	2-F-Ph	H	O	424.1
123	H	Et	H	Me	2-benzothieryl	H	O	462.2
124	H	Et	H	Me	2,6-F ₂ -Ph	H	O	442.2
125	H	Et	H	Me	3,4-F ₂ -Ph	H	O	442.2
126	H	Et	H	Me	2,4-Cl ₂ -Ph	H	O	473
127	H	Et	H	Me	1-naphthyl	H	O	456.3
128	Me	Et	H	Me		H	O	0.90 (t, 3H), 3.45 (bs, 4H), 3.74 (s, 3H)

Example 129**Preparation of ethyl (5-methoxy-2,3-dihydro-1H-inden-1-ylidene)ethanoate**


To a solution of 5-methoxyindanone (150 g, 0.91 mol) in anhydrous tetrahydrofuran (4.5 L), was added zinc (30 mesh, 103.64 g, 1.59 mol) and copper(I) chloride (4.53 g, 0.045 mol). The suspension was stirred under Ar atmosphere and refluxed for 15 minutes; approximately a 25% portion of ethyl bromoacetate (133 mL, 1.18 mol) was added to the refluxing mixture in a slow dropwise fashion. After allowing to cool and stirring overnight at rt, TLC showed the presence of desired product, indicating the formation of reactive zinc species. The remainder of ethyl bromoacetate was added dropwise; an exotherm was observed (internal temperature increased to 35°C). After 4 hours, TLC showed complete reaction. After the solids settled to the bottom of the flask, the liquid was siphoned off leaving a small amount behind to cover the solids. The flask was re-charged with 5-methoxyindanone (157.6 g, 1.86 mol total), anhydrous tetrahydrofuran (4.5 L), and zinc (80.92 g, 2.73 mol total). Ethyl bromoacetate (140 mL, 2.36 mol total) was added dropwise. An exotherm was observed (internal temperature increased to 35°C). When the stirred mixture cooled to rt, TLC showed the reaction to be complete. The solids were allowed to settle and the liquid was siphoned off. The combined reaction solutions were concentrated *in vacuo* to a volume of ~ 2L. The liquid was then poured into sufficient 1N aqueous hydrochloric acid (cooled in ice water) to bring the pH to 1. The product was extracted with ethyl acetate (2 x 1 L, 1 x 500 mL). The combined extracts were washed with water, brine (1 L each), dried over sodium sulfate, filtered, and concentrated *in vacuo* to afford a dark red oil which solidified gradually (438.3 g; theoretical yield = 432 g). ¹H NMR (CDCl₃): δ 7.5 (d, 1H), 6.8 (m, 2H), 6.2 (t, 1 H), 4.2 (q, 2H), 3.8 (s, 3H), 3.3 (m, 2H), 3.0 (t, 2H), 1.3 (t, 3H). MS (CI) *m/z* 233 [M+H]⁺.

Example 130**Preparation of ethyl (5-methoxy-2,3-dihydro-1H-inden-1-yl)acetate**


The crude product of Example 129 was dissolved in absolute ethanol (2.6 L) and hydrogenated at 40 psi of hydrogen over 10% palladium on carbon (21.6 g). Filtration through Celite and concentration of the filtrate afforded 433.3 g of brown oil (99% yield for 2 steps). ^1H NMR (CDCl_3): δ 7.1 (dd, 1H), 6.8 (d, 1H), 6.7 (dd, 1H), 4.2 (q, 2H), 3.8 (s, 3H), 3.5 (m, 1H), 2.9 (m, 2H), 2.7 (dd, 1H), 2.4 (m, 2H), 1.7 (m, 1H), 1.3 (t, 3H). MS (CI) m/z 235 $[\text{M}+\text{H}]^+$.

Example 131**Preparation of (5-methoxy-2,3-dihydro-1H-inden-1-yl)acetic acid**

To a solution of the crude ester (416 g, 1.77 mol) prepared in Example 130 in 1 L EtOH, was added a solution of NaOH (142 g, 3.54 mol) in 1.5 L water. The cloudy reaction mixture was heated to reflux, during which time the color changed to dark red, and the reaction became homogeneous. After 1 hour, the reaction was cooled to rt, and the EtOH was removed under reduced pressure. The basic aqueous layer was washed with Et_2O (3 x 500 mL), then acidified with conc. HCl to pH ~4 upon which an oil residue formed. The mixture was extracted with Et_2O (4 x 500 mL). The combined extracts were washed with water (2 x 300 mL), brine, then dried over Na_2SO_4 . Filtration and evaporation of solvent under reduced pressure gave the title compound (305 g, 83%) as a yellow solid after overnight drying under vacuum. ^1H NMR (CDCl_3) δ 7.34(d, 1H), 6.71(s, 1H), 6.65(dd, 1H), 3.71(s, 3H), 3.47(m, 1H), 2.80(m, 3H), 2.35(m, 2H), 1.71(m, 1H). MS (CI) m/z 207 $[\text{M}+\text{H}]^+$.

Example 132**Preparation of [(1S)-5-methoxy-2,3-dihydro-1H-inden-1-yl]acetic acid**

To a solution of the acid (341.0 g, 1.65 mol) prepared in Example 131 in 8.2 L reagent grade acetone, was added (S)-(-)- α -methylbenzylamine (223.8 mL, 1.74 mol) dropwise at rt with stirring. A thick white precipitate formed during the addition. An additional 500 mL acetone was added and stirring continued for 1 hour. The solids were collected by filtration, washed with 300 mL acetone, and dried under suction. The solids were then suspended in acetone (8.2 L) and warmed to reflux until all solids dissolved. The solution was cooled slowly overnight, during which time a white precipitate formed. The suspension was cooled to 0°C, then filtered, and the solids were washed with 500 mL acetone. After drying under suction, a sample analyzed by HPLC showed 95% ee. The recrystallization process was repeated as above using 6.7 L acetone. HPLC analysis showed 99% ee. After drying under suction, 192 g salt were obtained. The salt was suspended in 2 L EtOAc and 1 L of 1 N HCl solution, and shaken in a separatory funnel, whereupon the salt dissolved. The organic layer was separated, washed with 1 N HCl (500 mL), water (2 x 300 mL), and brine, then dried over Na₂SO₄. The solvent was evaporated under reduced pressure, giving an oil which soon solidified. The title product (120.5 g, 35%) was obtained as an off-white solid after vacuum drying. ¹H NMR (CDCl₃) δ 7.10(d, 1H), 6.79(d, 1H), 6.73(dd, 1H), 3.79(s, 3H), 3.55(m, 1H), 2.89(m, 2H), 2.79(dd, 1H), 2.46(dd, 1H), 2.43(m, 1H), 1.80(m, 1H). MS (ESI) m/z 207 [M+H]⁺.


Example 133**Preparation of [(1S)-5-methoxy-2,3-dihydro-1H-inden-1-yl]acetic acid**

As an alternative to Example 132, the title compound may also be prepared via an enzymatic process. Thus, a cloudy mixture of the crude ester (500.0 g, 2.13 mol; 87% pure as determined by HPLC) prepared in Example 130, in 1 L reagent grade acetone, 2.5 L Phosphate Buffer (pH 7.0, 0.05 M) and 2.5 L deionized water was treated in one

portion with Amano Lipase PS (150 g), and the mixture stirred efficiently at rt overnight. HPLC analysis of an aliquot (homogeneous aliquot prepared by dissolving aliquot in IPA followed by filtration) showed one peak corresponding to unreacted R-ester and another peak corresponding to desired S-acid. Trace amounts of S-ester and R-acid were noted. 2 N HCl (500 mL, ensure a pH ~2) was added in one portion to the reaction and stirred for 20 minutes. The mixture was filtered and the solids were washed with EtOAc (2 x 500 mL), then water (500 mL). The combined filtrates were further diluted with 1 L EtOAc, and the layers stirred together vigorously. Stirring was stopped and the layers allowed to separate. Emulsions were noted, but could be broken with the addition of solid NaCl and stirring. The aqueous layer was removed, then extracted with EtOAc (3 x 1 L) in the same fashion. The combined organic extractions were washed with water (4 x 500 mL), then with brine. The resulting organic layer was extracted with a 5% Na₂CO₃ solution (8 x 500 mL). HPLC analysis of the organic layer showed that it contained none of the S-enantiomer acid. The combined Na₂CO₃ extracts were washed with EtOAc (2 x 1 L), then acidified to pH ~2 by the addition of 2N HCl. A white solid precipitated, accompanied by CO₂ evolution. The mixture was extracted with EtOAc (3 x 1 L). The combined extracts were washed with water (2 x 1 L) and brine, then dried over Na₂SO₄. HPLC analysis of this solution showed the material was 98% ee. The solvent was evaporated under reduced pressure, giving an oil which soon solidified. The title product (172.9 g) was obtained as an off-white solid after vacuum drying. This material was recrystallized from boiling hexanes (8.8 L). After overnight cooling, light yellow needles were collected via filtration, washed with hexanes (200 mL), and dried under suction. The title product (146.9 g, 38% from crude starting ester) was obtained as light yellow needles after vacuum drying. ¹H NMR results as above.

Example 134

Preparation of ethyl [(1S)-5-methoxy-2,3-dihydro-1H-inden-1-yl]acetate




To a solution of the acid (305 g, 1.48 mol) prepared in either Example 132 or 133 in 4.8 L absolute EtOH at rt under argon, was added chlorotrimethylsilane (413 mL, 3.25 mol) dropwise. An approximate 5°C rise in temperature was noted during the addition. The reaction was stirred overnight. EtOH was evaporated under reduced pressure,

giving a bi-phasic liquid mixture. This was diluted in 500 mL ice-water, then extracted with EtOAc (2 x 750 mL). The combined extracts were washed with water (3 x 300 mL), then with saturated NaHCO₃ (200 mL). The organic was washed once more with water (300 mL), then brine, and dried over Na₂SO₄. The title compound (354 g, 102%) was obtained as a light yellow oil after solvent removal and vacuum drying. ¹H NMR (CDCl₃) δ 7.07(d, 1H), 6.78(d, 1H), 6.71(dd, 1H), 4.18(q, 2H), 3.78(s, 3H), 3.52(m, 1H), 2.89(m, 2H), 2.72(dd, 1H), 2.37(o, 2H), 1.74(m, 1H), 1.28(t, 3H). MS (CI) *m/z* 235 [M+H]⁺.

Example 135


Preparation of ethyl [(1S)-5-hydroxy-2,3-dihydro-1H-inden-1-yl]acetate



To a cold solution (ice water bath) of the compound (346 g, 1.48 mol) prepared in Example 134 in 4.2 L CH₂Cl₂, was added AlCl₃ (984.6 g, 7.38 mol) portionwise under Ar such that the reaction temperature was maintained below 10°C. The light brown suspension was stirred 10 minutes, then EtSH (546 mL, 7.38 mol) was added dropwise at such a rate that the reaction temperature was maintained below 5°C. After 2.5 hours of stirring below 10°C, the reaction mixture was slowly poured into 6 L ice water with strong agitation. The organic layer was separated, and the aqueous layer was extracted with CH₂Cl₂ (3 x 1 L). The combined CH₂Cl₂ layers were washed with water (2 x 1 L), then dried over Na₂SO₄. The solvent was removed under reduced pressure, giving a brown oil, which was filtered through a pad of silica gel (eluted with 0-10% EtOAc/Hexanes). Fractions were collected and the title compound (314 g, 96%) was obtained as a thick yellow oil after solvent removal and vacuum drying. ¹H NMR (CDCl₃) δ 6.92(d, 1H), 6.62(d, 1H), 6.55(dd, 1H), 4.10(q, 2H), 3.43(q, 1H), 2.75(m, 2H), 2.64(dd, 1H), 2.31(dd, 1H), 2.29(m, 1H), 1.67(m, 1H), 1.20 (t, 3H). MS (CI) *m/z* 221 [M+H]⁺.

Example 136


Preparation of ethyl 2-((1S)-5-[2-[5-methyl-2-(4-methylphenyl)(1,3-oxazol-4-yl)]ethoxy]indanyl)acetate



A suspension of the ethyl [(1*S*)-5-hydroxy-2,3-dihydro-1*H*-inden-1-yl]acetate prepared in Example 135 (507.5 mg, 2.30 mmol), and 2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethanol prepared in Example 10 (500 mg, 2.30 mmol), TMAD (792.6 mg, 4.60 mmol), and Ph₃P (1.21 g, 4.60 mmol) in 15 mL anhydrous DCM was stirred at rt under Ar for 12 hours. DCM was removed under reduced pressure. Flash chromatograph of the residue over silica gel using 1% CH₃CN/CH₂Cl₂ gave ethyl 2-((1*S*)-5-{2-[5-methyl-2-(4-methylphenyl)(1,3-oxazol-4-yl)]ethoxy}indanyl)acetate (776.3 mg, 1.85 mmol, 80.5%). HPLC/MS (M+H)⁺ m/z 420.5.

Example 137


Preparation of 2-((1*S*)-5-{2-[5-methyl-2-(4-methylphenyl)(1,3-oxazol-4-yl)]ethoxy}indanyl)acetic acid



Ethyl 2-((1*S*)-5-{2-[5-methyl-2-(4-methylphenyl)(1,3-oxazol-4-yl)]ethoxy}indanyl)acetate (Example 136, 776.3 mg, 1.85 mmol) in THF (4.0 ml) was added to a mixture of aqueous LiOH (2 M, 3.7 ml, 7.4 mmol), water (2.0 ml), and EtOH (4.0 ml) at rt. The resulting mixture turned cloudy. This mixture was heated at 40°C (oil-bath temperature). The reaction was completed after 1.5 hours. After cooling to rt, 1 N HCl solution was slowly added to the mixture until pH 4.0. The compound was extracted with EtOAc (3 x 20 ml). The combined EtOAc layers were dried (Na₂SO₄) and evaporated. Flash chromatography of the residue gave 2-((1*S*)-5-{2-[5-methyl-2-(4-methylphenyl)(1,3-oxazol-4-yl)]ethoxy}indanyl)acetic acid (616.8 mg, 1.57 mmol, 85%) as a white solid. ¹H NMR (CDCl₃) δ 7.83(d, 2H), 7.21(d, 2H), 7.03(d, 1H), 6.74(d, 1H), 6.69(dd, 1H), 4.19(t, 2H), 3.45(q, 1H), 2.93(t, 2H), 2.78(m, 2H), 2.51(m, 2H), 2.30(s, 3H), 2.25(s, 3H), 1.53(m, 2H).


By using the methods described above for Examples 129-137 and by substituting the appropriate starting materials, compounds of Formula Ia, listed in Table 4 below, were similarly prepared.

Table 4
Preparative Examples of Compounds of Formula (Ia)



Ex. No.	R ³	R ⁴	LC/MS [M+H]
138	Me	4-MeO-Ph	408.5
139	Me	3-MeO-Ph	408.5
140	Me	4-Et-Ph	406.5
141	Me	4-CF ₃ -Ph	446.5
142	Me	2-naphthyl	428.5
143	Me	4-(<i>t</i> -Bu)-Ph	434.6
144	Me	4-(<i>n</i> -Bu)-Ph	434.6
145	Me		422.5
146	Me	3,4-(Me) ₂ -Ph	406.5
147	Me	4-Me-Ph	392.5
148	Me	3-F-Ph	396.5
149	Me	2-benzothienyl	434.5
150	Me	4- <i>i</i> -Pr-Ph	420.6
151	Me	cyc-Pentyl	370.5
152	Me	cyc-hexyl	384.5
153	Me	PhCH ₂	392.5

Ex. No.	R ³	R ⁴	LC/MS [M+H]
154	Me	4-F-3-Me-Ph	410.5
155	Me	3-F-4-Me-Ph	410.5
156	Me	4-F-Ph	396.5
157	Et	Ph	392.5
158	Me	3,4-(Cl) ₂ -Ph	447.4
159	<i>n</i> -Pr	Ph	406.5
160	Me	4-Ph-Ph	454.5
161	Me	3-Cl-Ph	412.4
162	Me	3-Me-Ph	392.5
163	Me	4-CN-Ph	403.4
164	Me	3-CN-Ph	403.4
165	Me	4-Cl-Ph	412.4
166	Me	3-CF ₃ -Ph	446.4
167	Et	4-Et-Ph	420.5
168	Et	4-Me-Ph	406.5
169	Et	4-MeO-Ph	422.4


Example 170**Preparation of methyl 4-bromo-3-oxopentanoate**

A dry three-neck flask under an Ar atmosphere was charged with a solution of methyl propionylacetate (20 g, 154 mmol) in CHCl₃ (100 mL). Using an addition funnel, bromine (7.9 mL, 24.6 g, 154 mmol) was added dropwise over a period of 2 hours at 0°C.

The reaction was then allowed to warm slowly to rt, and the reaction mixture was stirred overnight. A saturated solution of Na_2CO_3 (40 mL) was slowly added, and after stirring the reaction mixture for an additional 15 minutes, the solvents layers were separated and the aqueous layer was extracted with CH_2Cl_2 (50 mL). The combined organic layers were dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was then purified by silica gel flash chromatography (10:1 hexanes/EtOAc) to give the desired bromide as a light yellow oil (25 g, 78%). ^1H NMR (CDCl_3): δ 1.80 (d, 3H), 3.64-3.92 (m, 2H), 3.78 (s, 3H), 4.61 (q, 1H).

Example 171


Preparation of methyl (2-amino-5-methyl-1,3-thiazol-4-yl)acetate



To a solution of bromide of Example 170 (18 g, 86 mmol) in toluene (100 mL) was added thiourea (10.5 g, 138 mmol). The reaction mixture was heated to 100°C for 1 hour, cooled to rt, and the solvent removed under reduced pressure. The residue was dissolved with CH_2Cl_2 (100 mL), a saturated solution NaHCO_3 (75 mL) added, and the mixture was vigorously stirred for 10 minutes. The organic layer was separated, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was then recrystallized from CH_2Cl_2 /hexanes to provide the product (10 g, 63%) as a white solid. ($\text{C}_7\text{H}_{10}\text{N}_2\text{O}_2\text{S}$): LC-MS, RT 0.76 min, M+H 187.0; ^1H NMR (CDCl_3): δ 2.23 (s, 3H), 3.70 (s, 2H), 3.75 (s, 3H), 4.83-4.95 (broad s, 2H).

Example 172

Preparation of methyl (2-bromo-5-methyl-1,3-thiazol-4-yl)acetate




To a solution of CuBr_2 (4.03 g, 18.1 mmol) and *t*-butyl nitrite (2.82 mL, 23.8 mmol) in MeCN (210 mL) was added the compound of Example 170 (2.95 g, 15.9 mmol) at -20°C . The reaction mixture was slowly warmed to 15°C , at which point the evolution of N_2 was observed. After stirring for an additional 2 hours at 15°C , the reaction mixture was diluted with Et_2O (400 mL) and washed with a 10% solution of HCl (200 mL). The solvent layers were separated, the aqueous re-extracted with Et_2O (2 x 300 mL), and the combined organic layers dried (MgSO_4), filtered, and concentrated under reduced pressure. The residue was then purified by silica gel flash chromatography (98:2,

hexanes/EtOAc) to afford bromide Example **172** (1.6 g, 40%) as a colorless oil that solidifies upon standing. (C₇H₈BrNO₂S): LC-MS, RT 2.56 min., M+H 250.3; ¹H NMR (CDCl₃): δ 2.26 (s, 3H), 3.60 (s, 2H), 3.61 (s, 3H).

Example 173


Preparation of 2-(2-bromo-5-methyl-1,3-thiazol-4-yl)ethanol



To a solution of ester prepared in Example **172** (3.80 g, 15.2 mmol) in CH₂Cl₂ (100 mL) was added DIBAL-H (33.4 mL, 33.4 mmol of a 1.0 M solution in toluene) at -78°C. After 15 minutes, the solution was warmed to 0°C and stirred for an additional 90 minutes. An aqueous solution of 2 N HCl (50 mL) was then added dropwise to quench the excess DIBAL-H. The solvent layers were separated and the aqueous layer extracted with CH₂Cl₂ (2 x 200 mL). The combined organic layers were dried (MgSO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel flash chromatography (5:2 hexanes/EtOAc) to yield the product (2.5 g, 74%) as a yellowish oil that solidifies upon standing. (C₆H₈BrNOS) LC-MS, RT 1.38 min., M+H 221.0; ¹H NMR (CDCl₃): δ 2.31 (s, 3H), 2.82 (t, 2H), 2.90-3.00 (broad s, 1H), 3.89 (t, 2H).


Example 174

Preparation of ethyl {(1S)-5-[2-(2-bromo-5-methyl-1,3-thiazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}acetate



Step 1. To a solution of Example **173** (975 mg, 4.39 mmol) and ethyl [(1S)-5-hydroxy-2,3-dihydro-1H-inden-1-yl]acetate (1.06 g, 4.83 mmol) in THF (20 mL) were added Ph₃P (1.88 g, 7.46 mmol) and ADDP (1.96 g, 7.46 mmol). The mixture was vigorously stirred at rt for 72 hours, the solvent removed under reduced pressure, and the residue purified by silica gel flash chromatography (6:1 hexanes/EtOAc) to yield the product (1.4 g, 76%) as a colorless oil that solidifies upon standing. (C₁₉H₂₂BrNO₃S) LC-MS, RT 3.92 min., M+H 424.5; ¹H NMR (CDCl₃): δ 1.26 (t, 3H), 1.65-1.81 (m, 1H), 2.28-2.45 (m, 2H), 2.37 (s, 3H), 2.69 (dd, 1H), 2.75-2.93 (m, 2H), 3.07 (t, 2H), 3.44-3.56 (m, 1H), 4.15 (t, 2H), 4.18 (q, 2H), 6.67 (dd, 1H), 6.73 (d, 1H), 7.03 (d, 1H).


Preparation of ethyl ((1S)-5-{2-[2-(4-isopropylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetate



Step 2. To a mixture of toluene (15 mL) and 1,4-dioxane (3 mL), were added the compound of step 1 (300 mg, 0.708 mmol), 4-isopropylbenzene boronic acid (464 mg, 2.83 mmol), and PdCl₂(dppf).CH₂Cl₂ (52 mg, 0.071 mmol). A flow of Ar was passed through the mixture for 30 minutes, then a 2 N solution of Na₂CO₃ (3.7 mL, 7.08 mmol) was added and the reaction was heated to 75°C for 18 hours. The reaction mixture was then cooled to rt, diluted with EtOAc (200 mL), and washed with a saturated solution of NaHCO₃ (50 mL). The organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel flash chromatography (8:1 hexanes/EtOAc), to provide the product (305 mg, 93%) as a colorless oil. (C₂₈H₃₃NO₃S): LC-MS, RT 5.17 min., M+H 464.5; ¹H NMR (CDCl₃): δ 1.17-1.31 (m, 3H), 1.26 (s, 3H), 1.27 (s, 3H), 1.65-1.82 (m, 1H), 2.30-2.43 (m, 2H), 2.46 (s, 3H), 2.72 (dd, 1H), 2.78-3.00 (m, 3H), 3.17 (t, 2H), 3.46-3.57 (m, 1H), 4.17 (q, 2H), 4.27 (t, 2H), 6.71 (d, 1H), 6.78 (s, 1H), 7.04 (d, 1H), 7.55 (AB quartet, 4H).

Example 175

Preparation of ((1S)-5-{2-[2-(4-isopropylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid




To a solution of Example 174 (305 mg, 0.657 mmol) in a mixture of THF (8 mL), water (8 mL), and EtOH (4 mL), was added LiOH (63 mg, 2.63 mmol). The reaction mixture was vigorously stirred for 24 hours, diluted with water (20 mL), and washed with Et₂O (10 mL). The aqueous phase was then acidified to pH ~1 using 1 N HCl, and then extracted with CH₂Cl₂ (4 x 50 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was then purified by silica gel flash chromatography (95:5 CH₂Cl₂/MeOH) to afford product (189 mg, 66%) as a white solid. (C₂₆H₂₉NO₃S): LC-MS, RT 3.95 min., M+H 436.4; ¹H NMR (CDCl₃): δ 1.25 (s, 3H), 1.28 (s, 3H), 1.70-1.82 (m, 1H), 2.32-2.43 (m, 2H), 2.45 (s, 3H), 2.74-2.98 (m, 4H),

3.18 (t, 2H), 3.47-3.54 (m, 1H), 4.28 (t, 2H), 6.72 (dd, 1H), 6.78 (s, 1H), 7.08 (d, 1H), 7.51 (AB quartet, 4H).

Example 176


Preparation of methyl [5-methyl-2-(4-methylphenyl)-1,3-thiazol-4-yl]acetate



To a solution of bromide of Example 170 (1.15 g, 5.52 mmol) in toluene (20 mL) was added 4-methyl thiobenzamide (1.0 g, 6.6 mmol). The reaction mixture was heated to reflux for 15 hours, cooled to rt, diluted with EtOAc (150 mL), and washed with a saturated solution of NaHCO₃ (50 mL), then with a saturated solution of NH₄Cl (50 mL). The organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was then purified by silica gel flash chromatography (9:1 hexanes/EtOAc) to afford the product as a pinkish oil that solidified upon standing (1.14 g, 62%). ¹H NMR (CDCl₃): δ 2.38 (s, 3H), 3.45 (s, 3H), 3.74 (s, 3H), 3.80 (s, 2H), 7.49 (AB quartet, 4H); R_f (0.4, eluant 9:1 hexanes/EtOAc).

Example 177

Preparation of 2-[5-methyl-2-(4-methylphenyl)-1,3-thiazol-4-yl]ethanol




To a solution of the thiazole of Example 176 (1.14 g, 4.37 mmol) in THF (60 mL) at 0°C, was added portion-wise LiAlH₄ (663 mg, 17.5 mmol). After 30 minutes, the reaction mixture was warmed to rt and stirred for an additional 60 minutes. The reaction mixture was then cooled to 0°C, and the excess LiAlH₄ was quenched by dropwise addition of water (5 mL), 1N NaOH (10 mL), and water (5 mL) sequentially. The mixture was then diluted with a saturated solution of Rochelle salt and extracted with EtOAc (4 x 75 mL). The combined organic phases were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by silica gel flash chromatography (3:2 hexanes/EtOAc) to afford the product as a white solid (830 mg, 82%). (C₁₃H₁₅NOS): LC-MS, RT 2.50 min., M+H 234.2; ¹H NMR (CDCl₃): δ 2.34 (s, 3H), 2.37 (s, 3H), 2.83 (t, 2H), 3.92-4.01 (broad t, 2H), 4.04-4.15 (broad s, 1H), 7.45 (AB quartet, 4H).

The following compounds below were synthesized using one of the two procedures of Examples 170-177 described above.

Example 178


((1S)-5-[2-(5-Methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl)acetic acid



(C₂₃H₂₃NO₃S): LC-MS RT 3.56 min., M + H 394.2; ¹H NMR (CDCl₃): δ 1.61-1.78 (m, 1H), 2.19-2.50 (m, 2H), 2.30 (s, 3H), 2.62-2.91 (m, 3H), 3.12 (t, 2H), 3.17-3.26 (m, 1H), 4.12 (t, 2H), 6.70 (d, 1H), 6.79 (s, 1H), 6.98 (d, 1H), 7.21-7.40 (m, 3H), 7.74-7.83 (m, 2H).

Example 179


((1S)-5-[2-[5-Methyl-2-(4-methylphenyl)-1,3-thiazol-4-yl]ethoxy]-2,3-dihydro-1H-inden-1-yl)acetic acid




(C₂₄H₂₅NO₃S): LC-MS, RT 3.57 min., M+H 408.5; ¹H NMR (CDCl₃): δ 1.61-1.68 (m, 1H), 2.29 (s, 3H), 2.36 (s, 3H), 2.25-2.37[hidden] (m, 2H), 2.63-2.79 (m, 3H), 3.09 (t, 2H), 3.35-3.47 (m, 1H), 4.18 (t, 2H), 6.60 (dd, 1H), 6.68 (s, 1H), 6.97 (d, 1H), 7.42 (AB quartet, 4H), 7.81-8.30 (br, 1H).

Example 180


((1S)-5-[2-[2-(1,3-Benzodioxol-5-yl)-5-methyl-1,3-thiazol-4-yl]ethoxy]-2,3-dihydro-1H-inden-1-yl)acetic acid




(C₂₄H₂₃NO₅S): LC-MS, RT 4.04 min., M+H 438.5; ¹H NMR (CDCl₃): δ 1.71-1.83 (m, 1H), 2.36-2.51 (m, 2H), 2.45 (s, 3H), 2.76-2.96 (m, 3H), 3.15 (t, 2H), 3.48-3.58 (m, 1H), 4.29 (t, 2H), 6.00 (s, 2H), 6.72 (dd, 1H), 6.78 (s, 1H), 6.82 (d, 1H), 7.07 (d, 1H), 7.32-7.40 (m, 2H).

Example 181**[(1S)-5-{2-[2-(4-Methoxyphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₄H₂₅NO₄S): LC-MS, RT 4.01 min., M+H 424.5; ¹H NMR (CDCl₃): δ 1.67-1.82 (m, 1H), 2.43 (s, 3H), 2.34-2.47 (m, 2H), 2.72-2.95 (m, 3H), 3.09 (t, 2H), 3.42-3.57 (m, 1H), 3.84 (s, 3H), 4.13 (t, 2H), 6.72 (d, 1H), 6.79 (s, 1H), 7.12 (d, 1H), 7.37 (AB quartet, 4H).

Example 182**[(1S)-5-(2-{5-Methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-4-yl}ethoxy)-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₄H₂₂F₃NO₃S): LC-MS, RT 4.47 min., M+H 462.4; ¹H NMR (DMSO-*d*₆): δ 1.63-1.81 (m, 1H), 2.28-2.43 (m, 2H), 2.50 (s, 3H), 2.69 (dd, 1H), 2.74-2.95 (m, 2H), 3.19 (t, 2H), 3.31-3.36 (m, 1H), 4.31 (t, 2H), 6.71 (dd, 1H), 6.78 (s, 1H), 7.08 (d, 1H), 7.87 (AB quartet, 4H).

Example 183**[(1S)-5-{2-[2-(4-Cyanophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₄H₂₂N₂O₃S): LC-MS, RT 3.43 min., M+H 419.6; ¹H NMR (CDCl₃): δ 1.68-1.85 (m, 1H), 2.31-2.49 (m, 2H), 2.51 (s, 3H), 2.77 (dd, 1H), 2.83-2.94 (m, 2H), 3.18 (t, 2H), 3.43-3.56 (m, 1H), 4.31 (t, 2H), 6.71 (dd, 1H), 6.79 (s, 1H), 7.10 (d, 1H), 7.86 (AB quartet, 4H).

Example 184**((1S)-5-{2-[2-(4-Isopropylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**


(C₂₆H₂₉NO₃S): LC-MS, RT 3.95 min., M+H 436.4; ¹H NMR (CDCl₃): δ 1.25 (s, 3H), 1.28 (s, 3H), 1.70-1.82 (m, 1H), 2.32-2.43 (m, 2H), 2.45 (s, 3H), 2.74-2.98 (m, 4H), 3.18 (t, 2H), 3.47-3.54 (m, 1H), 4.28 (t, 2H), 6.72 (dd, 1H), 6.78 (s, 1H), 7.08 (d, 1H), 7.51 (AB quartet, 4H).

Example 185**((1S)-5-{2-[2-(3-Chloro-4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**


(C₂₃H₂₁ClFNO₃S): LC-MS, RT 3.89 min., M+H 446.4; ¹H NMR (CDCl₃): δ 1.68-1.86 (m, 1H), 2.32-2.46 (m, 2H), 2.50 (s, 3H), 2.80 (dd, 1H), 2.84-2.96 (m, 2H), 3.18 (t, 2H), 3.47-3.59 (m, 1H), 4.32 (t, 2H), 6.72 (d, 1H), 6.82 (s, 1H), 7.12 (d, 1H), 7.23 (t, 1H), 7.72-7.82 (m, 1H), 7.97-8.04 (m, 1H).

Example 186**((1S)-5-{2-[2-(3,4-Dichlorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**


(C₂₃H₂₁Cl₂NO₃S): LC-MS, RT 4.12 min., M+H 462.0; ¹H NMR (CDCl₃): δ 1.74-1.88 (m, 1H), 2.36-2.48 (m, 2H), 2.50 (s, 3H), 2.73-2.93 (m, 3H), 3.19 (t, 2H), 3.48-3.55 (m, 1H), 4.30 (t, 2H), 6.71 (d, 1H), 6.79 (s, 1H), 7.09 (d, 1H), 7.52 (d, 1H), 7.61 (dd, 1H), 8.02 (d, 1H).

Example 187**((1S)-5-{2-[2-(4-Fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**


(C₂₃H₂₂FNO₃S): LC-MS, RT 3.58 min., M+H 412.4; ¹H NMR (CDCl₃): δ 1.70-1.77 (m, 1H), 2.37-2.45 (m, 1H), 2.44 (s, 3H), 2.70-2.90 (m, 4H), 3.16 (t, 2H), 3.47-3.52 (m, 1H), 4.27 (t, 2H), 6.70 (d, 1H), 6.76 (s, 1H), 7.00-7.10 (m, 3H), 7.82-7.87 (m, 2H).

Example 188**((1S)-5-{2-[2-(3,4-Dimethylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**

(C₂₅H₂₇NO₃S): LC-MS, RT 4.39 min., M+H 422.3; ¹H NMR (CDCl₃): δ 1.70-1.83 (m, 1H), 2.29 (s, 3H), 2.32 (s, 3H), 2.37-2.50 [hidden] (m, 2H), 2.46 (s, 3H), 2.70-2.90 (m, 3H), 3.32 (t, 2H), 3.45-3.60 (m, 1H), 4.30 (t, 2H), 6.73 (d, 1H), 6.79 (s, 1H), 7.07 (d, 1H), 7.17 (d, 1H), 7.59 (d, 1H), 7.68 (s, 1H).

Example 189**((1S)-5-{2-[2-(4-Acetylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**


(C₂₅H₂₅NO₄S): LC-MS, RT 4.01 min., M+H 436.3; ¹H NMR (CDCl₃): δ 1.70-1.82 (m, 1H), 2.37-2.49 (m, 2H), 2.50 (s, 3H), 2.63 (s, 3H), 2.70-2.90 (m, 3H), 3.20 (t, 2H), 3.45-3.60 (m, 1H), 4.30 (t, 2H), 6.72 (d, 1H), 6.78 (s, 1H), 7.08 (d, 1H), 7.95-8.03 (m, 4H).

Example 190**[(1S)-5-(2-{2-[4-(Dimethylamino)phenyl]-5-methyl-1,3-thiazol-4-yl}ethoxy)-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₅H₂₈N₂O₃S): LC-MS, RT 2.95 min., M+H 437.2; ¹H NMR (DMSO-*d*₆): δ 1.53-1.65 (m, 1H), 2.12-2.24 (m, 2H), 2.36 (s, 3H), 2.63-2.84 (m, 3H), 2.94 (s, 6H), 3.03 (t, 2H), 3.27-3.38 (m, 1H), 4.18 (t, 2H), 6.65 (d, 1H), 6.75 (s, 1H), 7.08 (d, 1H), 7.17 (AB quartet, 4H).

Example 191**[(1S)-5-(2-[2-(3-Amino-4-methylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy)-2,3-dihydro-1H-inden-1-yl]acetic acid**


TFA




C₂₄H₂₆N₂O₃S.C₂F₃O₂: LC-MS, RT 3.5 min., M+H 423.3; ¹H NMR (CD₃OD): δ 1.67-1.82 (m, 1H), 2.25-2.37 (m, 2H), 2.38 (s, 3H), 2.50 (s, 3H), 2.67-2.90 (m, 3H), 3.20 (t, 2H), 3.41-3.56 (m, 1H), 4.32 (t, 2H), 6.71 (d, 1H), 6.79 (s, 1H), 7.09 (d, 1H), 7.42 (d, 1H), 7.69 (dd, 1H), 7.77 (d, 1H).

Example 192**[(1S)-5-(2-[2-(2-Fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy)-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₃H₂₂FNO₃S): LC-MS, RT 4.25 min., M+H 412.2; ¹H NMR (CDCl₃): δ 1.70-1.82 (m, 1H), 2.37-2.48 (m, 2H), 2.49 (s, 3H), 2.74-2.94 (m, 3H), 3.21 (t, 2H), 3.42-3.60 (m, 1H), 4.31 (t, 2H), 6.72 (d, 1H), 6.79 (s, 1H), 7.06-7.35 (m, 4H), 8.21 (t, 1H).

Example 193**((1S)-5-{2-[2-(4-Chlorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**


(C₂₃H₂₂ClNO₃S): LC-MS, RT 4.44 min., M+H 428.2; ¹H NMR (CDCl₃): δ 1.70-1.81 (m, 1H), 2.35-2.45 (m, 2H), 2.46 (s, 3H), 2.74-2.89 (m, 3H), 3.17 (t, 2H), 3.42-3.60 (m, 1H), 4.28 (t, 2H), 6.71 (d, 1H), 6.77 (s, 1H), 7.07 (d, 1H), 7.36 (d, 2H), 7.79 (d, 2H).

Example 194**((1S)-5-{2-[2-(4-Ethoxyphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**


(C₂₅H₂₇NO₄S): LC-MS, RT 3.55 min., M+H 438.5; ¹H NMR (CDCl₃): δ 1.40 (t, 3H), 1.70-1.82 (m, 1H), 2.35-2.47 (m, 2H), 2.45 (s, 3H), 2.74-2.89 (m, 3H), 3.20 (t, 2H), 3.42-3.59 (m, 1H), 4.07 (q, 2H), 4.29 (t, 2H), 6.71 (d, 1H), 6.76 (s, 1H), 6.91 (d, 1H), 7.06 (d, 2H), 7.82 (d, 2H).

Example 195**((1S)-5-{2-[2-(3,4-Dimethoxyphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid**


(C₂₅H₂₇NO₅S): LC-MS, RT 3.86 min., M+H 454.2; ¹H NMR (CDCl₃): δ 1.67-1.82 (m, 1H), 2.37-2.48 (m, 2H), 2.49 (s, 3H), 2.71-2.87 (m, 3H), 3.27 (t, 2H), 3.42-3.57 (m, 1H), 3.93 (s, 3H), 3.96 (s, 3H), 4.29 (t, 2H), 6.35-6.64 (broad s, 1H), 6.67 (d, 1H), 6.75 (s, 1H), 6.89 (d, 1H), 7.05 (d, 1H), 7.39 (d, 1H), 7.56 (s, 1H).

Example 196**[(1S)-5-{2-[5-Methyl-2-(3-methylphenyl)-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₄H₂₅NO₃S): LC-MS, RT 3.71 min., M+H 408.2; ¹H NMR (CDCl₃): δ 1.70-1.82 (m, 1H), 2.38-2.52 (m, 2H), 2.40 (s, 3H), 2.47 (s, 3H), 2.75-2.87 (m, 3H), 3.19 (t, 2H), 3.45-3.60 (m, 1H), 4.29 (t, 2H), 6.72 (d, 1H), 6.78 (s, 1H), 7.07 (d, 1H), 7.19 (d, 1H), 7.30 (t, 1H), 7.64 (d, 1H), 7.75 (s, 1H).

Example 197**[(1S)-5-(2-{5-Methyl-2-[3-(trifluoromethyl)phenyl]-1,3-thiazol-4-yl}ethoxy)-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₄H₂₂F₃NO₃S): LC-MS, RT 3.90 min., M+H 462.1; ¹H NMR (CDCl₃): δ 1.70-1.82 (m, 1H), 2.38-2.48 (m, 2H), 2.49 (s, 3H), 2.75-2.87 (m, 3H), 3.19 (t, 2H), 3.44-3.59 (m, 1H), 4.30 (t, 2H), 6.72 (d, 1H), 6.79 (s, 1H), 7.07 (d, 1H), 7.52 (t, 1H), 7.61 (d, 1H), 8.01 (d, 1H), 8.13 (s, 1H).

Example 198**[(1S)-5-{2-[2-(3-Fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₃H₂₂FNO₃S): LC-MS, RT 3.66 min., M+H 412.1; ¹H NMR (CDCl₃): δ 1.70-1.82 (m, 1H), 2.39-2.47 (m, 2H), 2.48 (s, 3H), 2.76-2.87 (m, 3H), 3.18 (t, 2H), 3.45-3.60 (m, 1H), 4.30 (t, 2H), 6.72 (d, 1H), 6.78 (s, 1H), 7.04-7.09 (m, 2H), 7.36-7.42 (m, 1H), 7.58-7.62 (m, 2H).

Example 199**[(1S)-5-{2-[2-(3,5-Dimethylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₅H₂₇NO₃S): LC-MS, RT 3.88 min., M+H 422.2; ¹H NMR (CDCl₃): δ 1.72-1.84 (m, 1H), 2.36 (s, 6H), 2.37-2.45 (m, 2H), 2.46 (s, 3H), 2.75-2.87 (m, 3H), 3.19 (t, 2H), 3.45-3.60 (m, 1H), 4.28 (t, 2H), 6.72 (d, 1H), 6.79 (s, 1H), 7.01 (s, 1H), 7.07 (d, 1H), 7.48 (s, 2H).

Example 200**[(1S)-5-(2-{5-Methyl-2-[4-(trifluoromethoxy)phenyl]-1,3-thiazol-4-yl}ethoxy)-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₄H₂₂F₃NO₄S): LC-MS, RT 3.95 min., M+H 478.1; ¹H NMR (CDCl₃): δ 1.72-1.84 (m, 1H), 2.38-2.46 (m, 2H), 2.47 (s, 3H), 2.75-2.87 (m, 3H), 3.18 (t, 2H), 3.45-3.60 (m, 1H), 4.29 (t, 2H), 6.72 (d, 1H), 6.77 (s, 1H), 7.07 (d, 1H), 7.24 (d, 2H), 7.88 (d, 2H).

Example 201**[(1S)-5-{2-[2-(3-Methoxyphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl]acetic acid**


(C₂₄H₂₅NO₄S): LC-MS, RT 3.56 min., M+H 424.2; ¹H NMR (CDCl₃): δ 1.70-1.82 (m, 1H), 2.37-2.52 (m, 2H), 2.49 (s, 3H), 2.75-2.87 (m, 3H), 3.19 (t, 2H), 3.45-3.57 (m, 1H), 3.87 (s, 3H), 4.30 (t, 2H), 6.72 (d, 1H), 6.79 (s, 1H), 6.95 (d, 1H), 7.10 (d, 1H), 7.32 (t, 1H), 7.40-7.45 (m, 2H).

Example 202**((1S)-5-[2-[2-(1,1'-Biphenyl-4-yl)-5-methyl-1,3-thiazol-4-yl]ethoxy]-2,3-dihydro-1H-inden-1-yl)acetic acid**

(C₂₉H₂₇NO₃S): LC-MS, RT 3.96 min., M+H 470.3; ¹H NMR (CDCl₃): δ 1.70-1.81 (m, 1H), 2.38-2.48 (m, 2H), 2.49 (s, 3H), 2.75-2.87 (m, 3H), 3.20 (t, 2H), 3.43-3.59 (m, 1H), 4.31 (t, 2H), 6.72 (d, 1H), 6.79 (s, 1H), 7.08 (d, 1H), 7.36 (t, 1H), 7.45 (t, 2H), 7.61-7.65 (m, 4H), 7.93 (d, 2H).


Example 203**Preparation of ethyl {(1S)-5-[2-(4-methyl-2-phenyl-1,3-oxazol-5-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}acetate**

ADDP (0.205 g, 0.81 mmol) was added to a mixture of PPh₃ (0.212 g, 0.81 mmol), ethyl [(1S)-5-hydroxy-2,3-dihydro-1H-inden-1-yl]acetate (0.107 g, 0.49 mmol), and 2-(4-methyl-2-phenyl-1,3-oxazol-5-yl)ethanol (step 4, Example 51, 0.110 g, 0.54 mmol) in THF (5 mL). The reaction was stirred overnight at rt, and additional ADDP (0.136 g, 0.54 mmol) and PPh₃ (0.141 g, 0.54 mmol) were added with CH₂Cl₂ (5 mL). The solution was stirred for 24 hours at rt and filtered. The filtrate was evaporated and the resulting mixture was purified by Biotage using a gradient 0 to 50% EtOAc/hexane. Gave ethyl {(1S)-5-[2-(4-methyl-2-phenyl-1,3-oxazol-5-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}acetate (0.145 g, 66% yield) as yellowish oil. ES-MS *m/z* 406.2 ((MH)⁺); HPLC RT (min.) 3.89; ¹H NMR (Acetone-*d*₆) δ 7.85-7.82 (m, 2H), 7.36-7.30 (m, 3H), 6.94 (d, 1H), 6.65 (s, 1H), 6.60-6.55 (m, 1H), 4.10 (t, 2H), 3.98 (q, 2H), 3.31-3.27 (m, 1H), 3.03 (t, 2H), 3.27-2.51 (m, 3H), 2.24-2.14 (m, 2H), 2.18 (s, 3H), 1.58-1.53 (m, 1H), 1.08 (t, 3H).


Example 204**Preparation of {(1S)-5-[2-(4-methyl-2-phenyl-1,3-oxazol-5-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}acetic acid**

Ethyl {(1S)-5-[2-(4-methyl-2-phenyl-1,3-oxazol-5-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}acetate (0.135 g, 0.33 mmol) was dissolved in EtOH (6 mL) and LiOH (0.024 g, 1.0 mmol) was added. Water (3 mL) was added and THF was added until the cloudy solution became clear. The resulting mixture was stirred overnight at rt. HCl (2 N) was added to adjust the pH to 2, then extracted three times with ethyl acetate. The organic layers were combined, dried, and concentrated to give {(1S)-5-[2-(4-methyl-2-phenyl-1,3-oxazol-5-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl}acetic acid (0.039 g, 30.6% yield) as colorless oil. ES-MS m/z 378.2 ((MH)⁺); HPLC RT (min.) 3.22; ¹H NMR (Acetone-*d*₆) δ 8.1 (s br 1H) 8.0-7.95 (m, 2H), 7.52-7.43 (m, 3H), 7.15(d, 1H), 6.81 (s, 1H), 6.73 (d, 1H), 4.27 (t, 2H) 3.47-3.40 (m, 1H), 3.18 (t, 2H), 2.90-2.68 (m, 3H), 2.41-2.29 (m, 2H), 2.18 (s, 3H), 1.77-1.68 (m, 1H).


By using the procedure described above for Examples 51, 203, and 204 and substituting the appropriate starting materials, the following compounds were similarly prepared and characterized.

Example 205**Preparation of N-(4-methylbenzoyl)alanine**


¹H NMR (DMSO-*d*₆) δ 12.60 (s br, 1H), 8.57 (d, 1H), 7.81 (d, 2H), 7.28 (d, 2H), 4.38 (q, 1H), 2.35 (s, 3H), 1.38 (d, 3H).

Example 206**Preparation of *N*-(3-fluoro-4-methylbenzoyl)alanine**


^1H NMR (DMSO- d_6) δ 12.54 (s br, 1H), 8.67 (d, 1H), 7.65-7.62 (m, 2H), 7.39 (t, 1H), 4.38 (q, 1H), 2.27 (s, 3H), 1.38 (d, 3H).

Example 207**Preparation of *N*-[4-(trifluoromethyl)benzoyl]alanine**


^1H NMR (DMSO- d_6) δ 12.64 (s br, 1H), 8.91 (d, 1H), 8.08 (d, 2H), 7.85 (d, 2H), 4.42 (q, 1H), 1.40 (d, 3H).

Example 208**Preparation of ethyl 4-[(4-methylbenzoyl)amino]-3-oxopentanoate**


ES-MS m/z 278.38 ($(\text{MH})^+$); HPLC RT (min.) 2.04. ^1H NMR (Acetone- d_6) δ 8.08 (s br, 1H), 7.90 (d, 2H), 7.28 (d, 2H), 4.72-4.67 (m, 1H), 4.13 (q, 2H), 3.66 (s, 2H), 2.40 (s, 3H), 1.41 (d, 3H), 1.12 (t, 3H).

Example 209**Preparation of ethyl 4-[(3-fluoro-4-methylbenzoyl)amino]-3-oxopentanoate**


ES-MS m/z 296.4 ($(\text{MH})^+$); HPLC RT (min.) 2.26. ^1H NMR (Acetone- d_6) δ 7.75-7.60 (m, 2H), 7.38 (t, 1H), 4.20 (q, 2H), 3.65 (s, 2H), 2.23 (s, 3H), 1.45 (d, 3H), 1.20 (t, 3H).

Example 210**Preparation of ethyl 3-oxo-4-{[4-(trifluoromethyl)benzoyl]amino}pentanoate**


ES-MS m/z 332.4 ((MH)⁺); HPLC RT (min.) 2.45. ¹H NMR (Acetone-d₆) δ 8.14 (d, 2H), 7.84 (d, 2H), 4.80-4.74 (m, 2H), 4.20 (q, 2H), 3.70 (s, 2H), 1.48 (d, 3H), 1.21 (t, 3H).

Example 211**Preparation of ethyl [4-methyl-2-(4-methylphenyl)-1,3-oxazol-5-yl]acetate**

ES-MS m/z 260.2 ((MH)⁺); HPLC RT (min.) 2.96. ¹H NMR (Acetone-d₆) δ 7.86 (d, 2H), 7.30 (d, 2H), 4.15 (q, 2H), 3.81 (s, 2H), 2.37 (s, 3H), 2.14 (s, 3H), 1.24 (t, 3H).

Example 212**Preparation of ethyl [2-(3-fluoro-4-methylphenyl)-4-methyl-1,3-oxazol-5-yl]acetate**


ES-MS m/z 278.3 ((MH)⁺); HPLC RT (min.) 2.89. ¹H NMR (Acetone-d₆) δ 7.69 (d, 1H), 7.60 (d, 1H), 7.37 (t, 1H), 4.15 (q, 2H), 3.83 (s, 2H), 2.31 (s, 3H), 2.15 (s, 3H), 1.23 (t, 3H).

Example 213**Preparation of ethyl {4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-5-yl}acetate**

ES-MS m/z 314.3 ((MH)⁺); HPLC RT (min.) 3.27. ¹H NMR (Acetone-d₆) δ 8.18 (d, 2H), 7.84 (d, 2H), 4.17 (q, 2H), 3.88 (s, 2H), 2.20 (s, 3H), 1.23 (t, 3H).

Example 214


Preparation of 2-[4-methyl-2-(4-methylphenyl)-1,3-oxazol-5-yl]ethanol



ES-MS m/z 218.2 ((MH)⁺); HPLC RT (min.) 2.35. ¹H NMR (Acetone d₆) δ 7.85 (d, 2H), 7.27 (d, 2H), 3.99 (s br, 1H), 3.83 (t, 2H), 2.90 (t, 2H), 2.37 (s, 3H), 2.12 (s, 3H).

Example 215


Preparation of 2-[2-(3-fluoro-4-methylphenyl)-4-methyl-1,3-oxazol-5-yl]ethanol




ES-MS m/z 236.2 ((MH)⁺); HPLC RT (min.) 2.46. ¹H NMR (CDCl₃) δ 7.54 (d, 1H), 7.43 (d, 1H), 7.17 (t, 1H), 3.91 (d, 2H), 3.09 (s br, 1H), 2.88 (t, 2H), 2.29 (s, 3H), 2.13 (s, 3H).

Example 216


Preparation of 2-[4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-5-yl]ethanol




ES-MS m/z 272.2 ((MH)⁺); HPLC RT (min.) 2.71. ¹H NMR (CDCl₃) δ 8.03 (2, 2H), 7.66 (d, 2H), 3.95 (t, 2H), 2.96 (t, 2H), 2.21 (s, 3H), 1.97 (s br, 1H).

Example 217**Preparation of ethyl [(1S)-5-(2-{4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-5-yl}ethoxy)-2,3-dihydro-1H-inden-1-yl]acetate**

ES-MS m/z 474.5 ((MH)⁺); HPLC RT (min.) 4.10. ¹H NMR (Acetone-*d*₆) δ 8.16 (d, 2H), 7.83 (d, 2H), 7.09 (d, 1H), 6.80 (s, 1H), 6.72 (dd, 1H), 4.28 (t, 2H), 4.12 (q, 2H), 3.46-3.41 (m, 1H), 3.21 (t, 2H), 2.86-2.65 (m, 3H), 2.39-2.26 (m, 2H), 2.20 (s, 3H), 1.75-1.63 (m, 1H), 1.22 (t, 3H).

Example 218**Preparation of ethyl ((1S)-5-{2-[4-methyl-2-(4-methylphenyl)-1,3-oxazol-5-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetate**


TCL R_f = 0.22 Hexane/EtOAc 4:1

Example 219**Preparation of ethyl ((1S)-5-{2-[2-(3-fluoro-4-methylphenyl)-4-methyl-1,3-oxazol-5-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetate**

ES-MS m/z 438.2 ((MH)⁺); HPLC RT (min.) 4.18. ¹H NMR (Acetone-*d*₆) δ 6.67 (dd, 1H), 7.59 (dd, 1H), 7.37 (t, 1H), 7.08 (d, 1H), 6.80 (s, 1H), 6.72 (dd, 1H), 4.26 (t, 2H), 4.12 (q, 2H), 3.46-3.38 (m, 1H), 3.17 (t, 2H), 2.89-2.65 (m, 3H), 2.39-2.23 (m, 5H), 2.17 (s, 3H), 1.75-1.63 (m, 1H), 1.23 (t, 3H).

Example 220


Preparation of ((1S)-5-{2-[4-methyl-2-(4-methylphenyl)-1,3-oxazol-5-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid




ES-MS m/z 392.2 ((MH)⁺); HPLC RT (min.) 3.36. ¹H NMR (Acetone-*d*₆) δ 7.72 (d, 2H), 7.15 (d, 2H), 6.99 (d, 1H), 6.67 (s, 1H), 6.59 (dd, 1H), 4.12 (t, 2H), 3.33-3.28 (m, 1H), 3.03 (t, 2H), 2.73-2.54 (m, 3H), 2.27-2.21 (m, 5H), 2.02 (s, 3H), 1.64-1.54 (m, 1H).

Example 221



Preparation of ((1S)-5-{2-[2-(3-fluoro-4-methylphenyl)-4-methyl-1,3-oxazol-5-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetic acid



ES-MS m/z 410.2 ((MH)⁺); HPLC RT (min.) 3.49. ¹H NMR (Acetone-*d*₆) δ 7.68 (dd, 1H), 7.59 (dd, 1H), 7.36 (t, 1H), 7.12 (d, 1H), 6.80 (s, 1H), 6.72 (dd, 1H), 4.26 (t, 2H), 3.47-3.41 (m, 1H), 3.18 (t, 2H), 2.86-2.67 (m, 3H), 2.40-2.28 (m, 5H), 2.17 (s, 3H), 1.18-1.65 (m, 1H).

Example 222**Preparation of [(1S)-5-(2-{4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-5-yl}ethoxy)-2,3-dihydro-1H-inden-1-yl]acetic acid**


ES-MS m/z 446.5 ($(MH)^+$); HPLC RT (min.) 3.47. 1H NMR (Acetone- d_6) δ 8.17 (d, 2H), 7.84 (d, 2H), 7.13 (s, 1H), 6.80 (s, 1H), 6.72 (dd, 1H), 4.28 (t, 2H), 3.46-3.41 (m, 1H), 3.21 (t, 2H), 2.86-2.67 (m, 3H), 2.40-2.28 (m, 2H), 2.20 (s, 3H), 1.77-1.67 (m, 1H).

Example 223**Preparation of (2S)-2-[(1S)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl]propanoic acid and (2R)-2-[(1R)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl]propanoic acid****Step 1. Preparation of (2S)-2-[(1S)-5-methoxy-2,3-dihydro-1H-inden-1-yl]propanoic acid and (2R)-2-[(1R)-5-methoxy-2,3-dihydro-1H-inden-1-yl]propanoic acid**

The starting acid (Example 2b) was reacted using a similar procedure as described in Example 4, under 60 psi H_2 , and using 4.5 g starting material, 1.04 g catalyst, and 4.5 mL triethylamine in 45 mL ethanol and 5 mL THF. The standard extractive workup gave 3.22 g product. LC/MS retention time 2.41 min., NMR (d_6 -


DMSO): 0.87 (d, 3H, α -methyl), 1.75 (m, 1H), 2.04 (m, 1H), 3.66 (s, 3H, methoxy), 6.65 (m, 1H, aryl), 6.76 (s, 1H, aryl), 7.04 (d, 1H, aryl), 12.18 (bs, 1H, acid.)

Step 2: Preparation of methyl (2S)-2-[(1S)-5-methoxy-2,3-dihydro-1H-inden-1-yl]propanoate and methyl (2R)-2-[(1R)-5-methoxy-2,3-dihydro-1H-inden-1-yl]propanoate




The compound was prepared by the reaction of 1.5 g starting acid, 0.93 mL iodomethane, and 1.75 g sodium bicarbonate in 10 mL methanol under standard esterification conditions as described in Example 6. Workup gave 1.53 g, 96%. (NMR (CD_2Cl_2): 1.05 (d, 3H, α -methyl), 1.88 (m, 1H), 2.19 (m, 1H), 3.44 (m, 1H), 3.68 (s, 3H, methoxy), 3.77 (s, 3H, ester).

Step 3. Preparation of: methyl (2S)-2-[(1S)-5-hydroxy-2,3-dihydro-1H-inden-1-yl]propanoate and methyl (2R)-2-[(1R)-5-hydroxy-2,3-dihydro-1H-inden-1-yl]propanoate




Using the demethylation conditions as described in Example 7 (1.53 g starting material, 4.35 g AlCl_3 , and 2.4 mL ethanethiol in 20 mL dichloromethane), 1.21 g of product (84%) was obtained. (NMR (CD_2Cl_2): 1.05 (d, 3H, α -methyl), 1.88 (m, 1H), 2.18 (m, 1H), 3.45 (m, 1H), 3.67 (s, 3H, ester), 6.60 (m, 1H, aryl), 6.69 (s, 1H, aryl), 6.93 (d, 1H, aryl).

Step 4: Preparation of methyl (2S)-2-[(1S)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl]propanoate and methyl (2R)-2-[(1R)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl]propanoate




Using the standard Mitsunobu coupling procedure as described in Example 11 (0.100 g starting phenol, 0.110 g oxazolyethanol, 0.143 g triphenylphosphine, and 0.137 g ADDP in 2 mL dichloromethane), 0.107 g (58%) of product was obtained after chromatography in 15% EtOAc/hexane. NMR (CD₂Cl₂): 1.62-1.87 (m, 4H), 2.40 (s, 3H, oxazole methyl), 2.98 (t, 2H, methylene), 3.23 (m, 1H), 3.63 (s, 3H, ester), 6.60 (s, 1H, aryl), 6.64 (m, 1H, aryl), 7.42 (m, 3H, aryl), 8.00 (m, 2H, aryl).

Step 5. (2S)-2-[(1S)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl]propanoic acid and (2R)-2-[(1R)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1H-inden-1-yl]propanoic acid




The LiOH hydrolysis conditions were applied to 0.090 g of starting ester, yielding 0.082 g (95%) product. NMR (CD₃OD): 0.4-0.75 (m, 4H), 1.18 (s, 3H), 1.75 (t, 2H, methylene), 2.00 (m, 1H), 2.99 (t, 2H, methylene), 5.39 (s, 1H, aryl), 5.48 (m, 1H, aryl), 5.83 (d, 1H, aryl), 6.27 (m, 3H, aryl), 6.76 (m, 2H, aryl).

Using the methods described above and the appropriate starting materials, additional (2S, 1S) and (2R, 1R) were similarly prepared, either as diastereomeric (i.e., syn, {(2S, 1S)/(2R, 1R)}) and or anti {(2R, 1S)/(2S, 1R)}) mixtures, or as individual enantiomers. These compounds are summarized in Table 5.


Table 5

Ex. No.	R ³	R ⁴	X	Isomer	HPLC RT (min)	LC-MS [M+H] ⁺
224	Me	3,4-(Cl) ₂ -Ph	O	2S,1S	4.10	460.0
225	Me	3,4-(Cl) ₂ -Ph	O	syn racemate	4.10	460.0
226	Me	3,4-(Me) ₂ -Ph	O	syn racemate	4.32	420.4
227	Me	3,4-(Me) ₂ -Ph	O	2S,1S	4.32	420.4
228	Me	3-Me-Ph	O	syn racemate	4.19	406.3
229	Me	4-CF ₃ -Ph	O	syn racemate	3.73	460.2
230	Me	4-CF ₃ -Ph	O	2S,1S	3.73	460.2
231	Me	4-CF ₃ -Ph	O	2R,1R	3.73	460.2
232	Me	4-Cl-Ph	O	syn racemate	3.61	426.2
233	Me	4-Et-Ph	O	syn racemate	3.70	420.3
234	Me	4-Et-Ph	O	2S,1S	3.70	420.3
235	Me	4-Et-Ph	O	2R,1R	3.70	420.3
236	Me	4-Et-Ph	O	syn/anti mixture	3.70	420.3
237	Me	4-Et-Ph	O	2R,1S	3.70	420.3
238	Me	4-Et-Ph	O	2S,1R	3.70	420.3
239	Me	4-MeO-Ph	O	syn racemate	3.37	422.3
240	Me	4-MeO-Ph	O	2R,1R	3.37	422.3
241	Me	4-MeO-Ph	O	2S,1S	3.37	422.3
242	Me	4-n-Bu-Ph	O	syn racemate	4.08	448.4
243	Me	4-t-Bu-Ph	O	2S,1S	4.59	448.4
244	Et	4-t-Bu-Ph	O	syn racemate	4.59	448.4
245	Me	4-MeO-Ph	O	2S,1S	3.58	-
246	Me	4-Cl-Ph	S	syn racemate	3.84	442.2
247	Me	4-Me-Ph	S	syn racemate	4.34	422.3

Example 248**Preparation of ethyl [(1S)-5-(2-{2-[4'-(5-acetyl-2-thienyl)-1,1'-biphenyl-4-yl]-5-methyl-1,3-oxazol-4-yl}ethoxy)-2,3-dihydro-1H-inden-1-yl]acetate**

To a solution containing ethyl ((1S)-5-{2-[2-(4-bromophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1H-inden-1-yl)acetate (0.100 g, 0.21 mmol) [prepared from 2-[5-methyl-2-(4-bromophenyl)-1,3-oxazol-4-yl]ethanol and ethyl [(1S)-5-hydroxy-2,3-dihydro-1H-inden-1-yl]acetate (Example 135)], 1,1'-bis(diphenylphosphino)-ferrocene]dichloro palladium(II) (16.9 mg, 0.02 mmol), and 5-acetyl-2-thienylboronic acid (0.062 g, 0.41 mmol) in degassed toluene and dioxane (4:1, 2 mL) was added aqueous 2 M sodium carbonate (0.5 mL). The mixture was heated at 85°C for 16 hours. Solvents were evaporated under vacuum and the residue was dissolved in methanol and acetonitrile and filtered through a C8 reverse phase extraction cartridge. Solvents were evaporated and the residue was dissolved in acetonitrile and purified by HPLC to obtain ethyl [(1S)-5-(2-{2-[4'-(5-acetyl-2-thienyl)-1,1'-biphenyl-4-yl]-5-methyl-1,3-oxazol-4-yl}ethoxy)-2,3-dihydro-1H-inden-1-yl]acetate in 46% yield. (50 mg, 0.09 mmol) MS (electro spray) 530.4 (M+H)⁺, ¹H NMR (CDCl₃) δ 1.24 (t, 3H), 1.71 (m, 1H), 2.37 (m, 5H), 2.57 (s, 3H), 2.68 (m, 1H), 2.83 (m, 2H), 3.03 (m, 2H), 3.48 (m, 1H), 4.17(m, 4H), 6.67 (m, 2H), 7.02 (d, 1H), 7.39 (d, 1H), 7.67 (d, 1H), 7.73 (d, 2H), 8.01 (d, 2H).

Other compounds, prepared by using analogous starting materials and the method described in Example 248 together with the hydrolysis described in Example 11, are described below in Table 6.

Table 6.

Ex. No.	R	R ¹	R ²	R ³	R ⁴	R ⁵	X	LC-MS [M+H] ⁺
249	H	H	H	Me		H	O	493.3
250	H	H	H	Me		H	O	484.2
251	H	H	H	Me		H	O	502.2

EVALUATION OF COMPOUNDS

Demonstration of the activity of the compounds of the present invention may be accomplished through *in vitro*, *ex vivo*, and *in vivo* assays that are well known in the art. For example, to demonstrate the efficacy of a pharmaceutical agent for the treatment of diabetes and related disorders such as Syndrome X, impaired glucose tolerance, impaired fasting glucose, and hyperinsulinemia or atherosclerotic disease and related disorders such as hypertriglyceridemia and hypercholesteremia, the following assays may be used.

Insulin Receptor Binding in 3T3-L1 Cells Treated with Compounds

3T3-L1 cells were seeded at 9300 cells per well in Costar flat bottom TC and incubated for 1 week until they were 2 days post-confluent (e.g., cells have reached maximum density). The cells were then treated for 2 days with differentiation media (Dulbecco's Modified Eagle Medium (DMEM), 100 µg/ml Penicillin/Streptomycin, 2 mM L-Glutamine, 10% Fetal Bovine Serum) containing 0.5 µM human Insulin-like Growth Factor (IGF-1) and test compounds. After treatment, the media was replaced with differentiation media, and the cells were incubated for 4 days. The cells were then assayed for insulin receptor activity. After washing the cells with buffer, they were incubated with 0.1 nM ¹²⁵I-insulin and (+/-) 100 nM unlabeled insulin, and incubated at rt for 1 hour. The cells were

then washed 3x with buffer, dissolved with 1N NaOH, and counted on a gamma counter. An EC50 value was determined if a plateau was attained and percent maximum stimulation was assessed.

***In Vivo* Assays**

Method for Measuring Blood Glucose Levels

db/db mice (obtained from Jackson Laboratories, Bar Harbor, ME) were bled (by either eye or tail vein) and grouped according to equivalent mean blood glucose levels. They were dosed orally (by gavage in a pharmaceutically acceptable vehicle) with the test compound once daily for 14 days. At this point, the animals were bled again by eye or tail vein and blood glucose levels were determined. In each case, glucose levels were measured with a Glucometer Elite XL (Bayer Corporation, Elkhart, IN).

Method for Measuring Triglyceride Levels

hApoA1 mice (obtained from Jackson Laboratories, Bar Harbor, ME) were bled (by either eye or tail vein) and grouped according to equivalent mean serum triglyceride levels. They were dosed orally (by gavage in a pharmaceutically acceptable vehicle) with the test compound once daily for 8 days. The animals were then bled again by eye or tail vein, and serum triglyceride levels were determined. In each case, triglyceride levels were measured using a Technicon Axon Autoanalyzer (Bayer Corporation, Tarrytown, NY).

Method for Measuring HDL-Cholesterol Levels

To determine plasma HDL-cholesterol levels, hApoA1 mice are bled and grouped with equivalent mean plasma HDL-cholesterol levels. The mice are orally dosed once daily with vehicle or test compound for 7 days, and then bled again on day 8. Plasma is analyzed for HDL-cholesterol using the Synchron Clinical System (CX4) (Beckman Coulter, Fullerton, CA).

Method for Measuring Total Cholesterol, HDL-Cholesterol, Triglycerides, and Glucose Levels

In another *in vivo* assay, obese monkeys are bled, then orally dosed once daily with vehicle or test compound for 4 weeks, and then bled again. Serum is analyzed for total cholesterol, HDL-cholesterol, triglycerides, and glucose using the Synchron Clinical System (CX4) (Beckman Coulter, Fullerton, CA). Lipoprotein subclass analysis is

performed by NMR spectroscopy as described by Oliver et al., (Proc. Natl. Acad. Sci. USA 98:5306-5311, 2001).

Method for Measuring an Effect on Cardiovascular Parameters

Cardiovascular parameters (e.g., heart rate and blood pressure) are also evaluated. SHR rats are orally dosed once daily with vehicle or test compound for 2 weeks. Blood pressure and heart rate are determined using a tail-cuff method as described by Grinsell et al., (Am. J. Hypertens. 13:370-375, 2000). In monkeys, blood pressure and heart rate are monitored as described by Shen et al., (J. Pharmacol. Exp. Therap. 278:1435-1443, 1996).

Compounds of the present invention were tested in the above assays and by the resulting activity profiles, they were found to have an effect on blood glucose levels and serum triglyceride levels, and therefore, a potential utility in the treatment of diabetes and related disorders such as Syndrome X, impaired glucose tolerance, impaired fasting glucose, and hyperinsulinemia or cardiovascular disease and related disorders such as hypertriglyceridemia and hypercholesteremia.

Pharmaceutical Compositions

Based on the above tests, or other well known assays used to determine the efficacy for treatment of conditions identified above in mammals, and by comparison of these results with the results of known medicaments that are used to treat these conditions, the effective dosage of the compounds of this invention can readily be determined for treatment of each desired indication. The amount of the active ingredient to be administered in the treatment of one of these conditions can vary widely according to such considerations as the particular compound and dosage unit employed, the mode of administration, the period of treatment, the age and sex of the patient treated, and the nature and extent of the condition treated.

The total amount of the active ingredient to be administered may generally range from about 0.001 mg/kg to about 200 mg/kg, and preferably from about 0.01 mg/kg to about 200 mg/kg body weight per day. A unit dosage may contain from about 0.05 mg to about 1500 mg of active ingredient, and may be administered one or more times per day. The daily dosage for administration by injection, including intravenous, intramuscular, subcutaneous, and parenteral injections, and use of infusion techniques may be from about 0.01 to about 200 mg/kg. The daily rectal dosage regimen may be from 0.01 to

200 mg/kg of total body weight. The transdermal concentration may be that required to maintain a daily dose of from 0.01 to 200 mg/kg.

Of course, the specific initial and continuing dosage regimen for each patient will vary according to the nature and severity of the condition as determined by the attending diagnostician, the activity of the specific compound employed, the age of the patient, the diet of the patient, time of administration, route of administration, rate of excretion of the drug, drug combinations, and the like. The desired mode of treatment and number of doses of a compound of the present invention or a pharmaceutically acceptable salt thereof may be ascertained by those skilled in the art using conventional treatment tests.

The compounds of this invention may be utilized to achieve the desired pharmacological effect by administration to a patient in need thereof in an appropriately formulated pharmaceutical composition. A patient, for the purpose of this invention, is a mammal, including a human, in need of treatment for a particular condition or disease. Therefore, the present invention includes pharmaceutical compositions which are comprised of a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound identified by the methods described herein, or a pharmaceutically acceptable salt or ester thereof. A pharmaceutically acceptable carrier is any carrier which is relatively non-toxic and innocuous to a patient at concentrations consistent with effective activity of the active ingredient so that any side effects ascribable to the carrier do not vitiate the beneficial effects of the active ingredient. A pharmaceutically effective amount of a compound is that amount which produces a result or exerts an influence on the particular condition being treated. The compounds identified by the methods described herein may be administered with a pharmaceutically-acceptable carrier using any effective conventional dosage unit forms, including, for example, immediate and timed release preparations, orally, parenterally, topically, or the like.

For oral administration, the compounds may be formulated into solid or liquid preparations such as, for example, capsules, pills, tablets, troches, lozenges, melts, powders, solutions, suspensions, or emulsions, and may be prepared according to methods known to the art for the manufacture of pharmaceutical compositions. The solid unit dosage forms may be a capsule which can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers such as lactose, sucrose, calcium phosphate, and corn starch.

In another embodiment, the compounds of this invention may be tableted with conventional tablet bases such as lactose, sucrose, and cornstarch in combination with binders such as acacia, cornstarch, or gelatin; disintegrating agents intended to assist the break-up and dissolution of the tablet following administration such as potato starch,

alginic acid, corn starch, and guar gum; lubricants intended to improve the flow of tablet granulation and to prevent the adhesion of tablet material to the surfaces of the tablet dies and punches, for example, talc, stearic acid, or magnesium, calcium or zinc stearate; dyes; coloring agents; and flavoring agents intended to enhance the aesthetic qualities of the tablets and make them more acceptable to the patient. Suitable excipients for use in oral liquid dosage forms include diluents such as water and alcohols, for example, ethanol, benzyl alcohol, and polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance tablets, pills or capsules may be coated with shellac, sugar or both.

Dispersible powders and granules are suitable for the preparation of an aqueous suspension. They provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent, and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, those sweetening, flavoring and coloring agents described above, may also be present.

The pharmaceutical compositions of this invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil such as liquid paraffin or a mixture of vegetable oils. Suitable emulsifying agents may be (1) naturally occurring gums such as gum acacia and gum tragacanth, (2) naturally occurring phosphatides such as soy bean and lecithin, (3) esters or partial esters derived from fatty acids and hexitol anhydrides, for example, sorbitan monooleate, and (4) condensation products of said partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil such as, for example, arachis oil, olive oil, sesame oil, or coconut oil; or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent such as, for example, beeswax, hard paraffin, or cetyl alcohol. The suspensions may also contain one or more preservatives, for example, ethyl or *n*-propyl *p*-hydroxybenzoate; one or more coloring agents; one or more flavoring agents; and one or more sweetening agents such as sucrose or saccharin.

Syrups and elixirs may be formulated with sweetening agents such as, for example, glycerol, propylene glycol, sorbitol, or sucrose. Such formulations may also contain a demulcent, and preservative, flavoring and coloring agents.

The compounds of this invention may also be administered parenterally, that is,

subcutaneously, intravenously, intramuscularly, or interperitoneally, as injectable dosages of the compound in a physiologically acceptable diluent with a pharmaceutical carrier which may be a sterile liquid or mixture of liquids such as water, saline, aqueous dextrose and related sugar solutions; an alcohol such as ethanol, isopropanol, or hexadecyl alcohol; glycols such as propylene glycol or polyethylene glycol; glycerol ketals such as 2,2-dimethyl-1,1-dioxolane-4-methanol, ethers such as poly(ethyleneglycol) 400; an oil; a fatty acid; a fatty acid ester or glyceride; or an acetylated fatty acid glyceride with or without the addition of a pharmaceutically acceptable surfactant such as a soap or a detergent, suspending agent such as pectin, carbomers, methycellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agent and other pharmaceutical adjuvants.

Illustrative of oils which can be used in the parenteral formulations of this invention are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, sesame oil, cottonseed oil, corn oil, olive oil, petrolatum, and mineral oil. Suitable fatty acids include oleic acid, stearic acid, and isostearic acid. Suitable fatty acid esters are, for example, ethyl oleate and isopropyl myristate. Suitable soaps include fatty alkali metal, ammonium, and triethanolamine salts and suitable detergents include cationic detergents, for example, dimethyl dialkyl ammonium halides, alkyl pyridinium halides, and alkylamine acetates; anionic detergents, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates; nonionic detergents, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylenepolypropylene copolymers; and amphoteric detergents, for example, alkyl-beta-aminopropionates, and 2-alkylimidazoline quarternary ammonium salts, as well as mixtures.

The parenteral compositions of this invention may typically contain from about 0.5% to about 25% by weight of the active ingredient in solution. Preservatives and buffers may also be used advantageously. In order to minimize or eliminate irritation at the site of injection, such compositions may contain a non-ionic surfactant having a hydrophile-lipophile balance (HLB) of from about 12 to about 17. The quantity of surfactant in such formulation ranges from about 5% to about 15% by weight. The surfactant can be a single component having the above HLB or can be a mixture of two or more components having the desired HLB.

Illustrative of surfactants used in parenteral formulations are the class of polyethylene sorbitan fatty acid esters, for example, sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol.

The pharmaceutical compositions may be in the form of sterile injectable aqueous suspensions. Such suspensions may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents such as, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents which may be a naturally occurring phosphatide such as lecithin, a condensation product of an alkylene oxide with a fatty acid, for example, polyoxyethylene stearate, a condensation product of ethylene oxide with a long chain aliphatic alcohol, for example, heptadecaethyleneoxycetanol, a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol such as polyoxyethylene sorbitol monooleate, or a condensation product of an ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride, for example polyoxyethylene sorbitan monooleate.

The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent. Diluents and solvents that may be employed are, for example, water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile fixed oils are conventionally employed as solvents or suspending media. For this purpose, any bland, fixed oil may be employed including synthetic mono or diglycerides. In addition, fatty acids such as oleic acid may be used in the preparation of injectables.

A composition of the invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions may be prepared by mixing the drug with a suitable non-irritation excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such material are, for example, cocoa butter and polyethylene glycol.

Another formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art (see, e.g., U.S. Patent No. 5,023,252, incorporated herein by reference). Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

It may be desirable or necessary to introduce the pharmaceutical composition to the patient via a mechanical delivery device. The construction and use of mechanical delivery devices for the delivery of pharmaceutical agents is well known in the art. For example, direct techniques for administering a drug directly to the brain usually involve placement of a drug delivery catheter into the patient's ventricular system to bypass the

blood-brain barrier. One such implantable delivery system, used for the transport of agents to specific anatomical regions of the body, is described in U.S. Patent No. 5,011,472, incorporated herein by reference.

The compositions of the invention may also contain other conventional pharmaceutically acceptable compounding ingredients, generally referred to as carriers or diluents, as necessary or desired. Any of the compositions of this invention may be preserved by the addition of an antioxidant such as ascorbic acid or by other suitable preservatives. Conventional procedures for preparing such compositions in appropriate dosage forms can be utilized.

Commonly used pharmaceutical ingredients which may be used as appropriate to formulate the composition for its intended route of administration include: acidifying agents, for example, but are not limited to, acetic acid, citric acid, fumaric acid, hydrochloric acid, nitric acid; and alkalinizing agents such as, but are not limited to, ammonia solution, ammonium carbonate, diethanolamine, monoethanolamine, potassium hydroxide, sodium borate, sodium carbonate, sodium hydroxide, triethanolamine, triolamine.

Other pharmaceutical ingredients include, for example, but are not limited to, adsorbents (e.g., powdered cellulose and activated charcoal); aerosol propellants (e.g., carbon dioxide, CCl_2F_2 , $\text{F}_2\text{ClC-CClF}_2$ and CClF_3); air displacement agents (e.g., nitrogen and argon); antifungal preservatives (e.g., benzoic acid, butylparaben, ethylparaben, methylparaben, propylparaben, sodium benzoate); antimicrobial preservatives (e.g., benzalkonium chloride, benzethonium chloride, benzyl alcohol, cetylpyridinium chloride, chlorobutanol, phenol, phenylethyl alcohol, phenylmercuric nitrate and thimerosal); antioxidants (e.g., ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, hypophosphorus acid, monothioglycerol, propyl gallate, sodium ascorbate, sodium bisulfite, sodium formaldehyde sulfoxylate, sodium metabisulfite); binding materials (e.g., block polymers, natural and synthetic rubber, polyacrylates, polyurethanes, silicones and styrene-butadiene copolymers); buffering agents (e.g., potassium metaphosphate, potassium phosphate monobasic, sodium acetate, sodium citrate anhydrous and sodium citrate dihydrate); carrying agents (e.g., acacia syrup, aromatic syrup, aromatic elixir, cherry syrup, cocoa syrup, orange syrup, syrup, corn oil, mineral oil, peanut oil, sesame oil, bacteriostatic sodium chloride injection and bacteriostatic water for injection); chelating agents (e.g., edetate disodium and edetic acid); colorants (e.g., FD&C Red No. 3, FD&C Red No. 20, FD&C Yellow No. 6, FD&C Blue No. 2, D&C Green No. 5, D&C Orange No. 5, D&C Red No. 8, caramel and ferric oxide red); clarifying agents (e.g., bentonite); emulsifying agents (but are not limited to,

acacia, cetomacrogol, cetyl alcohol, glyceryl monostearate, lecithin, sorbitan monooleate, polyethylene 50 stearate); encapsulating agents (e.g., gelatin and cellulose acetate phthalate); flavorants (e.g., anise oil, cinnamon oil, cocoa, menthol, orange oil, peppermint oil and vanillin); humectants (e.g., glycerin, propylene glycol and sorbitol); levigating agents (e.g., mineral oil and glycerin); oils (e.g., arachis oil, mineral oil, olive oil, peanut oil, sesame oil and vegetable oil); ointment bases (e.g., lanolin, hydrophilic ointment, polyethylene glycol ointment, petrolatum, hydrophilic petrolatum, white ointment, yellow ointment, and rose water ointment); penetration enhancers (transdermal delivery) (e.g., monohydroxy or polyhydroxy alcohols, saturated or unsaturated fatty alcohols, saturated or unsaturated fatty esters, saturated or unsaturated dicarboxylic acids, essential oils, phosphatidyl derivatives, cephalin, terpenes, amides, ethers, ketones and ureas); plasticizers (e.g., diethyl phthalate and glycerin); solvents (e.g., alcohol, corn oil, cottonseed oil, glycerin, isopropyl alcohol, mineral oil, oleic acid, peanut oil, purified water, water for injection, sterile water for injection and sterile water for irrigation); stiffening agents (e.g., cetyl alcohol, cetyl esters wax, microcrystalline wax, paraffin, stearyl alcohol, white wax and yellow wax); suppository bases (e.g., cocoa butter and polyethylene glycols (mixtures)); surfactants (e.g., benzalkonium chloride, nonoxynol 10, octoxynol 9, polysorbate 80, sodium lauryl sulfate and sorbitan monopalmitate); suspending agents (e.g., agar, bentonite, carbomers, carboxymethylcellulose sodium, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, kaolin, methylcellulose, tragacanth and veegum); sweetening e.g., aspartame, dextrose, glycerin, mannitol, propylene glycol, saccharin sodium, sorbitol and sucrose); tablet anti-adherents (e.g., magnesium stearate and talc); tablet binders (e.g., acacia, alginic acid, carboxymethylcellulose sodium, compressible sugar, ethylcellulose, gelatin, liquid glucose, methylcellulose, povidone and pregelatinized starch); tablet and capsule diluents (e.g., dibasic calcium phosphate, kaolin, lactose, mannitol, microcrystalline cellulose, powdered cellulose, precipitated calcium carbonate, sodium carbonate, sodium phosphate, sorbitol and starch); tablet coating agents (e.g., liquid glucose, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose, ethylcellulose, cellulose acetate phthalate and shellac); tablet direct compression excipients (e.g., dibasic calcium phosphate); tablet disintegrants (e.g., alginic acid, carboxymethylcellulose calcium, microcrystalline cellulose, polacrillin potassium, sodium alginate, sodium starch glycollate and starch); tablet glidants (e.g., colloidal silica, corn starch and talc); tablet lubricants (e.g., calcium stearate, magnesium stearate, mineral oil, stearic acid and zinc stearate); tablet/capsule opaquants (e.g., titanium dioxide); tablet

polishing agents (e.g., carnuba wax and white wax); thickening agents (e.g., beeswax, cetyl alcohol and paraffin); tonicity agents (e.g., dextrose and sodium chloride); viscosity increasing agents (e.g., alginic acid, bentonite, carbomers, carboxymethylcellulose sodium, methylcellulose, povidone, sodium alginate and tragacanth); and wetting agents (e.g., heptadecaethylene oxycetanol, lecithins, polyethylene sorbitol monooleate, polyoxyethylene sorbitol monooleate, and polyoxyethylene stearate).

The compounds identified by the methods described herein may be administered as the sole pharmaceutical agent or in combination with one or more other pharmaceutical agents where the combination causes no unacceptable adverse effects. For example, the compounds of this invention can be combined with known anti-obesity, or with known antidiabetic or other indication agents, and the like, as well as with admixtures and combinations thereof.

The compounds identified by the methods described herein may also be utilized, in free base form or in compositions, in research and diagnostics, or as analytical reference standards, and the like. Therefore, the present invention includes compositions which are comprised of an inert carrier and an effective amount of a compound identified by the methods described herein, or a salt or ester thereof. An inert carrier is any material which does not interact with the compound to be carried and which lends support, means of conveyance, bulk, traceable material, and the like to the compound to be carried. An effective amount of compound is that amount which produces a result or exerts an influence on the particular procedure being performed.

Formulations suitable for subcutaneous, intravenous, intramuscular, and the like; suitable pharmaceutical carriers; and techniques for formulation and administration may be prepared by any of the methods well known in the art (see, e.g., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa., 20th edition, 2000)

The following examples are presented to illustrate the invention described herein, but should not be construed as limiting the scope of the invention in any way.

Capsule Formulation

A capsule formula is prepared from:

Compound of this invention	40 mg
Starch	109 mg
Magnesium stearate	1 mg

The components are blended, passed through an appropriate mesh sieve, and filled into hard gelatin capsules.

Tablet Formulation

A tablet is prepared from:

Compound of this invention	25 mg
Cellulose, microcrystalline	200 mg
Colloidal silicon dioxide	10 mg
Stearic acid	5.0 mg

The ingredients are mixed and compressed to form tablets. Appropriate aqueous and non-aqueous coatings may be applied to increase palatability, improve elegance and stability or delay absorption.

Sterile IV Solution

A 5 mg/ml solution of the desired compound of this invention is made using sterile, injectable water, and the pH is adjusted if necessary. The solution is diluted for administration to 1-2 mg/ml with sterile 5% dextrose and is administered as an IV infusion over 60 minutes.

Intramuscular suspension

The following intramuscular suspension is prepared:

Compound of this invention	50 mg/ml
Sodium carboxymethylcellulose	5 mg/ml
TWEEN 80	4 mg/ml
Sodium chloride	9 mg/ml
Benzyl alcohol	9 mg/ml

The suspension is administered intramuscularly.

Hard Shell Capsules

A large number of unit capsules are prepared by filling standard two-piece hard galantine capsules each with 100 mg of powdered active ingredient, 150 mg of lactose, 50 mg of cellulose and 6 mg of magnesium stearate.

Soft Gelatin Capsules

A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil is prepared and injected by means of a positive displacement pump into molten gelatin to form soft gelatin capsules containing 100 mg of the active ingredient. The capsules are washed and dried. The active ingredient can be dissolved in a mixture of polyethylene glycol, glycerin and sorbitol to prepare a water miscible medicine mix.


Immediate Release Tablets/Capsules

These are solid oral dosage forms made by conventional and novel processes. These units are taken orally without water for immediate dissolution and delivery of the medication. The active ingredient is mixed in a liquid containing ingredient such as sugar, gelatin, pectin and sweeteners. These liquids are solidified into solid tablets or caplets by freeze drying and solid state extraction techniques. The drug compounds may be compressed with viscoelastic and thermoelastic sugars and polymers or effervescent components to produce porous matrices intended for immediate release, without the need of water.

It should be apparent to one of ordinary skill in the art that changes and modifications can be made to this invention without departing from the spirit or scope of the invention as it is set forth herein.

We claim:

1. A compound of Formula I,



wherein

R is H or C₁ - C₆ alkyl;

R¹ is H, COOR, C₃-C₈ cycloalkyl, or

C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;

R² is H, halo, or C₁-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro,

or

R² is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl, each of which may be unsubstituted or substituted with R⁶;

R³ is H, C₁-C₆ alkyl, or phenyl which may be unsubstituted or substituted with R⁶;

X is O or S;

R⁴ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl, either of which may be unsubstituted or substituted with fluoro, oxo, or C₁-C₆ alkoxy which may be unsubstituted or substituted with C₁-C₆ alkoxy, or phenyl optionally substituted with R⁶,

or either of which may be substituted with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶,

or

C₁-C₆ alkyl may also be substituted with C₃-C₈ cycloalkyl or with phenoxy which may be unsubstituted or substituted with R⁶ or with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl, each of which may be unsubstituted or substituted with R⁶,

or

R⁴ is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl, each of which may be unsubstituted or substituted with R⁶, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, benzodioxolyl, dihydrobenzofuranyl, indolyl, pyrimidinyl or phenoxy,

each of which may be unsubstituted or substituted with R⁶;

R⁵ is H, halo or C₁-C₆ alkyl optionally substituted with oxo;

R⁶ is halo, CF₃, C₁-C₆ alkyl optionally substituted with oxo or hydroxy, or

C₁-C₆ alkoxy optionally substituted with fluoro;

and pharmaceutically acceptable salts and esters thereof.

2. The compound of claim 1, wherein
- R is H or C₁ - C₆ alkyl;
- R¹ is H, COOR, C₃-C₈ cycloalkyl, or
- C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;
- R² is H, halo, or C₁-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro,
- or
- R² is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl, each of which may be unsubstituted or substituted with R⁶;
- R³ is H, C₁-C₆ alkyl, or phenyl which may be unsubstituted or substituted with R⁶;
- X is O;
- R⁴ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl, either of which may be unsubstituted or substituted with fluoro, oxo, or C₁-C₆ alkoxy which may be unsubstituted or substituted with C₁-C₆ alkoxy, or phenyl optionally substituted with R⁶,
- or either of which may be substituted with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,
- each of which may be unsubstituted or substituted with R⁶,
- or
- C₁-C₆ alkyl may also be substituted with C₃-C₈ cycloalkyl or with phenoxy which may be unsubstituted or substituted with R⁶ or with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl,

benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶,

or

R⁴ is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, benzodioxolyl, dihydrobenzofuranyl, indolyl, pyrimidinyl or phenoxy,

each of which may be unsubstituted or substituted with R⁶;

R⁵ is H, halo or C₁-C₆ alkyl optionally substituted with oxo;

R⁶ is halo, CF₃, C₁-C₆ alkyl optionally substituted with oxo or hydroxy, or

C₁-C₆ alkoxy optionally substituted with fluoro;

and pharmaceutically acceptable salts and esters thereof.

3. The compound of claim 1, wherein

R is H or C₁ - C₆ alkyl;

R¹ is H, COOR, C₃-C₈ cycloalkyl, or

C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;

R² is H, halo, or C₁-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro,

or

R² is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl,

isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl, each of which may be unsubstituted or substituted with R⁶;

R³ is H, C₁-C₆ alkyl, or phenyl which may be unsubstituted or substituted with R⁶;

X is S;

R⁴ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl, either of which may be unsubstituted or substituted with fluoro, oxo, or C₁-C₆ alkoxy which may be unsubstituted or substituted with C₁-C₆ alkoxy, or phenyl optionally substituted with R⁶,

or either of which may be substituted with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶,

or

C₁-C₆ alkyl may also be substituted with C₃-C₈ cycloalkyl or with phenoxy which may be unsubstituted or substituted with R⁶ or with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶,

or

R⁴ is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl,

indoliny, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperaziny, morpholiny, benzodioxolyl, dihydrobenzofuranyl, indolyl, pyrimidinyl or phenoxy,

each of which may be unsubstituted or substituted with R⁶;

R⁵ is H, halo or C₁-C₆ alkyl optionally substituted with oxo;

R⁶ is halo, CF₃, C₁-C₆ alkyl optionally substituted with oxo or hydroxy, or

C₁-C₆ alkoxy optionally substituted with fluoro;

and pharmaceutically acceptable salts and esters thereof.

4. The compound of claim 1, wherein

R is H;

R¹ is H, COOR, C₃-C₈ cycloalkyl, or

C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;

R² is H, halo, or C₁-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro;

R³ is H or C₁-C₆ alkyl, or phenyl which may be unsubstituted or substituted with R⁶;

X is S;

R⁴ is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrroliny, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyraziny, pyridaziny, piperaziny, morpholiny, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indoliny, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl,

pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, benzodioxolyl, dihydrobenzofuranyl, indolyl, pyrimidinyl or phenoxy,

each of which may be unsubstituted or substituted with R⁶;

R⁵ is H, halo or C₁-C₆ alkyl optionally substituted with oxo;

R⁶ is halo, CF₃, C₁-C₆ alkyl optionally substituted with oxo or hydroxy, or

C₁-C₆ alkoxy optionally substituted with fluoro;

and pharmaceutically acceptable salts and esters thereof.

5. The compound of claim 1, wherein

R is H;

R¹ is H, COOR, C₃-C₈ cycloalkyl, or

C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;

R² is H, halo, or C₁-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro;

R³ is H or C₁-C₆ alkyl, or phenyl which may be unsubstituted or substituted with R⁶;

X is O;

R⁴ is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, benzodioxolyl, dihydrobenzofuranyl, indolyl, pyrimidinyl or phenoxy,

each of which may be unsubstituted or substituted with R⁶;

R⁵ is H, halo or C₁-C₆ alkyl optionally substituted with oxo;

R⁶ is halo, CF₃, C₁-C₆ alkyl optionally substituted with oxo or hydroxy, or

C₁-C₆ alkoxy optionally substituted with fluoro;
and pharmaceutically acceptable salts and esters thereof.

6. The compound of claim 1, wherein

R is H;

R¹ is H, COOR, C₃-C₈ cycloalkyl, or

C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;

R² is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl, each of which may be unsubstituted or substituted with R⁶;

R³ is H, C₁-C₆ alkyl, or phenyl which may be unsubstituted or substituted with R⁶;

X is O;

R⁴ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl, either of which may be unsubstituted or substituted with fluoro, oxo, or C₁-C₆ alkoxy which may be unsubstituted or substituted with C₁-C₆ alkoxy, or phenyl optionally substituted with R⁶,

or either of which may be substituted with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶,

or

C₁-C₆ alkyl may also be substituted with C₃-C₈ cycloalkyl or with phenoxy which may be unsubstituted or substituted with R⁶ or with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl,

indoliny, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶,

or

R⁴ is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidiny, pyrroliny, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidiny, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidiny, pyraziny, pyridaziny, piperaziny, morpholiny, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indoliny, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R⁶, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidiny, piperidiny, tetrahydropyranyl, tetrahydrothiopyranyl, piperaziny, morpholiny, benzodioxolyl, dihydrobenzofuranyl, indolyl, pyrimidiny or phenoxy,

each of which may be unsubstituted or substituted with R⁶;

R⁵ is H, halo or C₁-C₆ alkyl optionally substituted with oxo;

R⁶ is halo, CF₃, C₁-C₆ alkyl optionally substituted with oxo or hydroxy, or

C₁-C₆ alkoxy optionally substituted with fluoro;

and pharmaceutically acceptable salts and esters thereof.

7. The compound of claim 1, wherein C-1' is in the S configuration or the R configuration.

8. The compound of claim 1, wherein the configurations of C-1' and C-2 are selected from the group consisting of C-1' is R and C-2 is R; C-1' is R and C-2 is S; C-1' is S and C-2 is S; and C-1' is S and C-2 is R.

9. The compound of claim 1 selected from the group consisting of:

5-{2-[2-(4-ethylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;

5-(2-{5-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl}ethoxy)-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[5-methyl-2-(2-naphthyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(4-*tert*-butylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(4-butylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(3,4-dimethylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(1-benzothien-2-yl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(4-isopropylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-[2-(2-cyclopentyl-5-methyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(3-fluoro-4-methylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid; and
5-{2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid.

10. The compound of claim 9, wherein C-1' is in the S configuration or the R configuration.

11. The compound of claim 1 selected from the group consisting of:

2-(5-{2-[2-(4-ethylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)propanoic acid;
2-(5-{2-[2-(3,4-dichlorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)propanoic acid;
2-(5-{2-[5-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)propanoic acid;
2-(5-{2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)propanoic acid;
2-(5-{2-[2-(4-ethylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)propanoic acid;

2-(5-{2-[2-(3,4-dimethylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)propanoic acid; and
2-(5-{2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)propanoic acid.

12. The compound of claim 11, wherein the configurations of C-1' and C-2 are selected from the group consisting of C-1' is R and C-2 is R; C-1' is R and C-2 is S; C-1' is S and C-2 is S; and C-1' is S and C-2 is R.

13. The compound of claim 1 selected from the group consisting of:

5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-[2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(3-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(1,3-benzodioxol-5-yl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(3-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(4-fluoro-3-methylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(3,4-dichlorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(1,1'-biphenyl-4-yl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(1,3-benzodioxol-5-yl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(4-methoxyphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-(2-{5-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-4-yl}ethoxy)-2,3-dihydro-1*H*-inden-1-yl acetic acid;

5-{2-[2-(4-cyanophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(4-isopropylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(3-chlorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic
acid;

5-{2-[2-(4-cyanophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic
acid;

5-{2-[2-(3-cyanophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic
acid;

5-{2-[2-(4-chlorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-(2-{5-methyl-2-[3-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl]ethoxy)-2,3-dihydro-1*H*-inden-
1-yl acetic acid;

5-{2-[2-(3-chloro-4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-
1-yl acetic acid;

5-{2-[2-(3,4-dichlorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(3,4-dimethylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(4-acetylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(3-amino-4-methylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-
1-yl acetic acid trifluoroacetate;

5-{2-[2-(2-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(4-chlorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(4-ethoxyphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-{2-[2-(3,4-dimethoxyphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-
yl acetic acid;

5-{2-[5-methyl-2-(3-methylphenyl)-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl
acetic acid;

5-(2-{5-methyl-2-[3-(trifluoromethyl)phenyl]-1,3-thiazol-4-yl}ethoxy)-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(3-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(3,5-dimethylphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-(2-{5-methyl-2-[4-(trifluoromethoxy)phenyl]-1,3-thiazol-4-yl}ethoxy)-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(3-methoxyphenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[2-(1,1'-biphenyl-4-yl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[5-ethyl-2-(4-ethylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[5-ethyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-{2-[5-ethyl-2-(4-methoxyphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-(2-{2-[4'-(hydroxymethyl)-1,1'-biphenyl-4-yl]-5-methyl-1,3-oxazol-4-yl}ethoxy)-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-(2-{2-[4-(5-acetyl-2-thienyl)phenyl]-5-methyl-1,3-oxazol-4-yl}ethoxy)-2,3-dihydro-1*H*-inden-1-yl acetic acid;
5-(2-{2-[3-(1*H*-indol-5-yl)phenyl]-5-methyl-1,3-oxazol-4-yl}ethoxy)-2,3-dihydro-1*H*-inden-1-yl acetic acid;
(5-{2-[2-(3-fluoro-4-methylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)acetic acid; and
(-5-{2-[5-methyl-2-(3-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)acetic acid.

14. The compound of claim 13, wherein C-1' is in the S configuration or the R configuration.

15. The compound of claim 1 selected from the group consisting of:
2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl}butanoic acid;

3-ethoxy-2-{5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl}-3-oxopropanoic acid;

2-(5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)propanoic acid;

2-(5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

4-methyl-2-(5-{2-[5-methyl-2-(4-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)pentanoic acid;

2-{5-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl}butanoic acid;

2-(5-{2-[2-(4-chlorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[5-methyl-2-(3-methylphenyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-(3,4-dichlorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[5-methyl-2-(2-naphthyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[5-methyl-2-(1-naphthyl)-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-(3-chlorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-cyclopentyl-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-cyclohexyl-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-(3-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-(4-ethylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-{2-[2-(1-benzothien-2-yl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-[5-(2-[5-methyl-2-[3-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl]ethoxy)-2,3-dihydro-1*H*-inden-1-yl]butanoic acid;

2-(5-[2-[2-(4-*tert*-butylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-[2-[2-(3-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-[2-[2-(4-isopropylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-[2-[2-(1,3-benzodioxol-5-yl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-[6-[2-(2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl]butanoic acid;

2-(5-[2-[2-(3,4-difluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-[5-(2-[5-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl]ethoxy)-2,3-dihydro-1*H*-inden-1-yl]butanoic acid;

2-(5-[2-[5-methyl-2-(phenoxymethyl)-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-[2-[2-(4-fluoro-3-methylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-[2-[2-(3-fluoro-4-methylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-[2-[2-(4-butylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-[2-[2-(3,4-dimethylphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-(5-[2-[5-methyl-2-(4-chlorophenoxymethyl)-1,3-oxazol-4-yl]ethoxy]-2,3-dihydro-1*H*-inden-1-yl)butanoic acid;

2-[6-chloro-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl]butanoic acid;

2-[5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-6-phenyl-2,3-dihydro-1*H*-inden-1-yl]butanoic acid;


2-[6-(4-chlorophenyl)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl]butanoic acid;

2-[6-(4-methoxyphenyl)-5-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]-2,3-dihydro-1*H*-inden-1-yl]butanoic acid; and

2-(5-{2-[2-(2,3-dihydro-1-benzofuran-6-yl)-5-methyl-1,3-oxazol-4-yl]ethoxy}-2,3-dihydro-1*H*-inden-1-yl)butanoic acid.

16. The compound of claim 15, wherein the configurations of C-1' and C-2 are selected from the group consisting of C-1' is R and C-2 is R; C-1' is R and C-2 is S; C-1' is S and C-2 is S; and C-1' is S and C-2 is R.

17. A compound of Formula II,



(II)

wherein

R is H or C₁ - C₆ alkyl;

R¹ is H, COOR, C₃-C₈ cycloalkyl, or

C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;

R² is H, halo, or C₁-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro,

or

R² is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl, each of which may be unsubstituted or substituted with R⁶;

R³ is H, C₁-C₆ alkyl, or phenyl which may be unsubstituted or substituted with R⁶;

R⁴ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl, either of which may be unsubstituted or substituted with fluoro, oxo, or C₁-C₆ alkoxy which may be unsubstituted or substituted with C₁-C₆ alkoxy, or phenyl optionally substituted with R⁶,

or either of which may be substituted with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl,

pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R^6 ,

or

C_1 - C_6 alkyl may also be substituted with C_3 - C_8 cycloalkyl or with phenoxy which may be unsubstituted or substituted with R^6 or with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R^6 ,

or

R^4 is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R^6 , or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, benzodioxolyl, dihydrobenzofuranyl, indolyl, pyrimidinyl or phenoxy,


each of which may be unsubstituted or substituted with R^6 ;

R^5 is halo or C_1 - C_6 alkyl optionally substituted with oxo;

R^6 is halo, CF_3 , C_1 - C_6 alkyl optionally substituted with oxo or hydroxy, or

C₁-C₆ alkoxy optionally substituted with fluoro;


R⁷ is H, C₁-C₆ alkyl optionally substituted with phenyl or oxo, C₁-C₆ trialkylsilyl, arylalkylsilyl, COR⁸, COOR⁸, or



X is O or S;

R⁸ is C₁-C₆ alkyl, or phenyl optionally substituted with C₁-C₆ alkyl, halo, or nitro; and the salts thereof.

18. A compound of Formula III,



wherein

R is H or C₁ - C₆ alkyl;

R¹ is H, COOR, C₃-C₈ cycloalkyl, or

C₁ - C₆ alkyl, C₂-C₆ alkenyl, or C₁-C₆ alkoxy each of which may be unsubstituted or substituted with fluoro, methylenedioxyphenyl, or phenyl which may be unsubstituted or substituted with R⁶;

R² is H, halo, or C₁-C₆ alkyl which may be unsubstituted or substituted with C₁-C₆ alkoxy, oxo, fluoro,

or

R² is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl, each of which may be unsubstituted or substituted with R⁶;

R^3 is H, C_1 - C_6 alkyl, or phenyl which may be unsubstituted or substituted with R^6 ;

R^5 is halo or C_1 - C_6 alkyl optionally substituted with oxo;

R^6 is halo, CF_3 , C_1 - C_6 alkyl optionally substituted with oxo or hydroxy, or


C_1 - C_6 alkoxy optionally substituted with fluoro;

R^7 is H, C_1 - C_6 alkyl optionally substituted with phenyl or oxo, C_1 - C_6 trialkylsilyl,


arylalkylsilyl, or COR^8 , $COOR^8$;

R^8 is C_1 - C_6 alkyl, or phenyl optionally substituted with C_1 - C_6 alkyl, halo, or nitro; and the salts thereof.

19. A process for the preparation of a compound of Formula V



comprising the step of a stereospecific hydrogenation of a compound of Formula IV



in the presence of hydrogen source and a catalyst,
wherein

R^9 is methoxy optionally substituted by fluoro,

C_2 - C_6 alkoxy, C_1 - C_6 alkyl, or C_4 - C_8 cycloalkyl each optionally substituted by fluoro, methylenedioxyphenyl or phenyl optionally substituted with R^{13} ;

R^{10} is hydrogen, fluoro, methyl optionally substituted with fluoro, oxo, or

C_2 - C_6 alkyl which may be unsubstituted or substituted with C_1 - C_6 alkoxy, oxo, fluoro, or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl,


each of which may be unsubstituted or substituted with R^{13} ,

or

R^{10} is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, or morpholinyl, each of which may be unsubstituted or substituted with R^{13} ;

R^{11} is halo or C_1 - C_6 alkyl optionally substituted with oxo;

R^{12} is hydrogen, methyl optionally substituted with fluoro or oxo, C_2 - C_6 alkyl optionally substituted with phenyl, fluoro, or oxo, C_1 - C_6 trialkylsilyl, arylalkylsilyl, COR^{14} , $COOR^{14}$, or



R^{13} is fluoro, CF_3 , C_1 - C_6 alkyl optionally substituted with hydroxy or oxo, or C_1 - C_6 alkoxy optionally substituted with fluoro;

R^{14} is C_1 - C_6 alkyl, or phenyl optionally substituted with C_1 - C_6 alkyl or fluoro;

R^{15} is hydrogen, C_1 - C_6 alkyl or phenyl substituted with R^{13} ;

R^{16} is methyl optionally substituted with fluoro, oxo or

with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazolinyl, quinoxazolinyl, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R^{13} , or

C_4 - C_8 cycloalkyl or C_2 - C_6 alkyl, either of which may be unsubstituted or substituted with fluoro, methoxy, C_2 - C_6 alkoxy optionally substituted with phenyl or C_1 - C_6 alkoxy, oxo or with, phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, piperazinyl, morpholinyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl,

isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl,
dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R^{13} , or

C_2-C_6 alkyl which may also be substituted with C_4-C_8 cycloalkyl or with phenoxy which may be unsubstituted or substituted with R^6 or with phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidiny, pyrroliny, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidiny, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidiny, pyraziny, pyridaziny, piperaziny, morpholiny, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indoliny, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R^{13} ,

or

R^{16} is phenyl, naphthyl, furyl, thienyl, pyrrolyl, tetrahydrofuryl, pyrrolidiny, pyrroliny, tetrahydrothienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, piperidiny, tetrahydropyranyl, tetrahydrothiopyranyl, pyrimidiny, pyraziny, pyridaziny, piperaziny, morpholiny, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indoliny, indazolyl, benzoxazolyl, benxothiazolyl, benzimidazolyl, benzisoxazolyl, benzisothiazolyl, benzodioxolyl, quinolyl, isoquinolyl, quinazoliny, quinoxazoliny, dihydrobenzopyranyl, dihydrobenzothiopyranyl, or 1,4-benzodioxanyl,

each of which may be unsubstituted or substituted with R^{13} , or with phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrrolidiny, piperidiny, tetrahydropyranyl, tetrahydrothiopyranyl, piperaziny, morpholiny, pyrimidiny, indolyl, or phenoxy

each of which may be unsubstituted or substituted with R^{13} , and

X is O or S.

20. The process of claim 19, wherein hydrogenation is performed in the presence of a base.

21. The process of claim 19, wherein said compound of Formula V is in the syn form.

22. The process of claim 19, wherein said compound of Formula IV is a racemic mixture.
23. The process of claim 19, wherein said compound of Formula IV is enantiomerically enriched.
24. The process of claim 23, wherein said compound of Formula V is at least about 40% de.
25. The process of claim 19, wherein said catalyst is transition metal based homogenous catalyst.
26. The process of claim 19, wherein said catalyst is selected from the group consisting of $\text{ClRh[P(Ph)}_3\text{]}_3$ (Wilkinson's catalyst), (1,5-cyclooctadiene)tricyclohexylphosphinepyridinoiridium(I) hexafluorophosphate, (1,5-cyclooctadiene)bis(methyldiphenylphosphine)iridium(I) hexafluorophosphate, (Crabtree's catalysts).
27. The process of claim 20, wherein said base is selected from the group consisting of mono($\text{C}_1\text{-C}_6$ alkyl)amines, di($\text{C}_1\text{-C}_6$ alkyl)amines, tri($\text{C}_1\text{-C}_6$ alkyl)amines, tertiary amines, inorganic bases, and optically active bases.
28. The process of claim 19, wherein said hydrogen source is hydrogen gas.
29. The process of claim 28, wherein said hydrogenation is performed under a hydrogen pressure from about atmospheric pressure to about 1,000 psi.
30. The process of claim 29, wherein said hydrogenation is performed under a hydrogen pressure from about 20 psi to about 100 psi.
31. A process for the preparation of a compound of Formula V, comprising the steps of:
forming diastereomeric salts of a compound of Formula IV by treatment with
a resolving agent;
separating said diastereomeric salts by crystallization with a crystallization
solvent; and

reducing the separated diastereomeric salts by hydrogenation in the presence of catalyst and a solvent,
wherein
the compounds of Formula V and Formula IV are defined as in claim 19.

32. The process of claim 31, further comprising the step of;
liberating the antipodes from the separated diastereomeric salts by treatment with an aqueous mineral acid.
33. The process of claim 31, wherein said reduction step is performed in the presence of a base.
34. The process of claim 31 or 32, wherein said resolving agent is an optically active base selected from the group consisting of quinine, cinchonine, (+)-alpha-methylbenzylamine, and (-)-alpha-methylbenzylamine.
35. The process of claim 31 or 32, wherein said crystallization solvent is selected from the group consisting of acetonitrile, acetone, *t*-butanol, 2-propanol, ethanol, methanol, and mixtures thereof.
36. A process for the preparation of a compound of Formula V, comprising the steps of:
reducing a compound of Formula IV to a compound of Formula V
by hydrogenation in the presence of a hydrogenation catalyst and
a solvent;
forming diastereomeric salts of the compound of Formula V by treatment with
a resolving agent;
separating the diastereomeric salts by crystallization in a crystallization
solvent; and
liberating the individual antipodes from the separated salts by treatment with
aqueous mineral acid,
wherein
the compounds of Formula V and Formula IV are defined as in claim 19.
37. The process of claim 36, wherein said reduction step is performed in the presence of a base.

38. The process of claim 36 or 37, wherein said hydrogenation catalyst is selected from the group consisting of $\text{ClRh}[\text{P}(\text{Ph})_3]_3$ (Wilkinson's catalyst), (1,5-cyclooctadiene)tricyclohexylphosphinepyridinoiridium(I) hexafluorophosphate, and (1,5-cyclooctadiene)bis(methyldiphenylphosphine)iridium(I) hexafluorophosphate (Crabtree's catalyst).
39. The process of claim 36 or 37, wherein said resolving agent is selected from the group consisting of quinine, cinchonine, (+)-alpha-methylbenzylamine, and (-)-alpha-methylbenzylamine.
40. The process of claim 36 or 37, wherein said crystallization solvent is selected from the group consisting of acetonitrile, acetone, *t*-butanol, 2-propanol, ethanol, methanol, and mixtures thereof.
41. A method of identifying compounds useful for the treatment of diabetes, diabetes-related disorders, obesity, and atherosclerotic disease comprising the step of determining the insulin sensitizing activity of said compound.
42. The method of claim 41 further comprising the steps of:
growing cells to 2-4 days post-confluency;
treating cells with differentiation media; and
assaying for insulin receptor binding activity.
43. The method of claim 42, wherein said differentiation media comprises Insulin-like Growth Factor (IGF-1) and test compounds.
44. The method of claim 43, wherein said Insulin-like Growth Factor is present in an amount from about 0.1 μM to about 1 μM .
45. The method of claim 42, wherein said cells are 3T3-L1 cells.
46. A pharmaceutical composition comprising an effective amount of a compound of claim 1, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
47. A pharmaceutical composition comprising an effective amount of a compound of

- claim 7 or 8, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
48. A pharmaceutical composition comprising an effective amount of a compound of claim 9, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
49. A pharmaceutical composition comprising an effective amount of a compound of claim 10, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
50. A pharmaceutical composition comprising an effective amount of a compound of claim 11, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
51. A pharmaceutical composition comprising an effective amount of a compound of claim 12, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
52. A pharmaceutical composition comprising an effective amount of a compound of claim 13, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
53. A pharmaceutical composition comprising an effective amount of a compound of claim 14, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
54. A pharmaceutical composition comprising an effective amount of a compound of claim 15, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
55. A pharmaceutical composition comprising an effective amount of a compound of claim 16, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier.
56. A pharmaceutical composition comprising an effective amount of a compound of claim 1, or a pharmaceutically acceptable salt or ester thereof, in combination with

a pharmaceutically acceptable carrier and one or more hypoglycemic agents.

57. The pharmaceutical composition of claim 56, wherein said hypoglycemic agent is selected from the group consisting of insulin, biguanidines, sulfonylureas, insulin secretagogues, α -glycosidase inhibitors, and β_3 -adrenoreceptor agonists.
58. A pharmaceutical composition comprising an effective amount of a compound of claim 9, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier and one or more hypoglycemic agents.
59. The pharmaceutical composition of claim 58, wherein said hypoglycemic agent is selected from the group consisting of insulin, biguanidines, sulfonylureas, insulin secretagogues, α -glycosidase inhibitors, and β_3 -adrenoreceptor agonists.
60. A pharmaceutical composition comprising an effective amount of a compound of claim 11, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier and one or more hypoglycemic agents.
61. The pharmaceutical composition of claim 60, wherein said hypoglycemic agent is selected from the group consisting of insulin, biguanidines, sulfonylureas, insulin secretagogues, α -glycosidase inhibitors, and β_3 -adrenoreceptor agonists.
62. A pharmaceutical composition comprising an effective amount of a compound of claim 13, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier and one or more hypoglycemic agents.
63. The pharmaceutical composition of claim 62, wherein said hypoglycemic agent is selected from the group consisting of insulin, biguanidines, sulfonylureas, insulin secretagogues, α -glycosidase inhibitors, and β_3 -adrenoreceptor agonists.
64. A pharmaceutical composition comprising an effective amount of a compound of claim 15, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier and one or more hypoglycemic agents.
65. The pharmaceutical composition of claim 64, wherein said hypoglycemic agent is selected from the group consisting of insulin, biguanidines, sulfonylureas, insulin

secretagogues, α -glycosidase inhibitors, and β_3 -adrenoreceptor agonists.

66. A pharmaceutical composition comprising an effective amount of a compound of claim 1, or a pharmaceutically acceptable salt or ester thereof, in combination with a pharmaceutically acceptable carrier and one or more agents selected from the group consisting of HMG CoA reductase inhibitor, bile acid binding agent, fibric acid derivative, agent that regulates hypertension, and agent that regulates body weight.
67. A composition comprising an effective amount of a compound of claim 1, or a salt or ester thereof, in combination with an inert carrier.
68. A composition comprising an effective amount of a compound of claim 9, or a salt or ester thereof, in combination with an inert carrier.
69. A composition comprising an effective amount of a compound of claim 11, or a salt or ester thereof, in combination with an inert carrier.
70. A composition comprising an effective amount of a compound of claim 13, or a salt or ester thereof, in combination with an inert carrier.
71. A composition comprising an effective amount of a compound of claim 15, or a salt or ester thereof, in combination with an inert carrier.
72. A method of treating diabetes comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
73. A method of treating Syndrome X comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
74. A method of treating diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
75. The method of claim 74, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose

tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.

76. A method of treating obesity comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
77. A method of treating cardiovascular disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
78. The method of claim 77, wherein said cardiovascular disease is selected from the group consisting of atherosclerotic disease, dyslipidemia, hypercholesterolemia, decreased HDL levels, hypertension, coronary heart disease, and coronary artery disease.
79. A method of treating cerebrovascular disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
80. A method of treating peripheral vessel disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
81. A method of treating lupus comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
82. A method of treating polycystic ovary disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
83. A method of treating carcinogenesis and hyperplasia comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1.
84. A method of treating diabetes comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1 in combination with one or more hypoglycemic agents.

85. A method of treating Syndrome X comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1 in combination with one or more hypoglycemic agents.
86. A method of treating diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1 in combination with one or more hypoglycemic agents.
87. The method of claim 86, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.
88. The method of any one of claims 84 to 87, wherein the compound of Formula I and the one or more hypoglycemic agents are administered as a single pharmaceutical dosage formulation.
89. A method of treating diabetes, Syndrome X, or diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 1 in combination with one or more agents selected from the group consisting of HMG CoA reductase inhibitor, bile acid binding agent, fibric acid derivative, agent that regulates hypertension, and agent that regulates body weight.
90. The method of claim 89, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.
91. A method of treating diabetes comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9.
92. A method of treating Syndrome X comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.

93. A method of treating diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.
94. The method of claim 93, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.
95. A method of treating obesity comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.
96. A method of treating cardiovascular disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.
97. The method of claim 96, wherein said cardiovascular disease is selected from the group consisting of atherosclerotic disease, dyslipidemia, hypercholesterolemia, decreased HDL levels, hypertension, coronary heart disease, and coronary artery disease.
98. A method of treating cerebrovascular disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.
99. A method of treating peripheral vessel disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.
100. A method of treating lupus comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.
101. A method of treating polycystic ovary disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.

102. A method of treating carcinogenesis and hyperplasia comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11.
103. A method of treating diabetes comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11 in combination with one or more hypoglycemic agents.
104. A method of treating Syndrome X comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11 in combination with one or more hypoglycemic agents.
105. A method of treating diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 9 or 11 in combination with one or more hypoglycemic agents.
106. The method of claim 105, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.
107. A method of treating diabetes comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
108. A method of treating Syndrome X comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
109. A method of treating diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
110. The method of claim 109, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.

111. A method of treating obesity comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
112. A method of treating cardiovascular disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
113. The method of claim 112, wherein said cardiovascular disease is selected from the group consisting of atherosclerotic disease, dyslipidemia, hypercholesterolemia, decreased HDL levels, hypertension, coronary heart disease, and coronary artery disease.
114. A method of treating cerebrovascular disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
115. A method of treating peripheral vessel disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
116. A method of treating lupus comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
117. A method of treating polycystic ovary disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
118. A method of treating carcinogenesis and hyperplasia comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13.
119. A method of treating diabetes comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13 in combination with one or more hypoglycemic agents.

120. A method of treating Syndrome X comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13 in combination with one or more hypoglycemic agents.
121. A method of treating diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 13 in combination with one or more hypoglycemic agents.
122. The method of claim 121, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.
123. A method of treating diabetes comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
124. A method of treating Syndrome X comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
125. A method of treating diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
126. The method of claim 125, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.
127. A method of treating obesity comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
128. A method of treating cardiovascular disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
129. The method of claim 128, wherein said cardiovascular disease is selected from the group consisting of atherosclerotic disease, dyslipidemia,

hypercholesterolemia, decreased HDL levels, hypertension, coronary heart disease, and coronary artery disease.

130. A method of treating cerebrovascular disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
131. A method of treating peripheral vessel disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
132. A method of treating lupus comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
133. A method of treating polycystic ovary disease comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
134. A method of treating carcinogenesis and hyperplasia comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15.
135. A method of treating diabetes comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15 in combination with one or more hypoglycemic agents.
136. A method of treating Syndrome X comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15 in combination with one or more hypoglycemic agents.
137. A method of treating diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound of claim 15 in combination with one or more hypoglycemic agents.
138. The method of claim 137, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose

tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.

139. A method of treating diabetes, Syndrome X, or diabetes-related disorders comprising the step of administering to a patient in need thereof a pharmaceutically effective amount of a compound which was identified by the method of claim 41.
140. The method of claim 139, wherein said diabetes-related disorder is selected from the group consisting of hyperglycemia, hyperinsulinemia, impaired glucose tolerance, impaired fasting glucose, dyslipidemia, hypertriglyceridemia, and insulin resistance.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/23614

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D263/32 A61K31/421 C07C69/734 C07D413/04 C07D277/24
C07D417/04 C07D413/10 A61P3/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 16120 A (DOMINIANNI SAMUEL J ;MATTHEWS DONALD P (US); MICHELLYS PIERRE YVES) 8 March 2001 (2001-03-08) page 2, line 28 -page 5, line 11; examples 26,27	1-140
A	EP 0 490 772 A (ADIR) 17 June 1992 (1992-06-17) Page 3, formula (IV)	17-19
	--- -/-- ---	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

7 October 2002

Date of mailing of the international search report

15/10/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Uselli, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/23614

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SUBRAMANIAM S: "THE EMERGING ROLE OF THIAZOLIDINEDIONES IN THE TREATMENT OF DIABETES-MELLITUS AND RELATED DISORDERS" CLINICAL AND EXPERIMENTAL HYPERTENSION, MARCEL DEKKER, NEW YORK, NY, US, vol. 21, no. 1/2, 1999, pages 121-136, XP001028567 ISSN: 1064-1963 figure 1	1-140
P,A	WO 02 18355 A (BROOKS DAWN ALISA ;CONNOR SCOTT EUGENE (US); DOMINIANNI SAMUEL JAM) 7 March 2002 (2002-03-07) page 4, line 1 -page 5, line 23	1-140

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/23614

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 72-140 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/23614

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0116120	A	08-03-2001	AU 7073400 A	26-03-2001
			EP 1206457 A1	22-05-2002
			WO 0116120 A1	08-03-2001
			US 6417212 B1	09-07-2002
EP 0490772	A	17-06-1992	FR 2670491 A1	19-06-1992
			AT 125537 T	15-08-1995
			AU 638368 B2	24-06-1993
			AU 8976291 A	18-06-1992
			CA 2057578 A1	15-06-1992
			DE 69111582 D1	31-08-1995
			DE 69111582 T2	04-04-1996
			DK 490772 T3	11-12-1995
			EP 0490772 A1	17-06-1992
			ES 2077199 T3	16-11-1995
			GR 3017596 T3	31-01-1996
			HK 56497 A	09-05-1997
			IE 914339 A1	17-06-1992
			JP 2009368 C	11-01-1996
			JP 6025217 A	01-02-1994
			JP 7035377 B	19-04-1995
			NZ 240975 A	26-08-1993
			PT 99794 A , B	30-04-1993
			US 5194437 A	16-03-1993
			ZA 9109845 A	30-09-1992
WO 0218355	A	07-03-2002	AU 8465801 A	13-03-2002
			WO 0218355 A1	07-03-2002

This Page Blank (uspto)